Lynn S. Bickley

BATES'

Pocket Guide to

Physical Examination

AND HISTORY TAKING

SIXTH EDITION

BATES'

Pocket Guide to Physical Examination

AND HISTORY TAKING

BATES'

Pocket Guide to Physical Examination

AND HISTORY TAKING

SIXTH EDITION

Lynn S. Bickley, MD

Associate Dean for Curriculum and Professor of Internal Medicine
School of Medicine
Texas Tech University Health Sciences Center
Lubbock, Texas

Peter G. Szilagyi, MD, MPH

Professor of Pediatrics
Chief, Division of General Pediatrics
University of Rochester School of Medicine and Dentistry
Rochester, New York

Acquisitions Editor: Peter Darcy
Development Editor: Renee Gagliardi
Senior Production Editor: Sandra Cherrey Scheinin
Director of Nursing Production: Helen Ewan
Senior Managing Editor/Production: Erika Kors
Design Coordinator: Joan Wendt
Art Director, Illustration: Brett MacNaughton
Manufacturing Coordinator: Karin Duffield
Inducer: Gaye Tarallo
Compositor: Circle Graphics

6th Edition

Copyright © 2009 Wolters Kluwer Health | Lippincott Williams & Wilkins.

Copyright © 2007, 2004, 2000 by Lippincott Williams & Wilkins. Copyright © 1995, 1991 by J. B. Lippincott Company. All rights reserved. This book is protected by copyright. No part of this book may be reproduced or transmitted in any form or by any means, including as photocopies or scanned-in or other electronic copies, or utilized by any information storage and retrieval system without written permission from the copyright owner, except for brief quotations embodied in critical articles and reviews. Materials appearing in this book prepared by individuals as part of their official duties as U.S. government employees are not covered by the abovementioned copyright. To request permission, please contact Lippincott Williams & Wilkins at 530 Walnut Street, Philadelphia PA 19106, via email at permissions@lww.com or via website at lww.com (products and services).

9 8 7 6 5 4 3 2 1

ISBN: 978-0-7817-8066-7 Printed in China

Care has been taken to confirm the accuracy of the information presented and to describe generally accepted practices. However, the authors, editors, and publisher are not responsible for errors or omissions or for any consequences from application of the information in this book and make no warranty, expressed or implied, with respect to the currency, completeness, or accuracy of the contents of the publication. Application of this information in a particular situation remains the professional responsibility of the practitioner; the clinical treatments described and recommended may not be considered absolute and universal recommendations.

The authors, editors, and publisher have exerted every effort to ensure that drug selection and dosage set forth in this text are in accordance with the current recommendations and practice at the time of publication. However, in view of ongoing research, changes in government regulations, and the constant flow of information relating to drug therapy and drug reactions, the reader is urged to check the package insert for each drug for any change in indications and dosage and for added warnings and precautions. This is particularly important when the recommended agent is a new or infrequently employed drug.

Some drugs and medical devices presented in this publication have Food and Drug Administration (FDA) clearance for limited use in restricted research settings. It is the responsibility of the health care provider to ascertain the FDA status of each drug or device

planned for use in his or her clinical practice.



To Robert A. Hoekelman, master pediatrician, whose legacy of blending science with humanism for faculty, students, and patients lives on in this book, which he helped pioneer.

CONTENTS

CHAPTER 1	Overview: Physical Examination and History Taking 1
CHAPIER 2	Clinical Reasoning, Assessment,
CHAITER Z	and Recording Your Findings 17
CHAPTER 3	Interviewing and the Health History 37
CHAPTER 4	Beginning the Physical Examination: General Survey, Vital Signs, and Pain 55
CHAPTER 5	Behavior and Mental Status 75
CHAPTER 6	The Skin, Hair, and Nails 93
CHAPTER 7	The Head and Neck 111
CHAPTER 8	The Thorax and Lungs 141
CHAPTER 9	The Cardiovascular System 163
CHAPTER 10	The Breasts and Axillae 181
CHAPTER 11	The Abdomen 193
CHAPTER 12	The Peripheral Vascular System 215
CHAPTER 13	Male Genitalia and Hernias 229
CHAPTER 14	Female Genitalia 241
CHAPTER 15	The Anus, Rectum, and Prostate 259
CHAPTER 16	The Musculoskeletal System 271
CHAPTER 17	The Nervous System 311
CHAPTER 18	Assessing Children: Infancy Through Adolescence 349
CHAPTER 19	The Pregnant Woman 389
CHAPTER 20	The Older Adult 403
Index 427	

INTRODUCTION

The Pocket Guide to Physical Examination and History Taking, 6/E is a concise, portable text that:

- Describes how to interview the patient and take the health history
- Provides an illustrated review of the physical examination
- Reminds students of common, typical findings
- Describes special techniques of assessment that students may need in specific instances
- Provides succinct aids to interpretation of selected findings

There are several ways to use the Pocket Guide:

To review and remember the content of a health history. To review and rehearse the techniques of examination. This can be done while learning a single section and again while combining the approaches to several body systems or regions into an integrated examination (see Chap. 1).

To review common variations of normal and selected abnormalities. Observations are keener and more precise when the examiner knows what to look, listen, and feel for.

To look up special techniques as the need arises. Maneuvers such as doing an Allen test are included in the relevant sections of the examination and highlighted by a shaded blue-green bar.

To look up additional information about possible findings, including abnormalities and standards of normal.

The *Pocket Guide* is not intended to serve as a primary text for learning the skills of taking a history or performing a physical examination. Its detail is insufficient for these purposes. It is intended instead as an aid for student review and recall and as a convenient, brief, and portable reference.

Overview: Physical Examination and History Taking

This chapter provides a road map to clinical proficiency in two critical areas: the health history and the physical examination.

For adults, the comprehensive history includes *Identifying Data and Source of the History, Chief Complaint(s), Present Illness, Past History, Family History, Personal and Social History,* and *Review of Systems.* New patients in the office or hospital merit a *comprehensive health history;* however, in many situations a more flexible *focused,* or *problem-oriented, interview* is appropriate. The components of the comprehensive health history structure the patient's story and the format of your written record, but the order shown on pp. 3–7 should not dictate the sequence of the interview. The interview is more fluid and should follow the patient's leads and cues, as described in Chapter 3.

The History and Physical Examination: Comprehensive or Focused?

Comprehensive Assessment

• Is appropriate for new patients in the office or hospital

- Provides fundamental and personalized knowledge about the patient
- Strengthens the clinicianpatient relationship
- Helps identify or rule out physical causes related to patient concerns
- Develops proficiency in the essential skills of physical examination

Focused Assessment

- Is appropriate for established patients, especially during routine or urgent care visits
- Addresses focused concerns or symptoms
- Assesses the specific body systems relevant to the patient's concerns

2 Overview: Physical Examination and History Taking

Be sure to distinguish subjective from objective data.

Subjective Data	Objective Data
What the patient tells you	What you detect during the examination
The history, from Chief Complaint through Review of Systems	All physical examination findings

COMPONENTS OF THE ADULT HEALTH HISTORY

Identifying Data

- Identifying data—such as age, gender, occupation, marital status
- Source of the history—usually the patient, but can be a family member or friend, letter of referral, or the medical record
- If appropriate, establish source of referral, because a written report may be needed

Reliability

Varies according to the patient's memory, trust, and mood

Chief Complaint(s)

 The one or more symptoms or concerns causing the patient to seek care

Present Illness

- Amplifies the Chief Complaint, describes how each symptom developed, gives the seven features of every symptom (see p. 3)
- Includes patient's thoughts and feelings about the illness
- Includes pertinent positives and negatives based on relevant portions of the Review of Systems (see pp. 5–7)
- May include medications, allergies, habits of smoking and alcohol, which are frequently pertinent to the present illness

Past History

- Lists childhood illnesses
- Lists adult illnesses with dates for at least four categories: medical; surgical; obstetric/gynecologic; and psychiatric

COMPONENTS OF THE ADULT HEALTH HISTORY (CONTINUED)

 Includes Health Maintenance practices such as immunizations, screening tests, lifestyle issues, and home safety

Family History

- Outlines or diagrams age and health, or age and cause of death, of siblings, parents, and grandparents.
- Documents presence or absence of specific illnesses in family, such as hypertension, coronary artery disease, etc.

Personal and Social History

 Described educational level, family of origin, current household, personal interests, and lifestyle

Review of Systems

 Documents presence or absence of common symptoms related to each major body system

THE COMPREHENSIVE ADULT HEALTH HISTORY

As you elicit the adult health history, be sure to include the following: date and time of history; identifying data, which include age, gender, marital status, and occupation; and reliability, which reflects the quality of information the patient provides.



CHIEF COMPLAINT(S)

Quote the patient's own words. "My stomach hurts and I feel awful"; or "I have come for my regular check-up."



PRESENT ILLNESS

This section is a complete, clear, and chronologic account of the problems prompting the patient to seek care. It should include the problem's onset, the setting in which it has developed, its manifestations, and any treatments.

4 Overview: Physical Examination and History Taking

Every principal symptom should be well characterized, with descriptions of the seven features listed below and *pertinent positives* and *negatives* from relevant areas of the Review of Systems.

- Location
- Quality
- Quantity or severity
- Timing, including onset, duration, and frequency
- Setting in which they occur
- · Aggravating and relieving factors
- Associated manifestations

In addition, list *medications*, including name, dose, route, and frequency of use; *allergies*, including *specific reactions* to each medication; *tobacco* use; and *alcohol* and *drug* use.



PAST HISTORY

List childhood illnesses, then list adult illnesses in each of four areas:

- Medical (e.g., diabetes, hypertension, hepatitis, asthma, HIV), with dates of onset; also information about hospitalizations with dates; number and gender of sexual partners; risky sexual practices
- Surgical (dates, indications, and types of operations)
- Obstetric/gynecologic (obstetric history, menstrual history, birth control, and sexual function)
- Psychiatric (illness and time frame, diagnoses, hospitalizations, and treatments)

Also discuss *health maintenance*, including *immunizations*, such as tetanus, pertussis, diphtheria, polio, measles, rubella, mumps, influenza, varicella, hepatitis B, *Haemophilus influenzae* type b, and pneumococcal vaccine; and *screening tests*, such as tuberculin tests, Pap smears, mammograms, stools for occult blood, and cholesterol tests, together with the results and the dates they were last performed.



FAMILY HISTORY

Outline or diagram the age and health, or age and cause of death, of each immediate relative, including grandparents, parents, siblings, children, and grandchildren. Record the following conditions as either *present or absent* in the family: hypertension, coronary artery disease, elevated cholesterol levels, stroke, diabetes, thyroid or renal disease, cancer (specify type), arthritis, tuberculosis, asthma or lung disease, headache, seizure disorder, mental illness, suicide, alcohol or drug addiction, and allergies, as well as symptoms that the patient reports.



PERSONAL AND SOCIAL HISTORY

Include occupation and the last year of schooling; home situation and significant others; sources of stress, both recent and long-term; important life experiences, such as military service; leisure activities; religious affiliation and spiritual beliefs; and activities of daily living (ADLs). Also include lifestyle habits such as exercise and diet, safety measures, and alternative health care practices.



REVIEW OF SYSTEMS

These questions go from "head to toe." Start with a fairly general question. This allows you to shift to more specific questions about systems that may be of concern. For example, "How are your ears and hearing?" "How about your lungs and breathing?" "Any trouble with your heart?" "How is your digestion?" The Review of Systems questions may uncover problems that the patient overlooked. Remember to move major health events to the Present Illness or Past History in your write-up.

Some clinicians do the *Review of Systems* during the physical examination. If the patient has only a few symptoms, this combination can be efficient. If he or she has multiple symptoms, the flow of both the history and the examination can be disrupted.

General. Usual weight, recent weight change, any clothes that fit more tightly or loosely than before; weakness, fatigue, fever.

Skin. Rashes, lumps, sores, itching, dryness, color change, changes in hair or nails, changes in size or color of moles.

Head, Eyes, Ears, Nose, Throat (HEENT). Head: Headache, head injury, dizziness, lightheadedness. Eyes: Vision, glasses or contact lenses, last examination, pain, redness, excessive tearing, double or blurred vision, spots, specks, flashing lights, glaucoma, cataracts. Ears: Hearing, tinnitus, vertigo, earache, infection, discharge. If hearing is decreased, use or nonuse of hearing aid. Nose and sinuses: Frequent colds, nasal stuffiness, discharge or itching, hay fever, nosebleeds, sinus trouble. Throat (or mouth and pharynx): Condition of teeth and gums; bleeding gums; dentures, if any, and how they fit; last dental examination; sore tongue; dry mouth; frequent sore throats; hoarseness.

Neck. Lumps, "swollen glands," goiter, pain, stiffness.

Breasts. Lumps, pain or discomfort, nipple discharge, self-examination practices.

Respiratory. Cough, sputum (color, quantity), hemoptysis, dyspnea, wheezing, pleurisy, last chest x-ray. You may wish to include asthma, bronchitis, emphysema, pneumonia, and tuberculosis.

Cardiovascular. Heart trouble, hypertension, rheumatic fever, heart murmurs, chest pain or discomfort, palpitations, dyspnea, orthopnea, paroxysmal nocturnal dyspnea, edema, past electrocardiographic or other heart test results.

Gastrointestinal. Trouble swallowing, heartburn, appetite, nausea, bowel movements, color and size of stools, change in bowel habits, rectal bleeding or black or tarry stools, hemorrhoids, constipation, diarrhea. Abdominal pain, food intolerance, excessive belching or passing of gas. Jaundice, liver or gallbladder trouble, hepatitis.

Urinary. Frequency of urination, polyuria, nocturia, urgency, burning or pain on urination, hematuria, urinary infections, kidney stones, incontinence; in males, reduced caliber or force of urinary stream, hesitancy, dribbling.

Genital. *Male:* Hernias, discharge from or sores on penis, testicular pain or masses, history of sexually transmitted diseases (STDs) and treatments, testicular self-examination practices. Sexual habits, interest, function, satisfaction,

birth control methods, condom use, problems. Exposure to HIV infection. Female: Age at menarche; regularity, frequency, and duration of periods; amount of bleeding, bleeding between periods or after intercourse, last menstrual period; dysmenorrhea, premenstrual tension; age at menopause, menopausal symptoms, postmenopausal bleeding. In patients born before 1971, exposure to diethylstilbestrol (DES) from maternal use during pregnancy. Vaginal discharge, itching, sores, lumps, STDs and treatments. Number of pregnancies, number and type of deliveries, number of abortions (spontaneous and induced); complications of pregnancy; birth control methods. Sexual preference, interest, function, satisfaction, problems (including dyspareunia). Exposure to HIV infection.

Peripheral Vascular. Intermittent claudication; leg cramps; varicose veins; past clots in veins; swelling in calves, legs, or feet; color change in fingertips or toes during cold weather; swelling with redness or tenderness.

Musculoskeletal. Muscle or joint pain, stiffness, arthritis, gout, backache. If present, describe location of affected joints or muscles, any swelling, redness, pain, tenderness, stiffness, weakness, or limitation of motion or activity; include timing of symptoms (e.g., morning or evening), duration, and any history of trauma. Neck or low back pain. Joint pain with systemic features such as fever, chills, rash, anorexia, weight loss, or weakness.

Psychiatric. Nervousness; tension; mood, including depression, memory change, suicide attempts, if relevant.

Neurologic. Changes in mood, attention, or speech; changes in orientation, memory, insight, or judgment; headache, dizziness, vertigo; fainting, blackouts, seizures, weakness, paralysis, numbness or loss of sensation, tingling or "pins and needles," tremors or other involuntary movements; seizures.

Hematologic. Anemia, easy bruising or bleeding, past transfusions, transfusion reactions.

Endocrine. Thyroid trouble, heat or cold intolerance, excessive sweating, excessive thirst or hunger, polyuria, change in glove or shoe size.

HE PHYSICAL EXAMINATION: APPROACH AND OVERVIEW

Conduct a comprehensive physical examination on most new patients or patients being admitted to the hospital. For more problem-oriented, or focused, assessments, the presenting complaints will dictate which segments you elect to perform.

- The key to a thorough and accurate physical examination is a systematic sequence of examination. With effort and practice, you will acquire your own routine sequence. This book recommends examining from the patient's right side.
- Apply the techniques of inspection, palpation, auscultation, and percussion to each body region, but be sensitive to the whole patient.
- Minimize the number of times you ask the patient to change position from supine to sitting, or standing to lying supine.
- For an overview of the physical examination, study the sequence that follows. *Note that clinicians vary in* where they place different segments, especially for the musculoskeletal and nervous systems.



BEGINNING THE EXAMINATION: SETTING THE STAGE

Take the following steps to prepare for the physical examination.

Preparing for the Physical Examination

- Reflect on your approach to the patient.
- Adjust the lighting and the environment.
- Make the patient comfortable.
- Determine the scope of the examination.
- Choose the sequence of the examination.
- Observe the correct examining position (the patient's right side) and handedness.

Think through your approach, your professional demeanor, and how to make the patient comfortable and relaxed. Wash your hands in the patient's presence before beginning.

THE PHYSICAL EXAMINATION: SUMMARY OF SUGGESTED SEQUENCE

General survey

- Vital signs
- Skin: upper torso, anterior and posterior
- Head and neck, including thyroid and lymph nodes
- Optional: Nervous system (mental status, cranial nerves, upper extremity motor strength, bulk, tone; cerebellar function)
- Thorax and lungs
- Breasts
- Musculoskeletal as indicated: upper extremities
- Cardiovascular, including JVP, carotid upstrokes and bruits, PMI. etc.
- Cardiovascular, for S₃ and murmur of mitral stenosis
 - Nervous system: lower extremity motor strength, bulk, tone: sensation: reflexes: Babinskis

- ♀/ ♀ Musculoskeletal, as indicated
 - Optional: Skin, anterior and posterior
 - Optional: Nervous system, including gait Optional: Musculoskeletal,
- comprehensive examination
- examination
 - Cardiovascular, for murmur of aortic insufficiency
- Optional: Thorax and lungs anterior
 - Breasts and axillae
 - Abdomen
 - Peripheral vascular; Optional: Skin-lower torso and extremities

Key to the Symbols for the Patient's Position

Sitting

Lying supine, with head of bed raised 30 degrees

Same, turned partly to left side

Standing

Lying supine, with hips flexed, abducted, and externally rotated, and knees flexed (lithotomy position)

- Lying on the left side (left lateral decubitus)

Sitting, leaning forward

Lying supine

Each symbol pertains until a new one appears. Two symbols separated by a slash indicate either or both positions.

Reflect on Your Approach to the Patient. Identify yourself as a student. Try to appear calm, organized, and competent, even if you feel differently. If you forget to do part of the examination, this is not uncommon, especially at first! Simply examine that area out of sequence, but smoothly.

Adjust Lighting and the Environment. Adjust the bed to a convenient height (be sure to lower it when finished!). Ask the patient to move toward you if this makes it easier to do your physical examination.

Good lighting and a quiet environment are important. *Tangential lighting* is optimal for structures such as the jugular venous pulse, the thyroid gland, and the apical impulse of the heart. It throws contours, elevations, and depressions, whether moving or stationary, into sharper relief.

Make the Patient Comfortable. Show concern for privacy and modesty.

- Close nearby doors and draw curtains before beginning.
- Acquire the art of draping the patient with the gown or draw sheet as you learn each examination segment in future chapters. The goal is to visualize one body area at a time.
- As you proceed, keep the patient informed, as when checking for the femoral pulse. Also try to gauge how much the patient wants to know.
- Make sure your instructions to the patient at each step are courteous and clear.
- Watch the patient's facial expression and even ask "Is it okay?" as you move through the examination.

When you have finished, tell the patient your general impressions and what to expect next. Lower the bed to avoid risk of falls and raise the bedrails if needed. As you leave, clean your equipment, dispose of waste materials, and wash your hands.

Determine the Scope of the Examination. Comprehensive or Focused? Choose whether to do a comprehensive or focused examination.

Choose the Sequence of the Examination. The sequence of the examination should

- maximize the patient's comfort
- avoid unnecessary changes in position, and
- enhance the clinician's efficiency.

In general, move from head to toe. An important goal as a student is to develop your own sequence with these principles in mind. See Chapter 1 for a suggested examination sequence.

Observe the Correct Examining Position and Handedness.

Note that it is more reliable to estimate jugular venous pressure from the right, the palpating hand rests more comfortably on the apical impulse, the right kidney is more frequently palpable than the left, and examining tables are frequently positioned to accommodate a righthanded approach. To examine the supine patient, you can examine the head, neck, and anterior chest. Then roll the patient onto each side to listen to the lungs. examine the back, and inspect the skin. Roll the patient back and finish the rest of the examination with the patient again supine.

THE COMPREHENSIVE ADULT PHYSICAL EXAMINATION

General Survey. Continue this survey throughout the patient visit. Observe general state of health, height, build, and sexual development. Note posture, motor activity, and gait; dress, grooming, and personal hygiene; and any odors of the body or breath. Watch facial expressions and note manner, affect, and reactions to persons and things in the environment. Listen to the patient's manner of speaking and note the state of awareness or level of consciousness.

Vital Signs. Ask the patient to sit on the edge of the bed or examining table, unless this position is contraindicated. Stand in front of the patient, moving to either side as needed. Measure the blood pressure. Count pulse and respiratory rate. If indicated, measure body temperature.

Skin. Observe the face. Identify any lesions, noting their location, distribution, arrangement, type, and color. Inspect and palpate the hair and nails. Study the patient's hands. Continue to assess the skin as you examine the other body regions.

HEENT. Darken the room to promote pupillary dilation and visibility of the fundi. Head: Examine the hair, scalp, skull, and face. Eyes: Check visual acuity and screen the visual fields. Note position and alignment of the eyes. Observe the evelids. Inspect the sclera and conjunctiva of each eve. With oblique lighting, inspect each cornea, iris, and lens. Compare the pupils, and test their reactions to light. Assess extraocular movements. With an ophthalmoscope, inspect the ocular fundi. Ears: Inspect the auricles, canals, and drums. Check auditory acuity. If acuity is diminished, check lateralization (Weber test) and compare air and bone conduction (Rinne test). Nose and sinuses: Examine the external nose; using a light and nasal speculum, inspect nasal mucosa, septum, and turbinates. Palpate for tenderness of the frontal and maxillary sinuses. Throat (or mouth and pharynx): Inspect the lips, oral mucosa, gums, teeth, tongue, palate, tonsils, and pharvnx. (You may wish to assess the Cranial Nerves at this point in the examination.)

Neck. Move behind the sitting patient to feel the thyroid gland and to examine the back, posterior thorax, and lungs. Inspect and palpate the cervical lymph nodes. Note any masses or unusual pulsations in the neck. Feel for any deviation of the trachea. Observe sound and effort of the patient's breathing. Inspect and palpate the thyroid gland.

Back. Inspect and palpate the spine and muscles.

Posterior Thorax and Lungs. Inspect and palpate the spine and muscles of the *upper* back. Inspect, palpate, and percuss the chest. Identify the level of diaphragmatic dullness on each side. Listen to the breath sounds; identify any adventitious (or added) sounds, and, if indicated, listen to the transmitted voice sound (see p. 148).

Breasts, Axillae, and Epitrochlear Nodes. The patient is still sitting. Move to the front again. Female: Inspect the breasts with patient's arms relaxed, then elevated, and then with her hands pressed on her hips. Male and Female: Inspect the axillae and feel for the axillary nodes; feel for the epitrochlear nodes.

A Note on the Musculoskeletal System: By now, you have made preliminary observations of the musculoskeletal system, including the hands, the upper back, and, in women, the shoulders' range of motion (ROM). Use these observations to decide whether a full musculoskeletal examination is warranted: with the patient still sitting, examine the hands, arms, shoulders, neck, and temporomandibular joints. Inspect and palpate the joints and check their ROM. (You may choose to examine upper-extremity muscle bulk, tone, strength, and reflexes at this time, or you may decide to wait.)

Palpate the breasts, while continuing your inspection.

- Anterior Thorax and Lungs. The patient position is **supine.** Ask the patient to lie down. Stand at the *right side* of the patient's bed. Inspect, palpate, and percuss the chest. Listen to the breath sounds, any adventitious sounds, and, if indicated, transmitted voice sounds.
- Cardiovascular System. Elevate head of bed to about 30°, adjusting as necessary to see the jugular venous pulsations. Observe the jugular venous pulsations, and measure the jugular venous pressure in relation to the sternal angle. Inspect and palpate the carotid pulsations. Listen for carotid bruits.
- ○→ / Ask the patient to roll partly onto the left side while you listen at the apex. Then have the patient roll back to supine while you listen to the rest of the heart. Ask the patient to sit, lean forward, and exhale while you listen for the murmur of aortic regurgitation. Inspect and palpate the precordium. Note the location, diameter, amplitude, and duration of the apical impulse. Listen at the apex and the lower sternal border with the bell of a stethoscope. Listen at each auscultatory area with the diaphragm. Listen for S₁ and S_2 and for physiologic splitting of \bar{S}_2 . Listen for any abnormal heart sounds or murmurs.
- O—/ Peripheral Vascular System. With the patient supine, palpate the femoral pulses and, if indicated, popliteal pulses. Palpate the inguinal lymph nodes. Inspect for edema, discoloration, or ulcers in the lower extremities. Palpate for pitting edema. With the patient standing, inspect for varicose veins.

14 Overview: Physical Examination and History Taking

• Abdomen. Lower the head of the bed to the flat position. The patient should be supine. Inspect, auscultate, and percuss. Palpate lightly, then deeply. Assess the liver and spleen by percussion and then palpation. Try to feel the kidneys; palpate the aorta and its pulsations. If you suspect kidney infection, percuss posteriorly over the costovertebral angles.

Lower Extremities. Examine the legs, assessing the three systems (see next page) while the patient is still supine. Each of these systems can be further assessed when the patient stands.

Nervous System. The patient is sitting or supine. The examination of the nervous system can also be divided into the upper-extremity examination (when the patient is still sitting) and the lower-extremity examination (when the patient is supine) after examination of the peripheral nervous system.

- Mental Status: If indicated and not done during the interview, assess orientation, mood, thought process, thought content, abnormal perceptions, insight and judgment, memory and attention, information and vocabulary, calculating abilities, abstract thinking, and constructional ability.
- Cranial Nerves: If not already examined, check sense of smell, funduscopic examination, strength of the temporal and masseter muscles, corneal reflexes, facial movements, gag reflex, strength of the trapezia and sternomastoid muscles, and protrusion of tongue.
- Motor System: Muscle bulk, tone, and strength of major muscle groups. Cerebellar function: rapid alternating movements (RAMs), point-to-point movements such as finger to nose (F → N) and heel to shin (H → S); gait. Observe patient's gait and ability to walk heel to toe, on toes, and on heels; to hop in place; and to do shallow knee bends. Do a Romberg test; check for pronator drift.
- Sensory System: Pain, temperature, light touch, vibrations, and discrimination. Compare right and left sides and distal with proximal areas on the limbs.
- Reflexes: Include biceps, triceps, brachioradialis, patellar, Achilles deep tendon reflexes; also plantar reflexes or Babinski reflex (see p. 328-331)

Additional Examinations. The rectal and genital examinations are often performed at the end of the physical examination. Patient positioning is as indicated.

Male Genitalia and Hernias. Examine the penis and scrotal contents. Check for hernias.

• Rectal Examination in Men. The patient is lying on his left side for the rectal examination. Inspect the sacrococcygeal and perianal areas. Palpate the anal canal, rectum, and prostate. (If the patient cannot stand, examine the genitalia before doing the rectal examination.)

Genital and Rectal Examination in Women. The patient is supine in the lithotomy position. Sit during the examination with the speculum, then stand during bimanual examination of uterus, adnexa, and rectum. Examine the external genitalia, vagina, and cervix. Obtain a Pap smear. Palpate the uterus and adnexa. Do a bimanual and rectal examination.



STANDARD AND UNIVERSAL PRECAUTIONS

The Centers for Disease Control and Prevention (CDC) have issued several guidelines to protect patients and examiners from the spread of infectious disease. All clinicians examining patients are well advised to study and observe these precautions at the CDC Web-sites. Advisories for standard and methicillin-resistant Staphylococcus aureus (MRSA) precautions and for universal precautions are briefly summarized below.

• Standard and MRSA precautions: Standard precautions are based on the principle that all blood, body fluids. secretions, excretions except sweat, nonintact skin, and mucous membranes may contain transmissible infectious agents. These practices apply to all patients in any setting. They include: hand hygiene; when to use gloves, gowns, and mouth, nose, and eye protection; respiratory hygiene and cough etiquette; patient isolation criteria; precautions relating to equipment, toys and solid surfaces, and handling of laundry; and safe needle injection practices.

16 Overview: Physical Examination and History Taking

Be sure to wash your hands before and after examining the patient. This will show your concern for the patient's welfare and display your awareness of a critical component of patient safety. Antimicrobial fast-drying soaps are often within easy reach. Change your white coat frequently since cuffs can become damp and smudged.

• Universal precautions: Universal precautions are a set of precautions designed to prevent transmission of HIV, hepatitis B virus (HBV), and other bloodborne pathogens when providing first aid or health care. The following fluids are considered potentially infectious: all blood and other body fluids containing visible blood, semen, and vaginal secretions; and cerebrospinal, synovial, pleural, peritoneal, pericardial, and amniotic fluids. Protective barriers include gloves, gowns, aprons, masks, and protective eyewear. All healthcare workers should observe the important precautions for safe injections and prevention of injury from needlesticks, scalpels, and other sharp instruments and devices. Report to your health service immediately if such injury occurs.

2

Clinical Reasoning, Assessment, and Recording Your Findings

ASSESSMENT AND PLAN: THE PROCESS OF CLINICAL REASONING

Because assessment takes place in the clinician's mind, the process of clinical reasoning often seems inaccessible to beginning students. As an active learner, ask your teachers and clinicians to elaborate on the fine points of their clinical reasoning and decision making.

As you gain experience, your thinking will begin at the outset of the patient encounter, not at the end. Listed below are principles underlying the process of clinical reasoning and certain explicit steps to help guide your thinking.

IDENTIFYING PROBLEMS AND MAKING DIAGNOSES: STEPS IN CLINICAL REASONING

- Identify abnormal findings. Make a list of the patient's symptoms, the signs you observed during physical examination, and any available results of laboratory reports.
- Localize these findings anatomically. The symptom of a scratchy throat and the sign of an erythematous inflamed pharynx, for example, clearly localize the problem to the pharynx. Some symptoms and signs, such as fatigue or fever, cannot be localized but are useful in the next steps.
- Interpret the findings in terms of the probable process. There are a number of *pathologic* processes, including congenital, inflammatory or infectious, immunologic, neoplastic, metabolic, nutritional, degenerative,

IDENTIFYING PROBLEMS AND MAKING DIAGNOSES: STEPS IN CLINICAL REASONING (CONTINUED)

vascular, traumatic, and toxic. Other problems are pathophysiologic, reflecting derangements of biologic functions, such as congestive heart failure. Still other problems are psychopathologic, such as headache as an expression of a somatization disorder.

- Make hypotheses about the nature of the patient's problems. Draw on your knowledge, experience, and reading about patterns of abnormalities and diseases. By consulting the clinical literature, you will embark on the lifelong goal of evidence-based decision making. The following steps should help:
 - 1. Select the most specific and critical findings to support your hypothesis.
 - 2. Match your findings against all the conditions you know that can produce them.
 - 3. Eliminate the diagnostic possibilities that fail to explain the findings.
 - 4. Weigh the competing possibilities and select the most likely diagnosis.
 - 5. Give special attention to potentially life-threatening and treatable conditions. One rule of thumb is always to include "the worst case scenario" in your list of differential diagnoses and make sure you have ruled out that possibility based on your findings and patient assessment.
- Test your hypotheses. You may need further history, additional maneuvers on physical examination, or laboratory studies or x-rays to confirm or to rule out your tentative diagnosis or to clarify which of a few possible diagnoses is most likely.
- Establish a working diagnosis. Make this at the highest level of explicitness and certainty that the data allow. You may be limited to a symptom, such as "tension headache, cause unknown." At other times, you can define a problem explicitly in terms of its structure, process, and cause, such as "bacterial meningitis, pneumococcal." Routinely listing Health Maintenance helps you track several important health concerns more effectively: immunizations, screening measures (e.g., mammograms, prostate examinations),

IDENTIFYING PROBLEMS AND MAKING DIAGNOSES: STEPS IN CLINICAL REASONING (CONTINUED)

instructions regarding nutrition and breast or testicular self-examinations, recommendations about exercise or use of seat belts, and responses to important life events.

• Develop a plan agreeable to the patient. Identify and record a Plan for each patient problem, ranging from tests to confirm or further evaluate a diagnosis; to consultations for subspecialty evaluation; to additions, deletions, or changes in medication: or to arranging a family meeting.

CASE OF MRS.

Now study the case of Mrs. N. Scrutinize the findings recorded, apply your clinical reasoning, and analyze the assessment and plan.

Health History

8/30/08 11:00 AM

Mrs. N is a pleasant, 54-year-old widowed saleswoman residing in Amarillo, Texas.

Referral, None

Source and Reliability. Self-referred; seems reliable.

Chief Complaint: "My head aches."

Present Illness: For about 3 months, Mrs. N has had increasing problems with frontal headaches. These are usually bifrontal, throbbing, and mild to moderately severe. She has missed work on several occasions because of associated nausea and vomiting. Headaches now average once a week, usually are related to stress, and last 4 to 6 hours. They are relieved by sleep and putting a damp towel over the forehead. There is little relief from aspirin. No associated visual changes, motorsensory deficits, or paresthesias.

"Sick headaches" with nausea and vomiting began at age 15, recurred throughout her mid-20s, then decreased to one every 2 or 3 months and almost disappeared.

The patient reports increased pressure at work from a new and demanding boss; she is also worried about her daughter (see Personal and Social History). She thinks her headaches may be like those in the past but wants to be sure, because her mother died following a stroke. She is concerned that they interfere with her work and make her irritable with her family. She eats three meals a day and drinks three cups of coffee per day; cola at night.

Medications. Aspirin, 1 to 2 tablets every 4 to 6 hours as needed. "Water pill" in the past for ankle swelling, none recently.

*Allergies. Ampicillin causes rash.

Tobacco. About 1 pack of cigarettes per day since age 18 (36 pack-years).

Alcohol/drugs. Wine on rare occasions. No illicit drugs.


Past History

Childhood Illnesses. Measles, chickenpox. No scarlet fever or rheumatic fever.

Adult Illnesses. Medical: Pyelonephritis, 1998, with fever and right flank pain; treated with ampicillin; developed generalized rash with itching several days later. Reports kidney x-rays were normal; no recurrence of infection. Surgical: Tonsillectomy, age 6; appendectomy, age 13. Sutures for laceration, 2001, after stepping on glass. Ob/Gyn: 3-3-0-3, with normal vaginal deliveries. 3 living children. Menarche age 12. Last menses 6 months ago. Little interest in sex, and not sexually active. No concerns about HIV infection. Psychiatric: None.

Health Maintenance. Immunizations: Oral polio vaccine, year uncertain; tetanus shots × 2, 1991, followed with booster 1 year later; flu vaccine, 2000, no reaction. Screening tests: Last Pap smear, 2004, normal. No mammograms to date.

^{*}You may wish to add an asterisk or underline important points.



OR

Father died at age 43 in train accident. Mother died at age 67 from stroke; had varicose veins, headaches.

One brother, 61, with hypertension, otherwise well; second brother, 58, well except for mild arthritis; one sister, died in infancy of unknown cause.

Husband died at age 54 of heart attack.

Daughter, 33, with migraine headaches, otherwise well; son, 31, with headaches; son, 27, well.

No family history of diabetes, tuberculosis, heart or kidney disease, cancer, anemia, epilepsy, or mental illness.

Personal and Social History: Born and raised in Lake City, finished high school, married at age 19. Worked as sales clerk for 2 years, then moved with husband to Amarillo, had 3 children. Returned to work 15 years ago because of financial pressures. Children all married. Four years ago Mr. N died suddenly of a heart attack, leaving little savings. Mrs. N has moved to small apartment to be near daughter, Dorothy. Dorothy's husband, Arthur, has an alcohol problem. Mrs. N's apartment now a haven for Dorothy and her 2 children, Kevin, 6 years, and

Linda, 3 years. Mrs. N feels responsible for helping them; feels tense and nervous but denies depression. She has friends but rarely discusses family problems: "I'd rather keep them to myself. I don't like gossip." No church or other organizational support. She is typically up at 7:00 A.M., works 9:00 to 5:30, eats dinner alone.

Exercise and diet. Gets little exercise. Diet high in carbohydrates.

Safety measures. Uses seat belt regularly. Uses sunblock. Medications kept in an unlocked medicine cabinet. Cleaning solutions in unlocked cabinet below sink. Mr. N's shotgun and box of shells in unlocked closet upstairs.

Review of Systems

General. *Has gained about 10 lbs in the past 4 years.

Skin. No rashes or other changes.

Head, Eyes, Ears, Nose, Throat (HEENT). See Present Illness. No history of head injury. Eyes: Reading glasses for 5 years, last checked 1 year ago. No symptoms. Ears: Hearing good. No tinnitus, vertigo, infections. Nose, sinuses: Occasional mild cold. No hay fever, sinus trouble. *Throat (or mouth and pharynx): Some bleeding of gums recently. Last dental visit 2 years ago. Occasional canker sore.

Neck. No lumps, goiter, pain. No swollen glands.

Breasts. No lumps, pain, discharge. Does breast self-exam sporadically.

Respiratory. No cough, wheezing, shortness of breath. Last chest x-ray, 1986, St. Mary's Hospital; unremarkable.

Cardiovascular. No known heart disease or high blood pressure; last blood pressure taken in 2003. No dyspnea, orthopnea, chest pain, palpitations. Has never had an electrocardiogram (ECG).

Gastrointestinal. Appetite good; no nausea, vomiting, indigestion. Bowel movement about once daily, *though sometimes has hard stools for 2 to 3 days when especially tense; no diarrhea or bleeding. No pain, jaundice, gallbladder or liver problems.

Urinary. No frequency, dysuria, hematuria, or recent flank pain; nocturia × 1, large volume. *Occasionally loses some urine when coughs hard.

Genital. No vaginal or pelvic infections. No dyspareunia.

Peripheral Vascular. Varicose veins appeared in both legs during first pregnancy. For 10 years, has had swollen ankles after prolonged standing; wears light elastic pantyhose; tried "water pill" 5 months ago, but it didn't help much; no history of phlebitis or leg pain.

Musculoskeletal. Mild, aching, low back pain, often after a long day's work; no radiation down the legs; used to do back exercises but not now. No other joint pain.

Psychiatric. No history of depression or treatment for psychiatric disorders. See also Present Illness and Personal and Social History.

No fainting, seizures, motor or sensory loss. Neurologic. Memory good.

Hematologic. Except for bleeding gums, no easy bleeding. No anemia.

Endocrine. No known thyroid trouble, temperature intolerance. Sweating average. No symptoms or history of diabetes.

Physical Examination

Mrs. N is a short, overweight, middle-aged woman, who is animated and responds quickly to questions. She is somewhat tense, with moist, cold hands. Her hair is fixed neatly and her clothes are immaculate. Her color is good, and she lies flat without discomfort.

Vital Signs. Ht (without shoes) 157 cm (5'2"). Wt (dressed) 65 kg (143 lb). BMI 26. BP 164/98 right arm, supine; 160/96 left arm, supine; 152/88 right arm, supine with wide cuff. Heart rate (HR) 88 and regular. Respiratory rate (RR) 18. Temperature (oral) 98.6°F.

Skin. Palms cold and moist, but color good. Scattered cherry angiomas over upper trunk. Nails without clubbing, cyanosis.

Head, Eyes, Ears, Nose, Throat (HEENT). Head: Hair of average texture. Scalp without lesions, normocephalic/atraumatic (NC/AT). Eyes: Vision 20/30 in each eye. Visual fields full by confrontation. Conjunctiva pink; sclera white. Pupils 4 mm (continued) constricting to 2 mm, round, regular, equally reactive to light. Extraocular movements intact. Disc margins sharp, without hemorrhages, exudates. No arteriolar narrowing or A-V nicking. Ears: Wax partially obscures right tympanic membrane (TM); left canal clear, TM with good cone of light. Acuity good to whispered voice. Weber midline. AC > BC. Nose: Mucosa pink, septum midline. No sinus tenderness. Mouth: Oral mucosa pink. Several interdental papillae red, slightly swollen. Dentition good. Tongue midline, with 3 × 4 mm shallow white ulcer on red base on undersurface near tip; tender but not indurated. Tonsils absent. Pharynx without exudates.

Neck. Neck supple. Trachea midline. Thyroid isthmus barely palpable, lobes not felt.

Lymph Nodes. Small (<1 cm), soft, nontender, and mobile tonsillar and posterior cervical nodes bilaterally. No axillary or epitrochlear nodes. Several small inquinal nodes bilaterally, soft and nontender.

Thorax and Lungs. Thorax symmetric with good excursion. Lungs resonant. Breath sounds vesicular with no added sounds. Diaphragms descend 4 cm bilaterally.

Cardiovascular. Jugular venous pressure 1 cm above the sternal angle, with head of examining table raised to 30°. Carotid upstrokes brisk, without bruits. Apical impulse discrete and tapping, barely palpable in the 5th left interspace, 8 cm lateral to the midsternal line. Good S₁, S₂; no S₃ or S₄. A II/VI medium-pitched midsystolic murmur at the 2nd right interspace; does not radiate to the neck. No diastolic murmurs.

Breasts. Pendulous, symmetric. No masses; nipples without discharge.

Abdomen. Protuberant. Well-healed scar, right lower quadrant. Bowel sounds active. No tenderness or masses. Liver span 7 cm in right midclavicular line; edge smooth, palpable 1 cm below right costal margin (RCM). Spleen and kidneys not felt. No costovertebral angle tenderness (CVAT).

Genitalia. External genitalia without lesions. Mild cystocele at introitus on straining. Vaginal mucosa pink. Cervix pink, parous, and without discharge. Uterus anterior, midline, smooth, not enlarged. Adnexa not palpated due to obesity and poor relaxation. No cervical or adnexal tenderness. Pap smear taken. Rectovaginal wall intact.

Rectal. Rectal vault without masses. Stool brown, negative for occult blood.

Extremities. Warm and without edema. Calves supple, nontender.


Peripheral Vascular. Trace edema at both ankles. Moderate varicosities of saphenous veins in both lower extremities. No stasis pigmentation or ulcers. Pulses (2 + = brisk, or normal):

	Radial	Femoral	Popliteal	Dorsalis Pedis	Posterior Tibial
RT	2+	2+	2+	2+	2+
LT	2+	2+	2+	Absent	2+

Musculoskeletal. No joint deformities. Good range of motion in hands, wrists, elbows, shoulders, spine, hips, knees, ankles.

Neurologic. Mental Status: Tense but alert and cooperative. Thought coherent. Oriented to person, place, and time. Cranial Nerves: II-XII intact. Motor: Good muscle bulk and tone. Strength 5/5 throughout (see p. 321 for grading system). Cerebellar: Rapid alternating movements (RAMs), point-to-point movements intact. Gait stable, fluid. Sensory: Pinprick, light touch, position sense, vibration, and stereognosis intact. Romberg negative. Reflexes:

			Brachio-			
	Biceps	Triceps	radialis	Patellar	Achilles	Plantar
RT LT	2+ 2+	2+ 2+	2+ 2+	2+ 2+/2+	1+ 1+	\downarrow



Laboratory Data

None Currently. See Plan.

ASSESSMENT AND PLAN

1. Migraine headaches. A 54-year-old woman with migraine headaches since childhood, with a throbbing vascular pattern and frequent nausea and vomiting. Headaches are associated with stress and relieved by sleep and cold compresses. There is no papilledema, and there are no motor or sensory deficits on the neurologic examination. The differential diagnosis includes tension headache, also associated with stress, but there is no relief with massage, and the pain is more throbbing than aching. There are no fever, stiff neck, or focal findings to suggest meningitis, and the lifelong recurrent pattern makes subarachnoid hemorrhage unlikely (usually described as "the worst headache of my life").

Plan:

- Discuss features of migraine vs. tension headaches.
- Discuss biofeedback and stress management.
- Advise patient to avoid caffeine, including coffee, colas, and other caffeinated beverages.
- Start NSAIDs for headache, as needed.
- If needed next visit, begin prophylactic medication, because patient is having more than three migraines per month.
- 2. Elevated blood pressure. Systolic hypertension with wide cuff is present. May be related to obesity, also to anxiety from first visit. No evidence of end-organ damage to retina or heart.

Plan:

- Discuss standards for assessing blood pressure.
- Recheck blood pressure in 1 month, using wide cuff.
- Check basic metabolic panel; review urinalysis.
- Introduce weight reduction and/or exercise programs (see #4).
- Reduce salt intake.
- 3. Cystocele with occasional stress incontinence. Cystocele on pelvic examination, probably related to bladder relaxation. Patient is perimenopausal. Incontinence reported with coughing, suggesting alteration in bladder neck

ASSESSMENT AND PLAN (CONTINUED)

anatomy. No dysuria, fever, flank pain. Not taking any contributing medications. Usually involves small amounts of urine, no dribbling, so doubt urge or overflow incontinence.

Plan:

- Explain cause of stress incontinence.
- Review urinalysis.
- Recommend Kegel's exercises.
- Consider topical estrogen cream to vagina next visit if no improvement.
- 4. Overweight. Patient 5'2", weighs 143 lbs. BMI is ~26.

Plan:

- Explore diet history; ask patient to keep food intake diary.
- Explore motivation to lose weight; set target for weight loss by next visit.
- Schedule visit with dietitian.
- Discuss exercise program, specifically, walking 30 minutes most days each week.
- 5. Family stress. Son-in-law with alcohol problem; daughter and grandchildren seeking refuge in patient's apartment, leading to tensions in these relationships. Patient also has financial constraints. Stress currently situational. No evidence of major depression at present.

Plan:

- Explore patient's views on strategies to cope with stress.
- Explore sources of support, including Al-Anon for daughter and financial counseling for patient.
- Continue to monitor for depression.
- 6. Occasional musculoskeletal low back pain. Usually with prolonged standing. No history of trauma or motor vehicle accident. Pain does not radiate; no tenderness or motor-sensory deficits on examination. Doubt disc or nerve root compression, trochanteric bursitis, sacroiliitis.

Plan:

• Review benefits of weight loss and exercises to strengthen low back muscles.

(continued)

ASSESSMENT AND PLAN (CONTINUED)

7. **Tobacco abuse.** 1 pack per day for 36 years.

Plan:

- Check peak flow or FEV₁/FVC on office spirometry.
- Give strong warning to stop smoking.
- Offer referral to tobacco cessation program.
- Offer patch, current treatment to enhance abstinence.
- 8. Varicose veins, lower extremities. No complaints currently.
- 9. History of right pyelonephritis, 1998.
- 10. Ampicillin allergy. Developed rash but no other allergic reaction.
- 11. Health maintenance. Last Pap smear 2004; has never had a mammogram.

Plan:

- Teach patient breast self-examination; schedule mammogram.
- Schedule Pap smear next visit.
- Provide three stool quaiac cards; next visit discuss screening colonoscopy.
- Suggest dental care for mild gingivitis.
- Advise patient to move medications and caustic cleaning agents to locked cabinet—if possible, above shoulder height.

APPROACHING THE CHALLENGES OF CLINICAL DATA

As you can see from the case of Mrs. N, organizing the patient's clinical data poses several challenges. The following guidelines will help you address these challenges.

• Clustering data into single vs. multiple problems. The patient's age may help. Young people are more likely to have a single disease, while older people tend to have multiple diseases. The *timing* of symptoms is often useful. For example, an episode of pharyngitis 6 weeks ago

probably is unrelated to fever, chills, pleuritic chest pain, and cough that prompt an office visit today.

If symptoms and signs are in a single system, one disease may explain them. Problems in different, apparently unrelated systems often require more than one explanation. Again, knowledge of disease patterns is necessary.

Some diseases involve *multisystem conditions*. To explain cough, hemoptysis, and weight loss in a 60-year-old plumber who has smoked cigarettes for 40 years, you probably even now would rank lung cancer high in your list of differential diagnoses.

- Sifting through an extensive array of data. Try to tease out separate clusters of observations and analyze one cluster at a time. You also can ask a series of key questions that may steer your thinking in one direction. For example, you may ask what produces and relieves the patient's chest pain. If the answer is exercise and rest, you can focus on the cardiovascular and musculoskeletal systems and set the gastrointestinal system aside.
- Assessing the quality of the data. To avoid errors in interpreting clinical information, acquire the habits of skilled clinicians, summarized in the following.

TIPS FOR ENSURING THE QUALITY OF YOUR PATIENT ASSESSMENT

- Ask open-ended questions and listen carefully and patiently to the patient's story.
- Craft a thorough and systematic sequence to history taking and physical examination.
- Keep an open mind toward the patient, the patient's story, and your physical findings.
- Always include "the worst-case scenario" in your list of possible explanations of the patient's problem, and make sure it can be eliminated safely.
- Analyze any mistakes in data collection or interpretation.
- Confer with colleagues and review the pertinent medical literature to clarify uncertainties.
- Apply principles of evidence-based data analysis to patient information and testing.

- Improving your assessment of clinical data and laboratory tests. Apply several key principles for selecting and using clinical data and tests: *reliability*, *validity*, *sensitivity*, *specificity*, and *predictive value*. Learn to apply these principles to your clinical findings and the tests you order.
- Displaying clinical data. To use these principles, it is important to display the data in the 2 × 2 format diagrammed on page 32. Always using this format will ensure the accuracy of your calculations of sensitivity, specificity, and predictive value.

PRINCIPLES OF TEST SELECTION AND USE

Reliability: The reproducibility of a measurement. It indicates how well repeated measurements of the same relatively stable phenomenon will give the same result, also known as **precision.** Reliability may be measured for one observer or more observers.

Example. If on several occasions one clinician consistently percusses the same span of a patient's liver dullness, intraobserver reliability is good. If, on the other hand, several observers find quite different spans of liver dullness on the same patient, interobserver reliability is poor.

Validity: The closeness with which a measurement reflects the true value of an object. It indicates how closely a given observation agrees with "the true state of affairs," or the best possible measure of reality.

Example. Blood pressure measurements by mercury-based sphygmomanometers are less valid than intra-arterial pressure tracings.

Sensitivity: Identifies the proportion of people who test positive in a group of people known to have the disease or condition, or the proportion of people who are *true positives* compared with the total number of people who actually have the disease. When the observation or test is negative in people who have the disease, the result is termed *false negative*. Good observations or tests have a sensitivity of more than 90% and

(continued)

PRINCIPLES OF TEST SELECTION AND USE (CONTINUED)

help "rule out" disease because false negatives are few. Such observations or tests are especially useful for screening.

Example. The sensitivity of Homan's sign in the diagnosis of deep venous thrombosis (DVT) of the calf is 50%. In other words, compared with a group of patients with DVT confirmed by phlebogram, a much better test, only 50% will have a positive Homan's sign, so this sign, if absent, is not helpful, because 50% of patients may have DVT.

Specificity: Identifies the proportion of people who test negative in a group known to be without a given disease or condition, or the proportion of people who are true negatives compared with the total number of people without the disease. When the observation or test is positive in people without the disease, the result is termed false positive. Good observations or tests have a specificity of more than 90% and help "rule in" disease, because the test is rarely positive when disease is absent, and false positives are few.

Example: The specificity of serum amylase in patients with possible acute pancreatitis is 70%. In other words, of 100 patients without pancreatitis, 70% will have a normal serum amylase; in 30%, the serum amylase will be falsely elevated.

Predictive value: Indicates how well a given symptom, sign, or test result—either positive or negative—predicts the presence or absence of disease. Positive predictive value is the probability of disease in a patient with a positive (abnormal) test, or the proportion of "true positives" out of the total population tested. Negative predictive value is the probability of not having the condition or disease when the test is negative (normal), or the proportion of "true negatives" out of the total population tested.

Examples. In a group of women with palpable breast nodules in a cancer screening program, the proportion with confirmed breast cancer would constitute the positive predictive value of palpable breast nodules for diagnosing breast cancer. In a group of women without

(continued)

PRINCIPLES OF TEST SELECTION AND USE (CONTINUED)

palpable breast nodules in a cancer screening program, the proportion without confirmed breast cancer constitutes the *negative predictive value* of absence of breast nodules.

Sensitivity, specificity, and predictive values are illustrated in a 2×2 table, as shown below in an example of 200 people, half of whom have the disease in question. In this example, the disease prevalence of 50% is much higher than in most clinical situations. Because the positive predictive value increases with prevalence, its calculated value here is unusually high.

Disease								
	Present	Absent						
+	95 true positive observations <i>a</i>	95 false positive observations b	105 total positive observations					
Observation –	5 C false negative observations	d 90 true negative observations	95 total negative observations					
	100 total persons with the disease	100 total persons without the disease	200 total persons					
Observation 5								
Specificity = $\frac{d}{b+d} = \frac{90}{90+10} \times 100 = 90\%$								
Positive pred	lictive value = -	$\frac{a}{+b} = \frac{95}{95 + 10}$	× 100 = 90.5%					
Negative pre	edictive value = -	$\frac{d}{c+d} = \frac{90}{90+5}$	× 100 = 94.7%					

ORGANIZING THE PATIENT RECORD

A clear, well-organized clinical record is one of the most important adjuncts to your patient care. Think about the order and readability of the record and the amount of detail needed. Use the following checklist to make sure your record is clear, informative, and easy to follow.

CHECKLIST FOR YOUR PATIENT RECORD

Is the order clear?

Order is imperative. Make sure that future readers, including you, can find specific points of information easily. Keep the subjective items of the history, for example, in the history; do not let them stray into the physical examination. Did you . . .

- Make the headings clear?
- Accent your organization with indentations and spacing?
- Arrange the *Present Illness* in chronologic order, starting with the current episode, then filling in relevant background information?

Do the data included contribute directly to the assessment?

Spell out the supporting data—both positive and negative for every problem or diagnosis that you identify.

Are pertinent negatives specifically described?

Often portions of the history or examination suggest a potential or actual abnormality.

Examples. For the patient with notable bruises, record the "pertinent negatives," such as the absence of injury or violence, familial bleeding disorders, or medications or nutritional deficits that might lead to bruising.

For the patient who is depressed but not suicidal, record both facts. In the patient with a transient mood swing, on the other hand, a comment on suicide is unnecessary.

(continued)

CHECKLIST FOR YOUR PATIENT RECORD (CONTINUED)

Are there overgeneralizations or omissions of important data?

Remember that data not recorded are data lost. No matter how vividly you can recall selected details today, you probably will not remember them in a few months. The phrase "neurologic exam negative," even in your own handwriting, may leave you wondering in a few months' time, "Did I really do the sensory exam?"

Is there too much detail?

Avoid burying important information in a mass of excessive detail, to be discovered by only the most persistent reader. Omit most negative findings unless they relate directly to the patient's complaints or to specific exclusions in your diagnostic assessment. Do not list abnormalities that you did not observe. Instead, concentrate on a few major ones, such as "no heart murmurs," and try to describe structures concisely and positively.

Examples. "Cervix pink and smooth" indicates you saw no redness, ulcers, nodules, masses, cysts, or other suspicious lesions, but the description is shorter and much more readable.

You can omit certain body structures even though you examined them, such as normal eyebrows and eyelashes.

Are phrases and short words used appropriately? Is there unnecessary repetition of data?

Omit unnecessary words, such as those in parentheses in the examples below. This saves valuable time and space.

Examples. "Cervix is pink (in color)." "Lungs are resonant (to percussion)." "Liver is tender (to palpation)." "Both (right and left) ears with cerumen." "II/VI systolic ejection murmur (audible)." "Thorax symmetric (bilaterally)."

Omit repetitive introductory phrases such as "The patient reports no . . . ," because readers assume the patient is the source of the history unless otherwise specified.

Use short words instead of longer, fancier ones when they mean the same thing, such as "felt" for "palpated" or "heard" for "auscultated."

(continued)

CHECKLIST FOR YOUR PATIENT RECORD (CONTINUED)

Describe what you observed, not what you did. "Optic discs seen" is less informative than "disc margins sharp," even if it marks your first glimpse as an examiner!


Is the written style succinct? Is there excessive use of abbreviations?

Records are scientific and legal documents, so they should be clear and understandable. Using words and brief phrases instead of whole sentences is common, but abbreviations and symbols should be used only if they are readily understood. Likewise, an overly elegant style is less appealing than a concise summary.

Be sure your record is legible; otherwise, all that you have recorded is worthless to your readers.

Are diagrams and precise measurements included where appropriate?

Diagrams add greatly to the clarity of the record. Examples. Study the examples below:



CHECKLIST FOR YOUR PATIENT RECORD (CONTINUED)

To ensure accurate evaluations and future comparisons, make measurements in centimeters, not in fruits, nuts, or vegetables.

Example. "1 × 1 cm lymph node" vs. "a pea-sized lymph node. . . " Or "2 × 2 cm mass on the left lobe of the prostate" vs. "a walnut-sized prostate mass."

Is the tone of the write-up neutral and professional?

It is important to be objective. Hostile, moralizing, or disapproving comments have no place in the patient's record. Never use words, penmanship, or punctuation that are inflammatory or demeaning.

Example. Comments such as "Patient DRUNK and LATE TO CLINIC AGAIN!!" are unprofessional and set a bad example for other providers reading the chart. They also might prove difficult to defend in a legal setting.

Once you have completed your assessment and written record, you will find it helpful to generate a Problem List that summarizes the patient's problems for the front of the office or hospital chart. A sample *Problem List* for Mrs. N is provided next.

Sample Problem List								
Date Entered	Problem No.	Problem						
8/30/08	1	Migraine headaches						
	2	Elevated blood pressure						
	3	Cystocele with occasional stress incontinence						
	4	Overweight						
	5	Family stress						
	6	Low back pain						
	7	Tobacco abuse						
	8	Varicose veins						
	9	History of right pyelonephritis						
	10	Allergy to ampicillin						
	11	Health maintenance						

3

Interviewing and the Health History

The health history is a conversation with a purpose. Unlike social conversation, in which you express your own needs and interests with responsibility only for yourself, the primary goal of the clinician–patient interview is the well-being of the patient. The purpose of patient history taking is three-fold: to establish a trusting and supportive relationship, to gather information, and to offer information.

The interviewing process differs significantly from the format for the health history presented in Chapter 1. Both are fundamental to your work with patients but serve different purposes.

- The health history format is a structured framework for organizing patient information in written or verbal form for other health care providers.
- The interview that actually generates this information is more fluid. It requires knowledge of the information you need to obtain, the ability to elicit accurate and detailed information, and interpersonal skills that allow you to respond to the patient's feelings and elicit the patient's story in his or her own words.

For new patients in the office, hospital, or long-term care setting, you will do a *comprehensive health history*, described for adults in Chapter 1. For patients who seek care for a specific complaint, such as painful urination, a more limited interview, tailored to that specific problem—sometimes called a *focused* or *problem-oriented history*, may be indicated.

GETTING READY: THE APPROACH

Interviewing patients to elicit their health history requires planning.

- Take time for self-reflection. As clinicians, we encounter a wide variety of people, each one unique. Because we bring our own values, assumptions, and biases to every encounter, we must look inward to clarify how our expectations and reactions may affect what we hear and how we behave. Self-reflection brings a deepening personal awareness to our work with patients and is one of the most rewarding aspects of providing patient care.
- Review your clinical behavior and appearance. Consciously or not, you send messages through your behavior. Posture, gestures, eve contact, and tone of voice all can express interest, attention, acceptance, and understanding. The skilled interviewer seems calm and unhurried, even when time is limited. Reactions that betray disapproval, embarrassment, impatience, or boredom block communication. Patients find cleanliness, neatness, conservative dress, and a nametag reassuring.
- Review the chart. Before seeing the patient, review the medical record, or chart. It often provides valuable information about past diagnoses and treatments; however, data may be incomplete or even disagree with what you learn from the patient, so be open to developing new approaches or ideas.
- Adjust the environment. Always consider the patient's privacy. Pull shut any bedside curtains. Suggest moving to an empty room rather than having a conversation that can be overheard.
- Set goals for the interview. Clarify your goals for the interview. A clinician must balance provider-centered goals with patient-centered goals. The clinician's task is to balance these multiple agendas.
- Take notes. Jot down short phrases, specific dates, or words rather than trying to put them into a final format. Maintain good eye contact, and whenever the patient is talking about sensitive or disturbing material, put down your pen.

LEARNING ABOUT THE PA THE SEQUENCE OF THE INTERVIEW

In general, an interview moves through several stages. Throughout this sequence, you, as the clinician, must always stay attuned to the patient's feelings, help the patient express them, respond to their content, and validate their significance. Whether the interview is comprehensive or focused, be attentive to the patient's feelings and affect.

• Greet the patient and establish rapport. Greet the patient by name and introduce yourself, giving your name. If possible, shake hands. If this is the first contact, explain your role, including your status as a student and how you will be involved in the patient's care. Using a title to address the patient (e.g., Mr. O'Neil, Ms. Wu) is always best. Avoid first names unless you have specific permission from the patient.

Whenever visitors are present, maintain confidentiality. Let the patient decide if visitors or family members should remain in the room, and ask for the patient's permission before conducting the interview in front of them.

Attend to the patient's comfort. Ask how he or she is feeling and if you are coming at a convenient time. Look for signs of discomfort, such as frequent changes of position or facial expressions that show pain or anxiety. Arranging the bed may make the patient more comfortable.

Consider the best way to arrange the room. Choose a distance that facilitates conversation and good eye contact. Try to sit at eye level with the patient. Move any physical barriers between you and the patient, such as desks or bedside tables, out of the way.

Give the patient your undivided attention. Try not to look down to take notes or read the chart, and spend enough time on small talk to put the patient at ease.

• Establish an agenda. It is important to identify both your own and the patient's issues at the beginning of the encounter. Often, you may need to focus the interview by asking the patient which problem is most pressing. For

example, "Do you have some special concerns today? Which one are you most concerned about?" Some patients may not have a specific complaint or problem. *It is still important to start with the patient's story*.

• **Invite the patient's story.** As you probe the patient's concern, begin with **open-ended questions** that allow full freedom of response: "Tell me more about" Avoid questions that restrict the patient to a minimally informative "yes" or "no" answer. *Listen to the patient's answers without interrupting*.

Train yourself to *follow the patient's leads*. Use verbal and nonverbal cues that prompt patients to recount their stories spontaneously. Use *continuers*, especially at the outset, such as nodding your head and using phrases such as "Uh huh," "Go on," and "I see."

• Identify and respond to the patient's emotional cues. Patients offer various clues to their concerns that may be direct or indirect, verbal or nonverbal; they may express them as ideas or emotions. Acknowledging and responding to these clues help build rapport, expand the clinician's understanding of the illness, and improve patient satisfaction. A taxonomy of the clues about the patient's perspective on illness is provided in the box below.

CLUES TO THE PATIENT'S PERSPECTIVE ON ILLNESS

- Direct statement(s) by the patient of explanations, emotions, expectations, and effects of the illness
- Expression of feelings about the illness
- Attempts to explain or understand symptoms
- Speech clues (e.g., repetition, prolonged reflective pauses)
- Sharing a personal story
- Behavior clues indicative of unidentified concerns, dissatisfaction, or unmet needs such as reluctance to accept recommendations, seeking a second opinion, or early appointment

(Source: Lang F, Floyd MR, Beine KL. Clues to patients' explanations and concerns about their illnesses: a call for active listening. Arch Fam Med 9(3):222-227,2000.)

• Expand and clarify the patient's story. Each symptom has attributes that must be clarified, including context, associations, and chronology, especially for pain. It is critical to understand fully every symptom's essential characteristics. Always elicit the seven features of every symptom.

THE SEVEN ATTRIBUTES OF A SYMPTOM

- 1. Location. Where is it? Does it radiate?
- 2. Quality. What is it like?
- 3. Quantity or severity. How bad is it? (For pain, ask for a rating on a scale of 1–10.)
- 4. Timing. When did (does) it start? How long did (does) it last? How often did (does) it occur?
- 5. Setting in which it occurs. Include environmental factors, personal activities, emotional reactions, or other circumstances that may have contributed to the illness.
- 6. Remitting or exacerbating factors. Does anything make it better or worse?
- 7. Associated manifestations. Have you noticed anything else that accompanies it?

Use language that is understandable and appropriate to the patient. Technical language confuses patients and blocks communication. Whenever possible, use the patient's words, making sure you clarify their meaning.

Learn to facilitate the patient's story by using different types of questions and the techniques of skilled interviewing on pp. 43-45. Often you will need to use directed questions (see pp. 43–44) that ask for specific information the patient has not already offered. In general, the patient interview moves back and forth from an open-ended question to a directed question and then on to another open-ended question.

Establishing the sequence and time course of the patient's symptoms is important. You can encourage a chronologic account by asking such questions as "What then?" or "What happened next?"

• Generate and test diagnostic hypotheses. As you listen to the patient's concerns, you will begin to generate

42 Interviewing and the Health History

hypotheses about what disease process might be the cause. Identifying the various attributes of the patient's symptoms and pursuing specific details are fundamental to recognizing patterns of disease and differentiating one disease from another.

• Create a shared understanding of the problem. The disease/illness model helps you understand the difference between your perspective and the patient's perspective. In this model, disease is the explanation that the clinician brings to the symptoms. It is the way that the clinician organizes what he or she learns from the patient into a coherent picture that leads to a clinical diagnosis and treatment plan. Illness can be defined as how the patient experiences symptoms. The health history interview needs to include both of these views of reality.

Learning how patients perceive illness means asking patient-centered questions in the four domains listed below, which follow the mnemonic "FIFE"—Feelings, Ideas, effect on Function, and Expectations. This is crucial to patient satisfaction, effective health care, and patient follow-through.

EXPLORING THE PATIENT'S PERSPECTIVE

- The patient's Feelings, including fears or concerns, about the problem
- The patient's Ideas about the nature and the cause of the problem
- The effect of the problem on the patient's life and **F**unction
- The patient's Expectations of the disease, of the clinician, or of health care, often based on prior personal or family experiences
- Negotiate a plan. Learning about the disease and conceptualizing the illness give you and the patient the basis for planning further evaluation (physical examination, laboratory tests, consultations, etc.).
- Plan for follow-up and closing. Make sure the patient fully understands the plans you have developed together.

You can say, "We need to stop now. Do you have any questions about what we've covered?" Review future evaluation, treatments, and follow-up. Give the patient a chance to ask any final questions.

BUILDING A THERAPEUTIC RELATIONSHIP: THE TECHNIQUES OF SKILLED INTERVIEWING

Skilled interviewing requires the use of specific learnable techniques. Practice these techniques and find ways to be observed or recorded so that you can receive feedback on your progress.

Active Listening. This requires listening closely to what the patient is communicating, being aware of the patient's emotional state, and using verbal and nonverbal skills to encourage the speaker to continue and expand.

Guided Questioning. Learn to adapt your questioning to the patient's verbal and nonverbal cues.

Adaptive Questioning: Options for CLARIFYING THE PATIENT'S STORY

- Moving from open-ended to focused questions
- Questioning to elicit a graded response
- Asking a series of questions, one at a time
- Offering multiple choices for answers
- Clarifying what the patient means
- Using continuers
- Echoing

Proceed from the general to the specific. Directed questions should not be leading questions that call for a "yes" or "no" answer: not "Did your stools look like tar?" but "Please describe your stools."

44 Interviewing and the Health History

Ask questions that require a graded response rather than a single answer. "What physical activity do you do that makes you short of breath?" is better than "Do you get short of breath climbing stairs?" Be sure to ask one question at a time. Try "Do you have any of the following problems?" Be sure to pause and establish eye contact as you list each problem.

Sometimes patients seem unable to describe symptoms. *Offer multiple-choice answers.*

For patients using words that are ambiguous, request clarification, as in "Tell me exactly what you meant by 'the flu.'"

Posture, actions, or words encourage the patient to say more but do not specify the topic. Nod your head or remain silent. Lean forward, make eye contact, and use **continuers** like "Mm-hmm," "Go on," or "I'm listening."

Repetition and **echoing** of the patient's words encourage the patient to express both factual details and feelings.

Nonverbal Communication. Being sensitive to nonverbal messages allows you both to "read the patient" more effectively and to send messages of your own. Pay close attention to eye contact, facial expression, posture, head position and movement such as shaking or nodding, interpersonal distance, and placement of the arms or legs, such as crossed, neutral, or open. Physical contact (like placing your hand on the patient's arm) can convey empathy or help the patient gain control of feelings. You also can mirror the patient's *paralanguage*, or qualities of speech such as pacing, tone, and volume, to increase rapport. Be sensitive to cultural variations in uses and meanings of nonverbal behaviors.

Empathic Responses. Patients may express—with or without words—feelings they have not consciously acknowledged. *To empathize with your patient you must first identify his or her feelings.* Inquire about them rather than assuming how the patient feels.

Respond with understanding and acceptance. Responses may be as simple as "I understand," "That sounds upsetting," or "You seem sad." Empathy also may be nonverbal—for example, offering a tissue to a crying patient.

Validation. An important way to make a patient feel accepted is to provide verbal support that legitimizes or validates the patient's emotional experience.

Reassurance. Avoid premature or false reassurance. Such reassurance may block further disclosures, especially if the patient feels that exposing anxiety is a weakness. The first step to effective reassurance is identifying and accepting the patient's feelings without offering reassurance at that moment.

Partnering. Express your desire to work with patients in an ongoing way. Reassure patients that regardless of what happens with their disease, as their provider you are committed to a continuing partnership. Even in your role as a student, such support can make a big difference.

Summarization. Giving a capsule summary lets the patient know that you have been listening carefully. It also clarifies what you know and what you don't know. Summarization allows you to organize your clinical reasoning and to convey your thinking to the patient, which makes the relationship more collaborative.

Transitions. Tell patients when you are changing directions during the interview. This gives patients a greater sense of control.

Empowering the Patient. The clinician–patient relationship is inherently unequal. Patients have many reasons to feel vulnerable: pain, worry, feeling overwhelmed with the health care system, lack of familiarity with the clinical evaluation process. Differences of gender, ethnicity, race, or class may also create power differentials. Ultimately patients must be empowered to take care of themselves and feel confident about following through on your advice. Review the principles below.

Principles for Empowering the Patient

- Inquire about the patient's perspective.
- Express interest in the person, not just the problem.
- Follow the patient's lead.
- Flicit emotional content.
- Share information with the patient (e.g., transitions).
- Make your clinical reasoning transparent to the patient.
- Reveal the limits of your knowledge.

ADAPTING YOUR INTERVIEW TO SPECIFIC SITUATIONS

Always remember the importance of listening to the patient and clarifying the patient's agenda.

Silent Patient. Silence has many meanings and purposes. Watch closely for nonverbal cues such as difficulty controlling emotions. You may need to shift your inquiry to symptoms of depression or begin an exploratory mental status examination. Silence may be the patient's response to how you are asking questions. Are you asking too many direct questions? Have you offended the patient?

Confusing Patient. Some patients have *multiple symptoms* or a somatization disorder. Focus on the meaning or function of the symptoms and guide the interview into a psychosocial assessment. At other times you may be baffled, frustrated, and confused. The history is vague and difficult to understand, and patients may describe symptoms in bizarre terms. Try to learn more about the unusual symptoms. Watch for delirium in acutely ill or intoxicated patients and for dementia in the elderly. When you suspect a psychiatric or neurologic disorder, shift to a mental status examination, focusing on level of consciousness, orientation, and memory.

Patient With Altered Capacity. Some patients cannot provide their own histories because of delirium, dementia, or other conditions. Others cannot relate certain parts of the history. In such cases, determine whether the patient has decisionmaking capacity, or the ability to understand information related to health, to make medical choices based on reason and a consistent set of values and to declare preferences about treatments. Many patients with psychiatric or cognitive deficits still retain the ability to make decisions.

For patients with capacity, obtain their consent before talking about their health with others. Maintain confidentiality and clarify what you can discuss with others. Your knowledge about the patient can be quite comprehensive, yet others may offer surprising and important information. Consider dividing the interview into two segments—one with the patient and the other with both the patient and a second informant. Also learn the tenets of the Health Insurance Portability and

Accountability Act (HIPAA) passed by Congress in 1996, which sets strict standards for disclosure for both institutions and providers when sharing patient information. These can be found at www.hhs.gov/ocr/hipaa/.

For patients with impaired capacity, find a surrogate informant or decision-maker to assist with the history. Check whether the patient has a durable power of attorney for health care or a health care proxy. If not, in many cases a spouse or family member can represent the patient's wishes.

Talkative Patient. With a garrulous, rambling patient. several techniques are helpful. For the first 5 or 10 minutes, listen closely. Does the patient seem obsessively detailed or unduly anxious? Is there a flight of ideas or disorganized thought process? Try to focus on what seems most important to the patient. "You've described many concerns. Let's focus on the hip pain first. Can you tell me what it feels like?"

Crying Patient. Usually crying is therapeutic, as is quiet acceptance of the patient's distress. Make a facilitating or supportive remark like "I'm glad that you got that out."

Angry or Disruptive Patient. Many patients have reasons to be angry: they are ill, they have suffered a loss, they lack accustomed control over their own lives, and they feel relatively powerless. They may direct this anger toward you. Accept angry feelings from patients and allow them to express such emotions without getting angry in return. Validate their feelings without agreeing with their reasons. "I understand that you felt very frustrated by the long wait and answering the same questions over and over." Some angry patients become hostile and disruptive. Before approaching them, alert security. It is important to stay calm, appear accepting, and avoid being challenging. Keep your posture relaxed and nonthreatening. Once you have established rapport, gently suggest moving to a different location.

Patient With a Language Barrier. The ideal interpreter is a neutral, objective person trained in both languages and cultures. Avoid using family members or friends as interpreters: confidentiality may be violated. As you begin working with the interpreter, make questions clear, short, and simple. Speak directly to the patient. Bilingual written questionnaires are valuable.

GUIDELINES FOR WORKING WITH AN INTERPRETER

- Choose a professional interpreter in preference to a hospital worker, volunteer, or family member. Use the interpreter as a resource for cultural information.
- Orient the interpreter to the components you plan to cover in the interview; include reminders to translate everything the patient says.
- Arrange the room so that you and the patient have eye contact and can read each other's nonverbal cues.
- Seat the interpreter next to you (or even behind you) and allow the interpreter and the patient to establish rapport.
- Address the patient directly. Reinforce your questions with nonverbal behaviors.
- Keep sentences short and simple. Focus on the most important concepts to communicate.
- Verify mutual understanding by asking the patient to report back what he or she has heard.
- Be patient. The interview will take more time yet provide less information.

Patient With Low Literacy. Assess ability to read. Some patients may try to hide their reading problems. Ask the patient to read whatever instructions you have written. Simply handing the patient written material upside-down to see if the patient turns it around may settle the question.

Patient With Impaired Hearing. Find out the patient's preferred method of communicating. Patients may use American Sign Language, a unique language with its own syntax, or various other communication forms combining signs and speech. Determine whether the patient identifies with the Deaf or Hearing culture. Handwritten questions and answers may be the best solution. When patients have partial hearing impairment or can read lips, face them directly, in good light. If the patient has a unilateral hearing loss, sit on the hearing side. If the patient has a hearing aid, make sure it is working. Eliminate background noise such as television.

Patient With Impaired Vision. Shake hands to establish contact and explain who you are and why you are there. If the room is unfamiliar, orient the patient to the surroundings.

Patient With Limited Intelligence. Patients of moderately limited intelligence usually can give adequate histories. Pay special attention to the patient's schooling and ability to function independently. How far has the patient gone in school? If he or she didn't finish, why not? Assess simple calculations, vocabulary, memory, and abstract thinking. For patients with severe mental retardation, obtain the history from the family or caregivers. Avoid "talking down" or using condescending behavior. The sexual history is equally important and often overlooked.

Patient With Personal Problems. Patients may ask you for advice about personal problems outside the range of health. Letting the patient talk through the problem is usually more valuable and therapeutic than any answer you could give.

SENSITIVE TOPICS THAT CALL FOR SPECIFIC APPROACHES

The Sexual History. You can introduce questions about sexual function and practices at multiple points in a patient's history. An orienting sentence or two is often helpful. "Now I'd like to ask you some questions about your sexual health and practices" or "I routinely ask all patients about their sexual function."

- "When was the last time you had intimate physical contact with someone?" "Did that contact include sexual intercourse?"
- "Do you have sex with men, women, or both?" The health implications of heterosexual, homosexual, or bisexual experiences are significant.
- "How many sexual partners have you had in the last 6 months?" "In the last 5 years?" "In your lifetime?"
- Because no explicit risk factors may be present, it is important to ask all patients "Do you have any concerns about HIV disease or AIDS?" Also ask about routine use of condoms.

Mental Health History. Cultural constructs of mental illness vary widely, causing marked differences in acceptance and attitudes. Ask open-ended questions initially: "Have you ever had any problem with emotional or mental illnesses?" Then move to more specific questions: "Have you ever visited a counselor or psychotherapist?" "Have you taken medication for emotional issues?" "Have you or a family member ever been hospitalized for a mental health problem?"

Be sensitive to reports of mood changes or symptoms such as fatigue, tearfulness, appetite or weight changes, insomnia, and vague somatic complaints. Two opening screening questions are: "Over the past 2 weeks, have you felt down, depressed, or hopeless?" and "Over the past 2 weeks, have you felt little interest or pleasure in doing things?" Ask about thoughts of suicide: "Have you ever thought about hurting yourself or ending your life?" Evaluate severity.

Many patients with schizophrenia or other psychotic disorders can function in the community and tell you about their diagnoses, symptoms, hospitalizations, and medications. Investigate their symptoms and assess any effects on mood or daily activities.

The Alcohol and Drug History. Clinicians should routinely ask about current and past use of alcohol or drugs, patterns of use, and family history. "What do you like to drink?" or "Tell me about your use of alcohol" are good opening questions that avoid the easy yes or no response. The most widely used screening questions are the CAGE questions. Two or more affirmative answers to the CAGE questions suggest alcoholism.

THE CAGE QUESTIONS

- Have you ever felt the need to Cut down on drinking?
- Have you ever felt Annoyed by criticism of drinking?
- Have you ever felt Guilty about drinking?
- Have you ever taken a drink first thing in the morning (Eye-opener) to steady your nerves or get rid of a hangover?

(Adapted from Mayfield D, McLeod G, Hall P. The CAGE questionnaire: validation of a new alcoholism screening instrument. Am J Psychiatry 131:1121–1123, 1974.)

Also ask about blackouts (loss of memory for events during drinking), seizures, accidents or injuries while drinking, job loss, marital conflict, or legal problems. Ask specifically about drinking while driving or operating machinery.

Ouestions about drugs are similar. "How much marijuana do you use? Cocaine? Heroin? Amphetamines?" (Ask about each one by name.) "How about prescription drugs such as sleeping pills?" "Diet pills?" "Painkillers?" Use the CAGE questions but relate them to drug use. With adolescents, it may be helpful to ask about substance use by friends or family members first. "A lot of young people are using drugs these days. How about at your school? Your friends?"

Family Violence. Many authorities recommend routine screening of all female patients for domestic violence. Start with general "normalizing" questions: "Because abuse is common in many women's lives, I've begun to ask about it routinely." "Are there times in your relationships that you feel unsafe or afraid?" Consider physical abuse in the following settings:

- If injuries are unexplained, seem inconsistent with the patient's story, are concealed by the patient, or cause embarrassment
- If the patient has delayed getting treatment for trauma
- If there is a past history of repeated injuries or "accidents"
- If the patient or a person close to the patient has a history of alcohol or drug abuse
- If a partner tries to dominate the interview, will not leave the room, or seems unusually anxious or solicitous

Death and the Dying Patient. Work through your own feelings with the help of reading and discussion. Kubler-Ross has described five stages in our response to loss or the anticipatory grief of impending death: denial and isolation, anger, bargaining, depression or sadness, and acceptance. These stages may occur sequentially or overlap in different combinations.

Dying patients rarely want to talk about their illnesses all the time, nor do they wish to confide in everyone they meet. Give them opportunities to talk and then listen receptively, but be supportive if they prefer to stay at a social level.

52 Interviewing and the Health History

Understanding the patient's wishes about treatment at the end of life is an important clinician responsibility. Even if discussions of death and dying are difficult, you must learn to ask specific questions. Ask about Do Not Resuscitate (DNR) status. Find out about the patient's frame of reference. "What experiences have you had with the death of a close friend or relative?" "What do you know about cardiopulmonary resuscitation (CPR)?" Assure patients that relieving pain and taking care of their other spiritual and physical needs will be a priority.

Encourage any adult, but especially the elderly or chronically ill, to establish a *health care proxy*, an individual who can act for the patient in life-threatening situations.

Society and Ethics: Cultural Humility. As you provide care for an ever-expanding and diverse group of patients, it is important to understand how culture shapes not just the patient's beliefs, but your own. *Culture* is a system of shared ideas, rules, and meanings that influences how we view the world, experience it emotionally, and behave in relation to other people. This definition of culture is broader than the term *ethnicity*. The influence of culture is not limited to minority groups—it is relevant to everyone.

Clinicians are increasingly challenged to adopt *cultural humility*, a "process that requires humility as individuals continually engage in self-reflection and self-critique as lifelong learners and reflective practitioners." This process includes "the difficult work of examining cultural beliefs and cultural systems of both patients and providers to locate the points of cultural dissonance or synergy that contribute to patients' health outcomes." It calls for clinicians to "bring into check the power imbalances that exist in the dynamics of (clinician)-patient communication" and maintain mutually respectful and dynamic partnerships with patients and communities. The three-point framework below will help you.

• **Self-awareness.** As clinicians, we face the task of bringing our own values and biases to a conscious level. *Values* are the standards we use to measure our own and others' beliefs and behaviors. *Biases* are the attitudes or feelings that we attach to perceived differences, for example, the

way an individual relates to time, which can be a culturally determined phenomenon. Are you always on time—a positive value in the dominant Western culture? Or do you tend to run a little late? How do you feel about people whose habits are opposite to yours? Think about the role of physical appearance. Do you consider yourself thin, midsize, or heavy? How do you feel about people who have different weights?

• Enhanced Communication and Learning From the Patient. Maintain an open, respectful, and inquiring attitude. "What did you hope to get from this visit?" If you have established rapport and trust, patients will be willing to teach you. Be ready to acknowledge your ignorance or bias. "I mistakenly made assumptions about you that are not right. I apologize. Would you be willing to tell me more about yourself and your future goals?"

Learn about different cultures: do some reading; go to movies that are made in different countries; learn about different consumer health agendas.

• Collaborative Partnerships. Communication based on trust, respect, and a willingness to reexamine assumptions helps allow patients to express concerns that run counter to the dominant culture. You, the clinician, must be willing to listen to and validate these feelings, and not let your own feelings prevent you from exploring painful areas. You also must be willing to reexamine your beliefs.

Sexuality in the Clinician-Patient Relationship. The emotional and physical intimacy of the clinician-patient relationship may lead to sexual feelings. If you become aware of such feelings, accept them as a normal human response, and bring them to the conscious level so they will not affect your behavior. Denving these feelings makes it more likely that you will act inappropriately. Any sexual contact or romantic relationship with patients is *unethical*; keep your relationship with the patient within professional bounds and seek help if vou need it.

Ethical Considerations. Fundamental maxims are as follows:

• Nonmaleficence or primum non nocere, commonly stated as "First, do no harm"

54 Interviewing and the Health History

- *Beneficence*, or the dictum that the clinician needs to "do good" for the patient. As clinicians, our actions need to be motivated by what is in the patient's best interest.
- *Autonomy*, whereby patients have the right to determine what is in their own best interest
- *Confidentiality*, meaning that we are obligated not to tell others what we learn from our patients

The Tavistock Principles guide behavior in health care for both individuals and institutions

THE TAVISTOCK PRINCIPLES

Rights: People have a right to health and health care.

Balance: Care of individual patients is central, but the health of populations is also our concern.

Comprehensiveness: In addition to treating illness, we have an obligation to ease suffering, minimize disability, prevent disease, and promote health.

Cooperation: Health care succeeds only if we cooperate with those we serve, each other, and those in other sectors.

Improvement: Improving health care is a serious and continuing responsibility.

Safety: Do no harm.

Openness: Being open, honest, and trustworthy is vital in health care.

Beginning the Physical Examination: General Survey, Vital Signs, and Pain

THE HEALTH HISTORY

Common or Concerning Symptoms

- Changes in weight
- Fatigue and weakness
- Fever, chills, night sweats
- Pain

Changes in Weight. Good opening questions include "How often do you check your weight?" and "How is it compared to a year ago?"

- Weight gain occurs when caloric intake exceeds caloric expenditure over time. It also may reflect abnormal accumulation of body fluids.
- Weight loss has many causes: decreased food intake, dysphagia, vomiting, and insufficient supplies of food; defective absorption of nutrients; increased metabolic requirements; and loss of nutrients through the urine, feces, or injured skin. Be alert for signs of malnutrition.

Fatigue and Weakness. Fatigue is a nonspecific symptom with many causes. Use open-ended questions to explore the attributes of the patient's fatigue, and encourage the patient to fully describe what he or she is experiencing.

Weakness differs from fatigue. It denotes a demonstrable loss of muscle power and will be discussed later with other neurologic symptoms.

Fever, Chills, and Night Sweats. Ask about fever if the patient has an acute or chronic illness. Find out whether the patient has used a thermometer to measure the temperature. Distinguish between subjective *chilliness* and a *shaking chill*, with shivering throughout the body and chattering of teeth. *Night sweats* raise concerns about tuberculosis or malignancy.

Focus your questions on the timing of the illness and its associated symptoms. Become familiar with patterns of infectious diseases that may affect your patient. Inquire about travel, contact with sick people, or other unusual exposures. Be sure to inquire about medications, since they may cause fever. In contrast, recent ingestion of aspirin, acetaminophen, corticosteroids, and nonsteroidal anti-inflammatory drugs may mask it.

Pain. Approximately 70 million Americans report persisting or intermittent pain, often underassessed. Adopt the comprehensive approach found on pp. 67–69.

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Optimal weight, nutrition, and diet
- Exercise

Optimal Weight, Nutrition, and Diet. Less than half of U.S. adults maintain a healthy weight (BMI ≥19 but ≤25). Obesity has increased in every segment of the population. More than 50% of people with type 2 diabetes and roughly 20% of those with hypertension or elevated cholesterol levels are overweight or obese. Increasing obesity in children has been linked to rising rates of childhood diabetes.

Diet recommendations hinge on assessment of the patient's motivation and readiness to lose weight and individual risk factors. The *Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults* recommend the following general guidelines:

 A 10% weight reduction over 6 months, or a decrease of 300 to 500 kcal/day, for people with BMIs between 27 and 35 • A weight loss goal of ½ to 1 pound per week, because more rapid weight loss does not lead to better results at 1 vear

Exercise. Thirty minutes of moderate activity (defined as walking 2 miles in 30 minutes, or its equivalent, on most days of the week) is recommended. Patients can increase exercise by such simple measures as parking further away from their place of work or using stairs instead of elevators.

TECHNIQUES OF EXAMINAT

EXAMINATION TECHNIQUES

GENERAL SURVEY

Apparent State of Health

Acutely or chronically ill, frail, robust, vigorous

Level of Consciousness. Is the patient awake, alert, and interactive?

If not, promptly assess level of consciousness (see p. 333).

Signs of Distress

 Cardiac or respiratory distress

Clutching the chest, pallor, diaphoresis; labored breathing, wheezing, cough

Pain

Wincing, sweating, protecting painful area

Anxiety or depression

Anxious face, fidgety movements, cold and moist palms; inexpressive or flat affect, poor eye contact, psychomotor slowing

Skin Color and Obvious Lesions. See Chapter 6, The Skin, Hair, and Nails, for details.

Pallor, cyanosis, jaundice, rashes, bruises

Dress, Grooming, and Personal Hygiene

- Is the patient wearing any unusual jewelry? Where? Is there any body piercing?
- Note patient's hair, fingernails, and use of cosmetics

Facial Expression. Watch for eye contact. Is it natural? Sustained and unblinking? Averted quickly? Absent?

Stare of hyperthyroidism; flat or sad affect of depression.
Decreased eye contact may be cultural or may suggest anxiety, fear, or sadness.

Odors of Body and Breath.Odors can be important

Odors can be important diagnostic clues.

Breath odor of alcohol, acetone, uremia, or liver failure. Fruity odor of diabetes. (Never assume that alcohol on a patient's breath explains changes in mental status or neurologic findings.)

Posture, Gait, and Motor Activity

Preference to sit up in left-sided heart failure and to lean forward with arms braced in chronic obstructive pulmonary disease (COPD)

HEIGHT AND WEIGHT

Height. Measure the patient's height in stocking feet. Note the build—muscular or unconditioned, tall or short. Observe the body proportions.

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

Weight. Is the patient emaciated? Plump? If obese, is there central or dispersed distribution of fat? Weigh the patient with shoes off.

Calculate the *Body Mass Index* (BMI), which incorporates estimated but more accurate measurements of body fat than weight alone.

More than 50% of U.S. adults are overweight (BMI > 25); nearly 25% are obese (BMI > 30). These excesses are proven risk factors for diabetes, heart disease, stroke, hypertension, osteoarthritis, sleep apnea syndrome, and some forms of cancer.

Methods to Calculate Body Mass Index (BMI)								
Unit of Measure	Method of Calculation							
Weight in pounds, height in inches	1. Body Mass Index Chart (see p. 60)							
	2. $\frac{\left(\frac{\text{Weight (lbs)} \times 700^*}{\text{Height (inches)}}\right)}{\text{Height (inches)}}$							
Weight in kilograms, height in meters squared	3. $\frac{\text{Weight (kg)}}{\text{Height (m}^2)}$							
Either	4. BMI Calculator at Web site www. nhlbisupport.com/bmi/bmicalc.htm							

^{*}Several organizations use 704.5, but the variation in BMI is negligible. Conversion formulas: 2.2 lbs = 1 kg; 1.0 inch = 2.54 cm; 100 cm = 1 meter

(Source: National Institutes of Health and National Heart, Lung and Blood Institute. Body Mass Index Calculator. Available at: www.nhlbisupport.com/bmi/bmicalc.htm. Accessed December 12, 2007.)

 If the BMI falls above 25, engage the patient in a 24-hour dietary recall and compare the intake of food groups and number of servings per day with

current recommendations. Or choose a screening tool and provide appropriate counseling or referral.

 If the BMI falls below 17, be concerned about possible anorexia nervosa, bulimia, or other medical

Body Mass Index Chart

	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
Hei (inc	ght hes)	Body Weight (pounds) Overweight Obese															
58	91	96	100	105	110	115	119	124	129	134	138	143	148	153	158	162	167
59	94	99	104	109	114	119	124	128	133	138	143	148	153	158	163	168	173
60	97	102	107	112	118	123	128	133	138	143	148	153	158	163	168	174	179
61	100	105	111	116	122	127	132	137	143	148	153	158	164	169	174	180	185
62	104	109	115	120	126	131	136	142	147	153	158	164	169	175	180	185	191
63	107	113	118	124	130	135	141	145	152	158	163	169	174	180	185	191	197
64	110	116	122	128	134	140	145	151	157	163	169	174	180	185	192	197	204
65	114	120	126	132	138	144	150	156	162	168	174	180	186	192	198	204	210
66	118	124	130	136	142	148	155	161	167	173	179	186	192	198	204	210	216
67	121	127	134	140	146	153	159	166	172	178	185	191	198	204	211	217	223
68	125	131	138	144	151	158	164	171	177	184	190	197	203	210	216	223	230
69	128	135	142	149	155	162	169	176	182	189	196	203	209	216	223	230	236
70	132	139	146	153	160	167	174	181	188	189	196	203	209	216	223	230	236
71	136	143	150	157	165	172	179	186	193	200	208	215	222	229	236	243	250
72	140	147	154	162	169	177	184	191	199	206	213	221	228	235	242	250	258
73	144	151	159	166	174	182	189	197	204	212	219	227	235	242	250	257	265
74	148	155	163	171	179	186	194	202	210	218	225	233	241	249	256	264	272
75	152	160	168	176	184	192	200	208	216	224	232	240	248	256	264	272	279
76	156	164	172	180	189	197	205	213	221	230	238	246	254	263	271	279	287

(Source: National Institutes of Health and National Heart, Lung and Blood Institute. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults, June 1998.)

conditions (see Table 4-1, p. 70).

If the BMI is \leq 35, measure the waist circumference just above the hip bones. The patient may have excess body fat if the waist measures ≥40 inches for men.

THE VITAL SIGNS: BLOOD PRESSURE, HEART RATE, RESPIRATORY RATE, AND TEMPERATURE

Blood Pressure. To measure blood pressure accurately, choose a cuff of appropriate size and ensure careful technique.

SELECTING THE CORRECT BLOOD PRESSURE CUFF

- Width of the inflatable bladder of the cuff should be about 40% of upper-arm circumference (about 12–14 cm in the average adult).
- Length of inflatable bladder should be about 80% of upper-arm circumference (almost long enough to encircle the arm)

STEPS TO ENSURE ACCURATE **BLOOD PRESSURE RECORDINGS**

- Ideally, ask the patient to avoid smoking or drinking caffeinated beverages for 30 minutes before the blood pressure is taken and to rest for at least 5 minutes.
- Check to make sure the examining room is quiet and comfortably warm.
- Make sure the arm selected is free of clothing. There should be no arteriovenous fistulas for dialysis, scarring (continued)

STEPS TO ENSURE ACCURATE BLOOD PRESSURE RECORDINGS (CONTINUED)

from prior brachial artery cutdowns, or signs of lymphedema (seen after axillary node dissection or radiation therapy).

- Palpate the brachial artery to confirm that it has a viable pulse.
- Position the arm so that the brachial artery, at the antecubital crease, is at heart level—roughly level with the 4th interspace at its junction with the sternum.
- If the patient is seated, rest the arm on a table a little above the patient's waist; if standing, try to support the patient's arm at the midchest level.

MEASURING BLOOD PRESSURE

- Center the inflatable bladder over the brachial artery.
 The lower border of the cuff should be about 2.5 cm above the antecubital crease. Secure the cuff snugly. Position the patient's arm so that it is slightly flexed at the elbow.
- To determine how high to raise the cuff pressure, first estimate the systolic pressure by palpation. As you feel the radial artery with the fingers of one hand, rapidly inflate the cuff until the radial pulse disappears. Read this pressure on the manometer and add 30 mm Hg to it. Use of this sum as the target for subsequent inflations prevents discomfort from unnecessarily high cuff pressures. It also avoids the occasional error caused by an auscultatory gap—a silent interval between the systolic and diastolic pressures.
- Deflate the cuff promptly.
- Now place the bell of a stethoscope lightly over the brachial artery, taking care to make an air seal with its full rim. Because the sounds to be heard (Korotkoff sounds) are relatively low in pitch, they are heard better with the bell.

(continued)

MEASURING BLOOD PRESSURE (CONTINUED)

- Inflate the cuff rapidly again to the level just determined, and then deflate it slowly, at a rate of about 2 to 3 mm Hg per second. Note the level at which you hear the sounds of at least two consecutive beats. This is the systolic pressure.
- Continue to lower the pressure slowly. The disappearance point, usually only a few mm Hg below the muffling point. is the best estimate of diastolic pressure.
- Read both the systolic and diastolic levels to the nearest 2 mm Hg. Wait 2 or more minutes and repeat. Average your readings. If the first two readings differ by more than 5 mm Hg, take additional readings.
- Take blood pressure in both arms at least once.
- In patients taking antihypertensive medications or with a history of fainting, postural dizziness, or possible depletion of blood volume, take the blood pressure in two positions supine and standing (unless contraindicated). A fall in systolic pressure of 20 mm Hg or more, especially when accompanied by symptoms, indicates orthostatic (postural) hypotension.

In 2003, the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure (JNC) categorized four levels of systolic blood pressure (SBP) and diastolic blood pressure (DBP).

JNC VII Blood Pressure Classification—Adults Older Than 18 Years		
Category	Systolic (mm Hg)	Diastolic (mm Hg)
Normal	<120	<80
Prehypertension	120–139	80–89
Hypertension		
Stage 1	140-159	90–99
Stage 2	≥160	≥100

Note that the blood pressure goal for patients with hypertension, diabetes, or renal disease is <130/80.

When the systolic and diastolic levels fall in different categories, use the higher category. For example, 170/92 mm Hg is Stage 2 hypertension; 135/100 mm Hg is Stage 1 hypertension. In *isolated systolic hypertension*, systolic blood pressure is ≥140 mm Hg, and diastolic blood pressure is <90 mm Hg.

Heart Rate. The radial pulse is used commonly to assess heart rate. With the pads of your index and middle fingers, compress the radial artery until you detect a maximal pulsation. If the rhythm is regular, count the rate for 15 seconds and multiply by 4. If the rate is unusually fast or slow, count it for 60 seconds. When the rhythm is irregular, evaluate the rate by auscultation at the cardiac apex (the apical pulse).



Heart Rhythm. Feel the radial pulse. Check the rhythm again by listening with your stethoscope at the cardiac apex. Is the rhythm regular or irregular? If irregular, try to identify a pattern: (1) Do early beats appear in a basically regular rhythm? (2) Does the irregularity vary consistently with respiration? (3) Is the rhythm totally irregular?

Respiratory Rate and Rhythm. Observe the rate, rhythm, depth, and effort

Palpation of an irregularly irregular rhythm reliably indicates *atrial fibrillation*. For all other irregular patterns, an ECG is needed to identify the arrhythmia.

See Table 4-5, p. 74, Abnormalities in Rate and Rhythm of Breathing.

of breathing. Count the number of respirations in 1 minute either by visual inspection or by subtly listening over the patient's trachea with your stethoscope during examination of the head and neck or chest. Normally, adults take 14 to 20 breaths per minute in a quiet, regular pattern.

Temperature. Average oral temperature, usually 37°C (98.6°F), fluctuates considerably from the early morning to the late afternoon or evening. Rectal temperatures are higher than oral temperatures by about 0.4 to 0.5°C (0.7 to 0.9°F) but also vary.

In contrast, axillary temperatures are lower than oral temperatures by approximately 1° but take 5 to 10 minutes to register and are considered less accurate than other measurements.

Oral temperatures: Choose either glass or electronic thermometer.

Glass thermometer: Shake the thermometer down to 35°C (96°F) or below, insert it under the tongue, instruct the patient to close Fever or pyrexia refers to an elevated body temperature. Hyperpyrexia refers to extreme elevation in temperature, above 41.1°C (106°F), while hypothermia refers to an abnormally low temperature, below 35°C (95°F) rectally.

Causes of fever include infection, trauma (such as surgery or crush injuries), malignancy, blood disorders (such as acute hemolytic

EXAMINATION TECHNIQUES

both lips, and wait 3 to 5 minutes. Then read the thermometer, reinsert for 1 minute, and read it again. Avoid breakage.

anemia), drug reactions, and immune disorders such as collagen vascular disease.

Electronic thermometer: Carefully place the disposable cover over the probe and insert the thermometer under the tongue for about 10 seconds.

Tympanic membrane temperature: Make sure the external auditory canal is free of cerumen. Position the probe in the canal. Wait 2 to 3 seconds until the digital reading appears. This method measures core body temperature, which is higher than the normal oral temperature by approximately 0.8°C (11.4°F).

Rectal thermometer: Ask the patient to lie on one side with the hip flexed. Select a rectal thermometer with a stubby tip, lubricate it, and insert it about 3 cm to 4 cm (1½ inches) into the anal canal, in a direction pointing to the umbilicus. Remove it after 3 minutes, then read. Alternatively, use an electronic thermometer after lubricating the probe cover. Wait about 10 seconds for the digital temperature recording to appear.

The chief cause of hypothermia is exposure to cold. Other predisposing causes include reduced movement as in paralysis, interference with vasoconstriction as from sepsis or excess alcohol, starvation, hypothyroidism, and hypoglycemia. Older adults are especially suspectible to hypothermia and also less likely to develop fever.

ACUTE AND CHRONIC PAIN

The experience of pain is complex and multifactorial. It involves sensory, emotional, and cognitive processing but may lack a specific physical etiology.

Chronic pain is defined in several ways: pain not associated with cancer or other medical conditions that persists for more than 3 to 6 months; pain lasting more than 1 month beyond the course of an acute illness or injury; or pain recurring at intervals of months or years. Chronic noncancer pain affects 5% to 33% of patients in primary care settings.

Adopt a comprehensive approach, carefully listening to the patient's description of the many features of pain and contributing factors. Accept the self-report, which experts state is the most reliable indicator of pain.

Location. Ask the patient to point to the pain. Lay terms may not be specific enough to localize the site of origin.

Severity. Use a consistent method to determine severity. Three scales are common: the Visual Analog

EXAMINATION TECHNIQUES

Scale, and two scales using ratings from 1 to 10—the Numeric Rating Scale and the Faces Pain Scale.

Associated Features. Ask the patient to describe the pain and how it started. Pursue the seven features of pain, as you would with any symptom.

Attempted Treatments, Medications, Related Illnesses, and Impact on Daily Activities. Be sure to ask about any treatments that the patient has tried, including medications, physical therapy, and alternative medicines. A comprehensive medication history helps you to identify drugs that interact with analgesics and reduce their efficacy.

Identify any comorbid conditions such as arthritis, diabetes, HIV/AIDS, substance abuse, sickle cell disease, or psychiatric disorders. These can significantly affect the patient's experience of pain.

Health Disparities. Be aware of the well-documented health disparities in pain treatment and delivery of care, which range from lower use of analgesics in emergency rooms for

EXAMINATION TECHNIQUES

African-American and Hispanic patients to disparities in use of analgesics for cancer, postoperative, and low back pain. Clinician stereotypes, language barriers, and unconscious clinician biases in decision making all contribute to these disparities. Critique your own communication style, seek information and best practice standards, and improve your techniques of patient education and empowerment.

ECORDING YOUR FINDINGS

Record the vital signs taken at the time of your examination. They are preferable to those taken earlier in the day by other providers. (Common abbreviations for blood pressure, heart rate, and respiratory rate are self-explanatory.)

Recording the Physical Examination— General Survey and Vital Signs

"Mrs. Scott is a young, healthy-appearing woman, well-groomed, fit, and in good spirits. Height is 5'4", weight 135 lbs, BP 120/80, HR 72 and regular, RR 16. temperature 37.5°C."

OR

"Mr. Jones is an elderly male who looks pale and chronically ill. He is alert, with good eye contact, but cannot speak more than two or three words at a time because of shortness of breath. He has intercostal muscle retraction when breathing and sits upright in bed. He is thin, with diffuse muscle wasting. Height is 6'2", weight 175 lbs, BP 160/95, HR 108 and irregular, RR 32 and labored, temperature 101.2°F." (Suggests COPD exacerbation)

INTERPRETAT

Eating Disorders and Excessively Low BMI

Anorexia Nervosa

Refusal to maintain minimally normal body weight (or BMI above 17.5 kg/m²)

Fear of appearing fat

Frequently starving but in denial; lacking insight

Often brought in by family members May present as failure to make expected weight gains in childhood or adolescence, amenorrhea in women, loss of libido or potency in men

Associated with depressive symptoms such as depressed mood, irritability, social withdrawal, insomnia, decreased libido

Additional features supporting diagnosis: self-induced vomiting or purging, excessive exercise, use of appetite suppressants and/or diuretics

Biologic complications

- Neuroendocrine changes: amenorrhea, hormonal alterations
- Cardiovascular disorders: bradycardia, hypotension, dysrhythmias, cardiomyopathy
- Metabolic disorders: hypokalemia, hypochloremic metabolic alkalosis, increased BUN, edema

Other: dry skin, dental caries, delayed gastric emptying, constipation, anemia, osteoporosis

Bulimia Nervosa

Repeated binge eating followed by self-induced vomiting, misuse of laxatives, diuretics, or other medications; fasting; or excessive exercise

Often with normal weight

Overeating at least twice a week during 3-month period; large amounts of food consumed in short period (~2 hrs)

Preoccupation with eating; craving and compulsion to eat; lack of control over eating; alternating with periods of starvation

Dread of fatness but may be obese Subtypes of

- Purging: bulimic episodes accompanied by self-induced vomiting or use of laxatives, diuretics, or enemas
- Nonpurging: bulimic episodes accompanied by compensatory behavior such as fasting, exercise without purging
- Biologic complications; see changes listed for anorexia nervosa

(Sources: World Health Organization. The ICD-10 Classification of Mental and Behavioral Disorders: Diagnostic Criteria for Research. Geneva: World Health Organization, 1993; American Psychiatric Association. DSM-IV-TR: Diagnostic and Statistical Manual of Mental Disorders, 4th ed. Text Revision. Washington, DC: American Psychiatric Association, 2000. Halmi KA: Eating disorders: In: Kaplan HI, Sadock BJ, eds. Comprehensive Textbook of Psychiatry, 7th ed. Philadelphia: Lippincott Williams & Wilkins, 2000:1663–1676.)

Nutrition Screening Checklist

Yes (2 pts)
` ' /
Yes (3 pts)
Yes (2 pts)
Yes (2 pts)
Yes (2 pts)
Yes (4 pts)
Yes (1 pt)
Yes (1 pt)
Yes (2 pts)
Yes (2 pts)
TOTAL

Instructions: Check "yes" for each condition that applies, then total the nutritional score. For total scores between 3 and 5 points (moderate risk) or ≥6 points (high risk), further evaluation is needed (especially for the elderly).

Source: American Academy of Family Physicians. The Nutrition Screening Initiative. Available at: www.aafp.org/PreBuilt/NSI_DETERMINE.pdf. Accessed January 23, 2008.

Nutrition Counseling: Sources of Nutrients Nutrient **Food Source** Dairy foods such as yogurt, milk, and Calcium natural cheeses Breakfast cereal, fruit juice with calcium supplements Dark-green leafy vegetables such as collards, turnip greens Iron Shellfish Lean meat, dark turkey meat Cereals with iron supplements Spinach, peas, lentils Enriched and whole-grain bread **Folate** Cooked dried beans and peas Oranges, orange juice Dark-green leafy vegetables Vitamin D Milk (fortified) Eggs, butter, margarine Cereals (fortified)

(Source: Adapted from Dietary Guidelines Committee. 2000 Report. Nutrition and Your Health: Dietary Guidelines for Americans. Washington, DC: Agricultural Research Service, U.S. Department of Agriculture, 2000.)


Patients With Hypertension: Recommended Changes in Diet

Dietary Change	Food Source
Increase foods high in potassium	Baked white or sweet potatoes, cooked greens such as spinach Bananas, plantains, many dried fruits, orange juice
Decrease foods high in sodium	Canned foods (soups, tuna fish) Pretzels, potato chips, pickles, olives Many processed foods (frozen dinners, ketchup, mustard) Batter-fried foods Table salt, including for cooking

(Source: Adapted from Dietary Guidelines Committee. 2000 Report. Nutrition and Your Health: Dietary Guidelines for Americans. Washington, DC: Agricultural Research Service, U.S. Department of Agriculture.)



Abnormalities in Rate and Rhythm of Breathing



Normal. In adults, 14 to 20 per minute; in infants, up to 44 per minute

Rapid Shallow Breathing (Tachypnea).
Many causes, including restrictive lung disease, pleural chest pain, and an elevated diaphragm

MMM

Rapid Deep Breathing (Hyperpnea, Hyperventilation). Many causes, including exercise, anxiety, metabolic acidosis, brainstem injury. *Kussmaul breathing*, due to metabolic acidosis, is deep but rate may be fast, slow, or normal.

 \mathcal{M}

Slow Breathing. May be secondary to diabetic coma, drug-induced respiratory depression, increased intracranial pressure

Hyperpnea Apnea

Cheyne-Stokes Breathing. Rhythmically alternating periods of hyperpnea and apnea. In infants and the aged, may be normal during sleep; also accompanies brain damage, heart failure, uremia, drug-induced respiratory depression

Mh

Ataxic (Biot's) Breathing.
Unpredictable irregularity of depth and rate. Causes include brain damage and respiratory depression



Sighing Breathing. Breathing punctuated by frequent sighs. When associated with other symptoms, it suggests the hyperventilation syndrome. Occasional sighs are normal.

5

Behavior and Mental Status

Empathic listening and close observation open a unique vista on the patient's outlook, concerns, and habits. Nevertheless, clinicians often miss clues of mental illness and harmful dysfunctional behaviors in patients. The prevalence of mental disorders in the U.S. population is 30%, yet only approximately 20% of affected patients receive treatment. Even for patients who obtain care, evidence suggests that adherence to treatment guidelines in primary care offices is less than 50%.

Often patients have more than one mental disorder, with symptoms that mirror medical illnesses. Clinicians are well-advised to look for depression or anxiety in patients with substance abuse, and to look for substance abuse in patients with depression or anxiety. "Difficult patients" are frequently those with multiple unexplained symptoms and underlying psychiatric conditions that are amenable to therapy. Without better "dual diagnosis," patient health, function, and quality of life are at risk.

MENTAL HEALTH DISORDERS AND UNEXPLAINED SYMPTOMS IN PRIMARY CARE SETTINGS

Mental Health Disorders in Primary Care

- Approximately 20% of primary care outpatients have mental disorders, but up to 50% to 75% of these disorders are undetected and untreated.
- Prevalence of mental disorders in primary care settings is roughly:
 - Anxiety—20%
 - Mood disorders including dysthymia and depressive and bipolar disorders—25%

(continued)

MENTAL HEALTH DISORDERS AND UNEXPLAINED SYMPTOMS IN PRIMARY CARE SETTINGS (CONTINUED)

- Depression—10%
- Somatoform disorder—10% to 15%
- Alcohol and substance abuse—15% to 20%.

Explained and Unexplained Symptoms

- Physical symptoms account for approximately half of office visits.
- Roughly one-third of physical symptoms are unexplained; in 20% to 25% of patients, physical symptoms become chronic or recurring.
- In patients with unexplained symptoms, the prevalence of depression and anxiety exceeds 50% and increases with the total number of reported physical symptoms, making detection and "dual diagnosis" important clinical goals.

(Sources: Ansseau M, Dierick M, Buntinkxz F, et al. High prevalence of mental disorders in primary care. J Affect Disord 78(1):49-55, 2004; Kroenke K. The interface between physical and psychological symptoms. Primary Care Companion. | Clin Psychiatry, 5 (Suppl 7):11–18, 2003; Kroenke K. Patients presenting with somatic complaints: epidemiology, psychiatric comorbidity, and management. Int J Methods Psychiatr Res 12(1):34–43, 2003; Kroenke K, Sharpe M, Sykes R. Revising the classification of somatoform disorders: key questions and preliminary recommendations. Psychosomatics 48(4):277-285, 2007; Kroenke K, Spitzer RL, deGruy, et al. A symptom checklist for screen for somatoform disorders in primary care. Psychosomatics 39(3):263-272, 1998; Spitzer RL, Kroenke K, Williams JBW, et al. Validation and utility of a self-report version of PRIME-MD—the PHQ Primary Care Study. JAMA 282(18): 1737-1744, 1999; Staab JP, Datto CJ, Weinreig RM, et al. Detection and diagnosis of psychiatric disorders in primary medical care settings. Med Clin N Am 85(3):579-596, 2001.)

Unexplained conditions lasting beyond 6 weeks are increasingly recognized as common chronic disorders that should prompt screening for depression, anxiety, or both. Experts recommend brief screening questions with high sensitivity and specificity for patients at risk, followed by more detailed investigation when indicated. Several groups of patients warrant brief screening because of high rates of coexisting depression and anxiety.

PATIENT IDENTIFIERS FOR MENTAL HEALTH SCREENING

- Medically unexplained physical symptoms—more than half have a depressive or anxiety disorder
- Multiple physical or somatic symptoms or "high symptom count"
- High severity of the presenting somatic symptom
- Persisting pain
- Symptoms for more than 6 weeks
- Physician rating as a "difficult encounter"
- Recent stress
- Low self-rated health
- High use of health care services
- Substance abuse

(Source: Kroenke K. Patients presenting with somatic complaints: epidemiology, psychiatric comorbidity, and management. Int J Methods Psychiatr Res 12(1):34–43, 2003; Kroenke K. The interface between physical and psychological symptoms. Primary Care Companion. J Clin Psychiatry 5(Suppl 7):11–18, 2003.)

Common or Concerning Symptoms

- Changes in attention, mood, or speech
- Changes in insight, orientation, or memory
- Anxiety, panic, ritualistic behavior, and phobias
- Delirium or dementia

Your assessment of mental status begins with the first words of the interview. As you gather the health history, you will quickly discern the patient's level of alertness and orientation, mood, attention, and memory. You will learn about the patient's insight and judgment, as well as any recurring or unusual thoughts or perceptions. For some, you will need to conduct a more formal evaluation of mental status.

Many of the terms used to describe the mental status examination are familiar to you from social conversation. Take the time to learn their precise meanings in the context of the formal evaluation of mental status (see next page).

Assess level of consciousness, general appearance and mood, and ability to pay attention, remember, understand, and speak.

See Table 5-2, Disorders of Mood, pp. 88–89.

Assess the patient's responses to illness and life circumstances, which often tell you about his or her *insight* and *judgment*. Test *orientation* and *memory*.

TERMINOLOGY:	THE MENTAL STATUS EXAMINATION
Level of consciousness Attention	Alertness or state of awareness of the environment The ability to focus or concentrate over time on one task or activity
Memory	The process of registering or recording information. <i>Recent or short-term memory</i> covers minutes, hours, or days; <i>remote or long-term memory</i> refers to intervals of years.
Orientation	Awareness of personal identity, place, and time; requires both memory and attention
Perceptions	Sensory awareness of objects in the envi- ronment and their interrelationships; also refers to internal stimuli (e.g., dreams)
Thought processes	The logic, coherence, and relevance of the patient's thoughts, or <i>how</i> people think
Thought content	What the patient thinks about, including level of insight and judgment
Insight	Awareness that symptoms or disturbed behaviors are normal or abnormal
Judgment	Process of comparing and evaluating alternatives; reflects values that may or may not be based on reality and social conventions or norms
	(continued)

TERMINOLOGY:	THE M	ENTAL
STATUS EXAMIN	ATION	(CONTINUED)

Affect	An observable, usually episodic, feeling
	tone expressed through voice, facial expression, and demeanor
Mood	A more sustained emotion that may color a
	person's view of the world (mood is to
	affect as climate is to weather)
Language	A complex symbolic system for expressing, receiving, and comprehending words;
	essential for assessing other mental functions
Higher cognitive	Assessed by vocabulary, fund of informa-
functions	tion, abstract thinking, calculations,
	construction of objects with two or
	three dimensions

Explore any unusual thoughts, preoccupations, beliefs, or perceptions as they arise during the interview.

All patients with documented or suspected brain lesions, psychiatric symptoms, or reports from family members of vague or changed behavioral symptoms need further systematic assessment.

See Table 5-3, Anxiety Disorders, pp. 90–91, and Table 5-4, Selected Psychotic Disorders, p. 92.

See Table 20-1, Delirium and Dementia, pp. 424–425.

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Screening for depression and suicidality
- Screening for dementia

Depression. Lifetime prevalence of *major depression* meeting formal diagnostic criteria in the United States is approximately

80 Behavior and Mental Status

16%, with an annual prevalence of approximately 6%. Primary care providers fail to diagnose major depression in up to 50% of affected patients, often missing early clues such as *low self-esteem, anhedonia* (lack of pleasure in daily activities), *sleep disorders*, and *difficulty concentrating* or *making decisions*. Failure to diagnose depression can have fatal consequences—suicide rates in patients with major depression are eight times higher than in the general population. Ask, "Over the past 2 weeks, have you felt down, depressed, or hopeless?" and "Over the past 2 weeks, have you felt little interest or pleasure in doing things?"

Suicide. Suicide rates are highest among men older than 85 years and are increasing among teenagers and young adults. More than half of patients committing suicide have visited their physicians in the prior month. More than 90% of suicide deaths occur in patients with depression or other mental health disorders or substance abuse. Risk factors include suicidal or homicidal ideation, intent, or plan; access to the means for suicide; current symptoms of psychosis or severe anxiety; any history of psychiatric illness (especially linked to a hospital admission); substance abuse; personality disorder; and prior history or family history of suicide. Patients with these risk factors should be immediately referred for psychiatric care and possibly hospitalization.

Alcohol and Substance Abuse. The interactions and comorbidity of alcohol and substance abuse with mental disorders and suicide are extensive. Alcohol, tobacco, and illicit drugs account for more illness, deaths, and disabilities than any other preventable condition. Lifetime prevalence of alcohol and illicit drug use in the United States is 13% and 3%. In recent U.S. surveys, 8% of those 12 years or older, or 19 million people, reported use of illicit drugs in the prior 30 days. An estimated 3% are dependent on or abuse illicit drugs; of these, 60% use marijuana. Because screening for alcohol and drug use is part of *every* patient history, information on screening is found in Chapter 3, Interviewing and the Health History.

TECHNIQUES OF EXAMINATION

Important Areas of Examination

- Appearance and behavior
- Speech and language
- Mood
- Thoughts and perceptions
- Cognition, including memory, attention, information and vocabulary, calculations, abstract thinking, and constructional ability

Observe patient's mental status throughout your interaction. Test specific functions if indicated during the interview or physical examination.

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

APPEARANCE AND BEHAVIOR

Assess the following:

- Level of Consciousness.
 Observe alertness and response to verbal and tactile stimuli.
- Posture and Motor Behavior.
 Observe pace, range, character, and appropriateness of movements.
- Dress, Grooming, and Personal Hygiene
- Facial Expressions. Assess during rest and interaction.
- Manner, Affect, and Relation to People and Things

Normal consciousness, lethargy, obtundation, stupor, coma (see pp. 332–333)

Restlessness, agitation, bizarre postures, immobility, involuntary movements

Fastidiousness, neglect

Anxiety, depression, elation, anger, responses to imaginary people or objects, withdrawal

SPEECH AND LANGUAGE

Note quantity, rate, loudness, clarity, and fluency of speech. If indicated, test for aphasia.

Aphasia, dysphonia, dysarthria, changes with mood disorders

Testing for Aphasia	
Word comprehension	Ask patient to follow a one-stage command, such as "Point to your nose." Try a two-stage command: "Point to your mouth, then your knee."
Repetition	Ask patient to repeat a phrase of one-syllable words (the most difficult repetition task): "No ifs, ands, or buts."
Naming	Ask patient to name the parts of a watch.
Reading comprehension	Ask patient to read a paragraph aloud.
Writing	Ask patient to write a sentence.

MOOD

Ask about the patient's spirits. Note nature, intensity, duration, and stability of any abnormal mood. If indicated, assess risk of suicide.

Happiness, elation, depression, anxiety, anger, indifference

THOUGHT AND PERCEPTIONS

Thought Processes. Assess logic, relevance, organization, and coherence.

Derailments, flight of ideas, incoherence, confabulation, blocking

Thought Content. Ask about and explore any unusual or unpleasant thoughts.

Obsessions, compulsions, delusions, feelings of unreality

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

Perceptions. **Ask** about any unusual perceptions (e.g., seeing or hearing things).

Illusions, hallucinations

Insight and Judgment.

Assess patient's insight into the illness and level of judgment used in making decisions or plans.

Recognition or denial of mental cause of symptoms; bizarre, impulsive, or unrealistic judgment

COGNITIVE FUNCTIONS

If indicated, assess:

Orientation to time, place, and person

Disorientation

Attention

- Digit span—ability to repeat a series of numbers forward and then backward
- Serial 7s—ability to subtract 7 repeatedly, starting with 100
- Spelling backward of a five-letter word, such as W-O-R-L-D

Remote Memory (e.g., birthdays, anniversaries, social security number, schools, jobs, wars)

Recent Memory (e.g., events of the day)

New Learning Ability ability to repeat three or four words after a few minutes of unrelated activity Poor performance of digit span, serial 7s, and spelling backward are common in dementia and delirium but have other causes, too.

Impaired in late stages of dementia

Recent memory and new learning ability impaired in dementia, delirium, and amnestic disorders

HIGHER COGNITIVE FUNCTIONS

If indicated, assess:

Information and Vocabulary.

Note range and depth of patient's information, complexity of ideas expressed, and vocabulary used. For the fund of information, you also may ask names of presidents, other political figures, or large cities.

These attributes reflect intelligence, education, and cultural background. They are limited by mental retardation but are fairly well preserved in early dementia.

Calculating Abilities, such as addition, subtraction, and multiplication

Poor calculation in mental retardation and dementia

Abstract Thinking—ability to respond abstractly to questions about

• The meaning of *proverbs*, such as "A stitch in time

saves nine"

- Concrete responses (observable details rather than concepts) are common in mental retardation, dementia, and delirium. Responses are sometimes bizarre in schizophrenia.
- The *similarities* of beings or things, such as a cat and a mouse or a piano and a violin

Constructional Ability. Ask patient

- To copy figures such as circle, cross, diamond, and box, and two intersecting pentagons, or
- To draw a clock face with numbers and hands

Impaired ability common in dementia and with parietal lobe damage

SPECIAL TECHNIQUE

Mini-Mental State Examination (MMSE). This brief test is useful in screening for cognitive dysfunction and dementia and following their course over time. For more detailed information regarding the MMSE, contact the Publisher, Psychological Assessment Resources, Inc., 16204 North Florida Avenue, Lutz, Florida 33549. Below are some sample questions.

MMSE SAMPLE ITEMS

Orientation to Time

"What is the date?"

Registration

"Listen carefully; I am going to say three words. You say them back after I stop. Ready? Here they are . . .

HOUSE (pause), CAR (pause), LAKE (pause). Now repeat those words back o me." [Repeat up to five times, but score only the first trial.]

Naming

"What is this?" [Point to a pencil or pen.]

Reading

"Please read this and do what it says." [Show examinee the words on the stimulus form.1

CLOSE YOUR EYES

Reproduced by special permission of the Publisher, Psychological Assessment Resources, Inc., 16204 North Florida Avenue, Lutz, Florida 33549, from the Mini Mental State Examination, by Marshal Folstein and Susan Folstein, Copyright 1975, 1998, 2001 by Mini Mental LLC, Inc. Published 2001 by Psychological Assessment Resources, Inc. Further reproduction is prohibited without permission of PAR, Inc. The MMSE can be purchased from PAR, Inc. by calling (800) 331-8378 or (813) 968-3003.

RECORDING YOUR FINDINGS

Recording the Examination—Mental Status

"Mental Status: The patient is alert, well-groomed, and cheerful. Speech is fluent and words are clear. Thought processes are coherent, insight is good. The patient is oriented to person, place, and time. Serial 7s accurate; recent and remote memory intact. Calculations intact."

OR

"Mental Status: The patient appears sad and fatigued; clothes are wrinkled. Speech is slow and words are mumbled. Thought processes are coherent but insight into current life reverses is limited. The patient is oriented to person, place, and time. Digit span, serial 7s, and calculations accurate but responses delayed. Clock drawing is good."

TABLE 5-1 Somatoform Disorders: Types and Approach

Management Guidelines for Patients With Unexplained Medical Symptoms

10

General Aspects	Show empathy and understanding for the complaints and frustrating experiences the patient has had so far (e.g., explain that medically unexplained symptoms are common). Develop a good patient—physician relationship; try to be the "coordinator" of diagnostic procedures and care.
Diagnosis	Explore not only the history of complaints and former treatments, but impairment, (health) anxiety, psychosocial issues. Use screeners and self-report questionnaires as economic instruments for detection; use symptom diaries to assess course and influencing factors on symptoms. When the patient presents with a new symptom, examine the relevant organ system. Show the results of investigations to explain the absence of pathology and to give clear reassurance that there is no serious physical disease. Avoid unnecessary diagnostic tests or surgical procedures.
Treatment	Provide regularly scheduled visits (e.g., every 4–6 weeks), especially in the case of a history of very frequent health care utilization. Explain that treatment is coping, not curing (when pathology cannot be found or does not

If referral is necessary to start psychotherapy or psychopharmacotherapy, prepare the patient for the treatment and reassure him/her that you will continue to be his/her "doctor." Suggest coping strategies like regular physical activity, relaxation, distraction.

explain degree of complaints).

(Source: Reif W, Martin A, Rauh E, et al. Evaluation of general practitioners' training: how to manage patients with unexplained physical symptoms. Psychosomatics 47(4):304–311, 2006.)

TABLE 5-2 Disorders of Mood

Major Depressive Episode

At least five of the symptoms listed below (including one of the first two) must be present during the same 2-week period; they must represent a change from the person's previous state.

- Depressed mood (may be an irritable mood in children and adolescents) most of the day, nearly every day
- Markedly diminished interest or pleasure in almost all activities most of the day, nearly every day
- Significant weight gain or loss (not dieting) or increased or decreased appetite nearly every day
- Insomnia or hypersomnia nearly every day
- Psychomotor agitation or retardation nearly every day
- Fatigue or loss of energy nearly every day
- Feelings of worthlessness or inappropriate guilt nearly every day
- Inability to think or concentrate or indecisiveness nearly every day
- Recurrent thoughts of death or suicide, or a specific plan for or attempt at suicide

The symptoms cause significant distress or impair social, occupational, or

Manic Episode

- A distinct period of abnormally and persistently elevated, expansive, or irritable mood must be present for at least a week (any duration if hospitalization is necessary). During this time, at least three of the symptoms listed below have been persistent and significant. (Four symptoms are required if the mood is only irritable.)
- Inflated self-esteem or grandiosity
- Decreased need for sleep (feels rested after sleeping 3 hours)
- More talkative than usual or pressure to keep talking
- Flight of ideas or racing thoughts
- Distractibility
- Increased goal-directed activity (either socially at work or school, or sexually) or psychomotor agitation
- Excessive involvement in pleasurable high-risk activities (buying sprees, foolish business ventures, sexual indiscretions)

The disturbance is severe enough to impair social or occupational functions or

(table continues next page)

TABLE 5-2 Disorders of Mood (continued)

other important functions. In severe cases, hallucinations and delusions may occur. relationships. It may necessitate hospitalization for the protection of self or others. In severe cases, hallucinations and delusions may occur.

Mixed Episode

A mixed episode, which must last at least 1 week, meets the criteria for both major and manic depressive episodes.

Hypomanic Episode

The mood and symptoms resemble those in a manic episode but are less impairing, do not require hospitalization, do not include hallucinations or delusions, and have a shorter minimum duration—4 days.

Dysthymic Disorder

A depressed mood and symptoms for most of the day, for more days than not, over at least 2 years (1 year in children and adolescents). Freedom from symptoms lasts no more than 2 months at a time.

Cyclothymic Episode

Numerous periods of hypomanic and depressive symptoms that last for at least 2 years (1 year in children and adolescents). Freedom from symptoms lasts no more than 2 months at a time.

(Tables 5-2, 5-3, and 5-4 are based, with permission, on the Diagnostic and Statistical Manual of Mental Disorders, Fourth ed., Text Revision [DSM IV-TR]. Washington, DC: American Psychiatric Association, 2000. For further details and criteria, the reader should consult this manual, its successor, or comprehensive textbooks of psychiatry.)

Anxiety Disorders

Panic Disorder. Recurrent, unexpected panic attacks, at least one of which has been followed by a month or more of persistent concern about further attacks, worry over their implications or consequences, or a significant change in behavior in relation to the attacks.

A panic attack is a discrete period of intense fear or discomfort that develops abruptly and peaks within 10 minutes. It involves at least four of the following symptoms: (1) palpitations, pounding heart, or accelerated heart rate; (2) sweating; (3) trembling or shaking; (4) shortness of breath or a sense of smothering; (5) a feeling of choking; (6) chest pain or discomfort; (7) nausea or abdominal distress; (8) feeling dizzy, unsteady, lightheaded, or faint; (9) feelings of unreality or depersonalization; (10) fear of losing control or going crazy; (11) fear of dying; (12) paresthesias (numbness or tingling); and (13) chills or hot flushes.

Agoraphobia. Anxiety about being in places or situations where escape may be difficult or embarrassing or help for sudden symptoms unavailable. Such situations are avoided, require a companion, or cause marked anxiety.

Specific Phobia. A marked, persistent, and excessive or unreasonable fear that is cued by the presence or anticipation of a specific object or situation, such as dogs, injections, or flying. The person recognizes the fear as excessive or unreasonable, but exposure to the cue provokes immediate anxiety. Avoidance or fear impairs the person's normal routine, occupational or academic functioning, or social activities or relationships.

Social Phobia. A marked, persistent fear of one or more social or performance situations that involve exposure to unfamiliar people or to scrutiny by others. Those afflicted fear that they will act in embarrassing or humiliating ways, as by showing their anxiety. Exposure creates anxiety and possibly a panic attack, and the person avoids precipitating situations. He or she recognizes the fear as excessive or unreasonable. Normal functioning, social activities, or relationships are impaired.

Obsessive-Compulsive Disorder. Obsessions or compulsions that cause marked anxiety or distress. Although recognized as

TABLE 5-3 Anxiety Disorders (continued)

excessive or unreasonable, they are time consuming and interfere with the person's normal routine and relationships.

Acute Stress Disorder. Exposure to a traumatic event that involved actual or threatened death or serious injury to self or others, leading to intense fear, helplessness, or horror. During or immediately after this event, the person has at least three dissociative symptoms: (1) a subjective sense of numbing, detachment, or absence of emotional responsiveness; (2) a reduced awareness of surroundings, as in a daze; (3) feelings of unreality; (4) feelings of depersonalization; and (5) amnesia for an important part of the event. The event is persistently reexperienced, as in thoughts, images, dreams, illusions, and flashbacks. The person is anxious, shows increased arousal, and avoids stimuli that evoke memories of the event. Causes marked distress or impairs social, occupational, or other important functions. Symptoms occur within 4 weeks of the event and last from 2 days to 4 weeks.

Posttraumatic Stress Disorder. The event, fearful response, and persistent reexperiencing of the traumatic event resemble acute stress disorder. Hallucinations may occur. The person has increased arousal, tries to avoid stimuli related to the trauma, and has numbing of general responsiveness. Causes marked distress and impaired social or occupational function, and lasts for more than a month.

Generalized Anxiety Disorder. Lacks a specific traumatic event or focus for concern. Excessive anxiety and worry are hard to control and generalize to a number of events or activities. At least three of the following symptoms are associated: (1) feeling restless, keyed up, or on edge; (2) being easily fatigued; (3) difficulty in concentrating or mind going blank; (4) irritability; (5) muscle tension; and (6) difficulty in falling or staying asleep, or restless, unsatisfying sleep. Causes significant distress or impairs daily function.

Selected Psychotic Disorder

Schizophrenia. Impairs major functioning at work or school, in interpersonal relations, or in self-care. Performance of one or more of these functions must decrease for a significant time to a level markedly below prior achievement. Person displays at least two of the following for a significant part of 1 month: (1) delusions; (2) hallucinations; (3) disorganized speech; (4) grossly disorganized or catatonic behavior; and (5) negative symptoms such as a flat affect, alogia (lack of content in speech), or avolition (lack of interest, drive, and ability to set and pursue goals). Continuous signs of the disturbance must persist for at least 6 months.

Subtypes of this disorder include paranoid, disorganized, and catatonic schizophrenia.

Schizophreniform Disorder. Symptoms are similar to those of schizophrenia but last <6 months. Functional impairment need not be present.

Schizoaffective Disorder. Features both a major mood disturbance and schizophrenia. Mood disturbance (depressive, manic, or mixed) present during most of the illness and must, for a time, be concurrent with symptoms of schizophrenia and demonstrate delusions or hallucinations for at least 2 weeks without prominent mood symptoms.

Delusional Disorder. Nonbizarre delusions involve situations in real life, such as having a disease, and persists for at least a month. Functioning is not markedly impaired and behavior is not obviously odd or bizarre. Symptoms of schizophrenia, except for tactile and olfactory hallucinations, are not present. Brief Psychotic Disorder. At least one of the following psychotic symptoms must be present: delusions, hallucinations, disordered speech such as frequent derailment or incoherence, or grossly disorganized or catatonic behavior. Disturbance lasts ≥1 day but <1 month, and person returns to prior functional level.

The Skin, Hair, and Nails

THE HEALTH HISTORY

Common or Concerning Symptoms

- Hair loss
- Rash
- Moles

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Risk factors for melanoma
- Avoidance of excessive sun exposure

Counsel patient to avoid unnecessary sun exposure and to use sunscreen with at least SPF-15. Teach the ABCD screen for dysplastic nevi/melanomas: Asymmetry, irregular Borders, variation in Color, and Diameter >6 mm. Survey skin at 3-year intervals for patients 20 to 40 years of age and annually for patients older than 40 years.

TECHNIQUES OF EXAMINATION

EXAMINATION TECHNIQUES

OSSIBLE FINDINGS

SKIN

Examine the entire skin surface under good lighting. Inspect and palpate each area.

Note:

 Color
 Cyanosis, jaundice, carotenemia, changes in melanin

Moisture Dry, oily

Temperature Cool, warm

• Texture Smooth, rough

 Mobility—ease with which a fold of skin can be moved

Decreased in edema

 Turgor—speed with which the fold returns into place

Decreased in dehydration

Note any lesions and their

Anatomical location and distribution

Generalized, localized

Patterns and shapes
 Linear, clustered, dermatomal

 Type Macule, papule, pustule, bulla, tumor

Color
 Red, white, brown, heliotrope

HAIR

Inspect and **palpate** the hair.

Note

Quantity Thin, thick

 Distribution Patchy or total alopecia

 Texture Fine, coarse

NAILS

Inspect and **palpate** the fingernails and toenails.

Note

Color Cyanosis, pallor

Shape Clubbing

Any lesions Paronychia, onycholysis

RECORDING YOUR FINDINGS

Recording the Physical Examination— The Skin, Hair, and Nails

"Color good. Skin warm and moist. Nails without clubbing or cyanosis. No suspicious nevi, rash, petechiae, or ecchymoses."

pigments)

AIDS TO INTERPRETATION

TABLE 6-1 Color Changes	in the Skin
Color/Mechanism	Selected Causes
Brown: Increased melanin (greater than a person's genetic norm)	Sun exposure Pregnancy (melasma) Addison's disease
Blue (cyanosis): Increased deoxyhemoglobin from hypoxia: • Peripheral • Central (arterial)	Anxiety or cold environment Heart or lung disease
Abnormal hemoglobin	Methemoglobinemia, sulfhemoglobinemia
 Red: Increased visibility of oxyhemoglobin from: Dilated superficial blood vessels or increased blood flow in skin Decreased use of oxygen in skin 	Fever, blushing, alcohol intake, local inflammation Cold exposure (e.g., cold ears)
Yellow: Increased bilirubin of jaundice (sclera looks yellow)	Liver disease, hemolysis of red blood cells
Carotenemia (sclera does not look yellow)	Increased carotene intake from yellow fruits and vegetables
Pale: Decreased melanin	Albinism, vitiligo, tinea versicolor
Decreased visibility of oxyhemoglobin from: • Decreased blood flow to skin • Decreased amount of oxyhemoglobin	Syncope or shock Anemia
Edema (may mask skin	Nephrotic syndrome

TABLE 6-2 **Primary Skin Lesions**

Flat, Nonpalpable Lesions With Changes in Skin Color



Macule—Small flat spot, up to 1.0 cm



Examples: Hemangioma, vitiligo



Patch-Flat spot, 1.0 cm or larger Example: Café-au-lait spot

Palpable Elevations: Solid Masses



Papule—Up to 1.0 cm Example: An elevated nevus



Plaque—Elevated superficial lesion 1.0 cm or larger, often formed by coalescence of papules

Primary Skin Lesions (continued)




Example: Psoriasis



Nodule-Marble-like lesion larger than 0.5 cm, often deeper and firmer than a papule Example: Dermatofibroma



Cyst—Nodule filled with expressible material, either liquid or semisolid Example: Epidermal inclusion cyst



Wheal—A somewhat irregular, relatively transient, superficial area of localized skin edema Examples: Mosquito bite, hives (urticaria)

TABLE 6-2 Primary Skin Lesions (continued)

Palpable Elevations With Fluid-Filled Cavities



Vesicle—Up to 1.0 cm; filled with serous fluid



Example: Herpes simplex



Bulla—1.0 cm or larger; filled with serous fluid



Example: Insect bite



Pustule—Filled with pus

Primary Skin Lesions (continued)



Examples: Acne, small pox (impetigo)



Burrow (scabies)—A minute, slightly raised tunnel in the epidermis, commonly found on the finger webs and on the sides of the fingers. It looks like a short (5-15 mm), linear or curved gray line and may end in a tiny vesicle. With a magnifying lens, look for the burrow of the mite that causes scabies.

Secondary Skin Lesions

May arise from primary lesions, overtreatment, excess scratching



Scale—A thin flake of dead, exfoliated epidermis



Examples: Ichthyosis, dandruff, dry skin, psoriasis

Secondary Skin Lesions (continued)



Crust—The dried residue of skin exudates such as serum, pus, or blood Example: Impetigo



Lichenification—Visible and palpable thickening of the epidermis and roughening of the skin with increased visibility of the normal skin furrows (often from chronic rubbing) Example: Neurodermatitis Scars—Connective tissue that arises from injury or disease



Example: Acne



Keloids—Hypertrophic scarring that extends beyond the borders of the initiating injury

(Sources of photos: Hemangioma, Café-au-Lait Spot, Elevated Nevus, Psoriasis [bottom], Dermatofibroma, Herpes Simplex, Insect Bite [bottom], Impetigo, Lichenification—Hall JC. Sauer's Manual of Skin Diseases, 9th ed. Philadelphia: Lippincott Williams & Wilkins, 2006; Vitiligo, Psoriasis [top], Epidermal Inclusion Cyst, Urticaria, Insect Bite [top], Acne, Ichthyosis, Psoriasis, Acne Scar, Keloids-Goodheart HP. Goodheart's Photoguide of Common Skin Disorders: Diagnosis and Management, 2nd ed. Philadelphia: Lippincott Williams & Wilkins, 2003; Small Pox—Ostler HB, Mailbach HI, Hoke AW, Schwab IR. Diseases of the Eye and Skin: A Color Atlas. Philadelphia: Lippincott Williams & Wilkins, 2004.)

Secondary Skin Lesions—Depressed



Erosion—Nonscarring loss of the superficial epidermis; surface is moist but does not bleed Example: Aphthous stomatitis, moist area after the rupture of a vesicle, as in chickenpox



Excoriation—Linear or punctate erosions caused by scratching Example: Cat scratches



Fissure—A linear crack in the skin, often resulting from excessive dryness Example: Athlete's foot



Ulcer—A deeper loss of epidermis and dermis; may bleed and scar Examples: Stasis ulcer of venous insufficiency, syphilitic chancre

^{*}These are secondary lesions (resulting from primary lesions). (Sources of photos: Erosion, Excoriation, Fissure—Goodheart HP. Goodheart's Photoguide of Common Skin Disorders: Diagnosis and Management, 2nd ed. Philadelphia: Lippincott Williams & Wilkins, 2003; Ulcer-Hall JC. Sauer's Manual of Skin Diseases, 9th ed. Philadelphia: Lippincott Williams & Wilkins, 2006.)


TABLE 6-5 Vascular and Purpuric Lesions of the Skin

Lesion

CHERRY ANGIOMA



SPIDER ANGIOMA



SPIDER VEIN



PETECHIA AND PURPURA



Features

Appearance: Bright or ruby red, may become brownish with age; 1-3 mm; round, flat, sometimes raised; may be surrounded by a pale halo

Distribution: Found on trunk or extremities

Significance: None; increase in size and number with aging

Appearance: Fiery red; very small to 2 cm; central body, sometimes raised, radiating with ervthema

Distribution: Face, neck, arms, and upper trunk, but almost never below the waist

Significance: Liver disease, pregnancy, vitamin B deficiency; normal in some people

Appearance: Bluish; varies from very small to several inches; may resemble a spider or be linear, irregular, or cascading

Distribution: Most often on the legs, near veins; also on anterior chest

Significance: Often accompanies increased pressure in the superficial veins, as in varicose veins

Appearance: Deep red or reddish purple; fades over time; 1–3 mm or larger; rounded, sometimes irregular, flat

Distribution: Varies

Significance: Blood outside the vessels; may suggest a bleeding disorder or, if petechiae, emboli to skin

(table continues next page)

Vascular and Purpuric Lesions of the Skin (continued)

Lesion

Features

ECCHYMOSIS



Appearance: Purple or purplish blue, fading to green, yellow, and brown over time; larger than petechiae; rounded, oval, or irregular

Distribution: Varies Significance: Blood outside the

vessels; often secondary to bruising or trauma; also seen in bleeding disorders

TABLE 6-6 Skin Tumors



Actinic Keratoses Superficial, flattened papules covered by a dry scale. Often multiple; may be round or irregular; pink, tan, or gravish. They appear on sun-exposed skin of older, fair-skinned persons. Though benign, 1 out of 1,000 per year develop into squamous cell carcinoma (suggested by rapid growth, induration, redness at the base, and ulceration). Typically on face and hands.



Seborrheic Keratoses Common, benign, yellowish to brown, raised lesions that feel slightly greasy and velvety or warty; have a "stuck-on" appearance. Typically multiple and symmetrically distributed on the trunk of older people, but also on the face and elsewhere.

(table continues next page)

TABLE 6-6 Skin Tumors (continued)







In blacks, may appear as small, deeply pigmented papules on cheeks and temples (dermatosis papulosa nigra).

Basal Cell Carcinoma Though malignant, grows slowly and seldom metastasizes. Most common in fair-skinned adults ≥ age 40: usually on the face. Initial translucent nodule spreads, leaving depressed center and firm elevated border. Telangiectatic vessels often visible.

Squamous Cell Carcinoma Usually on sun-exposed skin of fair-skinned adults \geq age 60. May develop in an actinic keratosis. Usually grows more quickly than a basal cell carcinoma, is firmer, and looks redder. The face and the back of the hand are often affected.

Kaposi's Sarcoma in AIDS May appear in many forms: macules, papules, plaques, or nodules almost anywhere on the body. Lesions are often multiple and may involve internal structures.

⁽Sources of photos: Basal Cell Carcinoma: Rapini R. Squamous Cell Carcinoma, Actinic Keratosis, and Seborrheic Keratosis-Hall JC. Sauer's Manual of Skin Diseases, 9th ed. Philadelphia: Lippincott Williams & Wilkins, 2006; Kaposi's Sarcoma in AIDS-DeVita VT Jr, Hellman S, Rosenberg SA [eds]. AIDS: Etiology, Diagnosis, Treatment, and Prevention. Philadelphia: JB Lippincott, 1985.)

Characteristics of Nevi (Moles)

Normal



Diameter smaller than 6 mm Symmetric; regular borders; even in color

Malignant Melanoma: "ABCD"



Asymmetric



Borders irregular



Color varied Diameter more than 6 mm

(Courtesy of American Cancer Society; American Academy of Dermatology)

TABLE 6-8 Hair Loss



Alopecia Areata

Clearly demarcated round or oval patches of hair loss, usually affecting young adults and children. There is no visible scaling or inflammation.



Trichotillomania

Hair loss from pulling, plucking, or twisting hair. Hair shafts are broken and of varying lengths. More common in children, often in settings of family or psychosocial stress.





Hair Loss (continued)



Tinea Capitis ("Ringworm")

Round scaling patches of alopecia. Hairs are broken off close to the surface of the scalp. Usually caused by fungal infection from tinea tonsurans. Mimics seborrheic dermatitis.



(Sources of photos: Alopecia Areata [top], Trichotillomania [top]—Hall JC. Sauer's Manual of Skin Diseases, 9th ed. Philadelphia: Lippincott Williams & Wilkins, 2006; Alopecia Areata [bottom], Tinea Capitis—Goodheart HP. Goodheart's Photoguide of Common Skin Disorders: Diagnosis and Management, 2nd ed. Philadelphia: Lippincott Williams & Wilkins, 2003; Trichotillomania [bottom]—Ostler HB, Mailbach HI, Hoke AW, Schwab IR. Diseases of the Eye and Skin: A Color Atlas. Philadelphia: Lippincott Williams & Wilkins, 2004.)

TABLE (

Fingernails

Clubbing



Dorsal phalanx rounded and bulbous; convexity of nail plate increased. Angle between plate and proximal nail fold increased to 180° or more. Proximal nail folds feel spongy. Many causes, including chronic hypoxia and lung cancer.

Paronychia



Inflammation of proximal and lateral nail folds, acute or chronic. Folds red, swollen, may be tender.

(table continues next page)

TABLE 6-9 Fingernails (continued)

Onycholysis



Painless separation of nail plate from nail bed, starting distally. Many causes.

Terry's Nails



Whitish with a distal band of reddish brown. Seen in aging and some chronic diseases.

Leukonychia



White spots caused by trauma. They grow out with nail(s).

Transverse White Lines



Curved white lines similar to curve of lunula.

They follow an illness and grow out with nails.

The Head and Neck

THE HEALTH HISTORY

Common or Concerning Symptoms

- Headache
- Change in vision
- Double vision, or diplopia
- Hearing loss, earache, tinnitus
- Vertigo
- Nosebleed, or epistaxis
- Sore throat, hoarseness
- Swollen glands
- Goiter

THE HEAD

Headache is a common symptom that always requires careful evaluation because a small fraction of headaches arise from lifethreatening conditions. Elicit a full description of the headache and all seven attributes of the patient's pain (see p. 41).

Is the headache one-sided or bilateral? Steady or throbbing? Continuous or comes and goes? Ask the patient to point to the area of pain or discomfort. Assess chronologic pattern and severity.

See Table 7-1, Primary
Headaches, p. 125, and
Table 7-2, Secondary
Headaches, pp. 126–127.
Tension and migraine
headaches are the most
common recurring headaches.

Tension headaches often arise in the temporal areas; cluster headaches may be retroorbital. Ask about associated symptoms. Inquire specifically about associated nausea and vomiting and neurologic symptoms such as change in vision or motor-sensory deficits.

Ask whether coughing, sneezing, or changing the position of the head has any effect (better, worse, or none) on the headache.

Ask about family history.

THE EYES

Ask "How is your vision?" and "Have you had any trouble with your eyes?" If the patient reports a change in vision, pursue the related details:

- Is the onset sudden or gradual?
- Is the problem worse during close work or at distances?

Changing or progressively severe headaches increase the likelihood of tumor, abscess, or other mass lesion. Extremely severe headaches suggest subarachnoid hemorrhage or meningitis.

Visual aura or scintillating scotomas may accompany migraine. Nausea and vomiting are common with migraine but also occur with brain tumor and subarachnoid hemorrhage.

Such maneuvers may increase pain from brain tumor and acute sinusitis.

Family history is often positive in patients with migraine.

Gradual blurring, often from refractive errors; also in hyperglycemia.

Sudden visual loss suggests retinal detachment, vitreous hemorrhage, or occlusion of the central retinal artery.

Difficulty with close work suggests *hyperopia* (farsightedness) or *presbyopia*

• Is there blurring of the entire field of vision or only parts? Is blurring central, peripheral, or only on one side?

• Has the patient seen lights flashing across the field of vision? Vitreous floaters?

Ask about pain in or around the eyes, redness, and excessive tearing or watering.

Check for *diplopia*, or double vision.

THE EARS

Ask "How is your hearing?"

Does the patient have special difficulty understanding people as they talk? What difference does a noisy environment make?

For complaints of earache, or pain in the ear, ask about associated fever, sore throat, cough, and concurrent upper respiratory infection.

(aging vision); difficulty with distances suggests myopia (near-sightedness).

Slow central loss in *nuclear* cataract and macular degeneration; peripheral loss in advanced open-angle glaucoma; one-sided loss in hemianopsia and quadrantic defects (p. 129).

These symptoms suggest detachment of vitreous from retina. Prompt eye consultation is indicated.

Diplopia in brainstem or cerebellar lesions, also from weakness or paralysis of one or more extraocular muscles.

See Table 7-8, Patterns of Hearing Loss, p. 135.

Sensorineural loss leads to difficulty understanding speech, and complaints that others mumble; noisy environments worsen hearing. In conductive loss, noisy environments may help.

Otitis externa if pain in the external ear; otitis media if pain associated with respiratory infection and in inner ear.

114 The Head and Neck

Tinnitus has no external stimulus—commonly, it manifests as a musical ringing or a rushing or roaring noise.

When associated with hearing loss and vertigo, tinnitus suggests *Ménière's disease*.

Ask about *vertigo*, the perception that the patient or the environment is rotating or spinning.

Vertigo in labrynthitis, CN VII lesions, brainstem lesions

THE NOSE AND SINUSES

Rhinorrhea, or drainage from the nose, frequently is associated with nasal congestion. Ask further about sneezing, watery eyes, and throat discomfort, and also itching in the eyes, nose, and throat.

Causes include viral infections, allergic rhinitis ("hay fever"), and vasomotor rhinitis. Itching favors an allergic cause.

For *epistaxis*, or bleeding from the nose, identify the source carefully—is bleeding from the nose or has the patient coughed up or vomited blood? Assess the site of bleeding, its severity, and associated symptoms.

Local causes of epistaxis include trauma (especially nose picking), inflammation, drying and crusting of the nasal mucosa, tumors, and foreign bodies.

THE MOUTH, THROAT, AND NECK

Sore throat is a frequent complaint. Ask about fever, swollen glands, and any associated cough.

Fever, pharyngeal exudates, and anterior cervical lymphadenopathy, especially without cough, suggest streptococcal pharyngitis, or "strep throat" (p. 139).

Hoarseness may arise from overuse of the voice, allergies, smoking, or inhaled irritants.

Also present in hypothyroidism, in laryngeal disease, or when extrapharyngeal lesions press on the laryngeal nerves

Assess thyroid function. Ask about goiter, temperature intolerance, and sweating.

With goiter, thyroid function may be increased, decreased, or normal. Cold intolerance in hypothyroidism; heat intolerance, palpitations, and involuntary weight loss in hyperthyroidism

$H \in A \cup I \cup H$ PROMOTION AND

Important Topics for Health Promotion and Counseling

- Changes in vision: cataracts, macular degeneration, glaucoma
- Hearing loss
- Oral health

Disorders of vision shift with age. Healthy young adults generally have refractive errors. Up to 25% of adults older than 65 years have refractive errors; cataracts, macular degeneration, and glaucoma also become more prevalent. Glaucoma is the leading cause of blindness in African-Americans and the second leading cause of blindness overall. Glaucoma causes gradual vision loss, with damage to the optic nerve, loss of visual fields, beginning usually at the periphery, and pallor and increasing size of the optic cup (enlarging to more than half the diameter of the optic disc).

More than one third of adults older than 65 years have detectable hearing deficits. Questionnaires and handheld audioscopes work well for periodic screening.

Be sure to promote *oral health*: up to half of all children 5 to 17 years of age have one to eight cavities, and the average U.S. adult has 10 to 17 decayed, missing, or filled teeth. More than half of all adults older than 65 years have no teeth! Inspect the oral cavity for decayed or loose teeth, inflammation of the gingiva, and signs of periodontal disease (bleeding, pus, receding gums, and bad breath). Counsel patients to use fluoride-containing toothpastes, brush, floss, and seek dental care at least annually.

TECHNIQUES OF EXAMINATION

EXAMINATION TECHNIQUES



THE HEAD

Examine the

 Hair, including quantity, distribution, and texture

 Scalp, including lumps or lesions

• Skull, including size and contour

• Face, including symmetry and facial expression

• Skin, including color, texture, hair distribution, and lesions

Coarse and sparse in myxedema, fine in hyperthyroidism

Pilar cysts, psoriasis, pigmented nevi

Hydrocephalus, skull depression from trauma

Facial paralysis; flat affect of depression, moods such as anger, sadness

Pale, fine, hirsute, acne, skin cancer

THE EYES

Test visual acuity in each eye.

Assess visual fields, if indicated.

Diminished acuity

Hemianopsia, quadrantic defects in cerebrovascular accidents (CVAs)

Exophthalmos, strabismus

Inspect the

• Position and alignment of eves

Eyebrows

Eyelids

Seborrheic dermatitis

Sty, chalazion, ectropion, ptosis, xanthelasma

Lacrimal apparatus

Swollen lacrimal sac

EXAMINATION TECHNIQUES

- Conjunctiva and sclera
- Red eye, jaundice
- Cornea, iris, and lens
- Corneal opacity, cataract

Examine pupils for

- Size, shape, and symmetry
- Miosis, mydriasis, anisocoria
- Reactions to light and if these are abnormal
- Absent in paralysis of CN III
- The near reaction

Useful in tonic and Argyll Robertson pupils: slows in tonic pupil; absent in Argyll Robertson pupils of syphilis






THE NEAR REACTION

Assess the extraocular muscles by observing

- The corneal reflections from a midline light
- Imbalance of extraocular muscles
- The six cardinal directions of gaze

Cranial nerve palsy, strabismus, nystagmus, lid lag of hyperthyroidism

Superior rectus (III) Lateral rectus (VI) Inferior rectus (III)



Superior rectus (III) Lateral rectus (VI) Inferior rectus (III)

Convergence

Poor in hyperthyroidism

Inspect the fundi with an ophthalmoscope.

TIPS FOR USING THE OPHTHALMOSCOPE

- Darken the room. Turn the lens disc to the large round beam of white light. Lower the brightness of the light beam to make the examination more comfortable for the patient.
- Turn the lens disc to the 0 diopter (a diopter measures the power of a lens to converge or diverge light).
- Hold the ophthalmoscope in your right hand to examine the patient's right eye; hold it in your left hand to examine the patient's left eye to avoid bumping the patient's nose.
- Brace the ophthalmoscope firmly against the medial aspect of your bony orbit, with the handle tilted laterally at about 20°. Instruct the patient to look slightly up and over your shoulder at a point directly ahead on the wall.
- Place yourself about 15 inches away from the patient and at an angle 15° lateral to the patient's line of vision. Look for the orange glow in the pupil—the red reflex. Note any opacities interrupting the red reflex. No red reflex suggests an opacity of the lens (cataract) or possibly the vitreous.
- Place the thumb of your other hand across the patient's eyebrow. Keeping the light beam focused on the red reflex, move in at a 15° angle toward the pupil until you almost touch the patient's eyelashes. Adjust the position of your ophthalmoscope and angle of vision as a unit until you see the fundus.

Inspect the fundi for the following:


Red reflex

Cataracts, artificial eye

Optic disc

Papilledema, glaucomatous cupping, optic atrophy. See Table 7-5, Abnormalities of the Optic Disc, p. 132.

POSSIBLE FINDINGS



- Arteries, veins, and A–V crossings
- Adjacent retina (note any lesions)
- Macular area
- Anterior structures

A-V nicking, copper wiring in hypertensive changes

Hemorrhages, exudates, cotton-wool patches, microaneurysms, pigmentation

Macular degeneration

Vitreous floaters, cataracts

TIPS FOR EXAMINING THE OPTIC DISC AND THE RETINA

- Locate the optic disc. Look for the round yellowish-orange structure.
- Now, bring the optic disc into sharp focus by adjusting the lens of your ophthalmoscope.
- Inspect the optic disc. Note the following features:
 - The sharpness or clarity of the disc outline
 - The color of the disc
 - The size of the physiologic cup (an enlarged cup suggests chronic open-angle glaucoma)

(continued)

TIPS FOR EXAMINING THE OPTIC DISC AND THE RETINA (CONTINUED)

- Venous pulsations in the retinal veins as they emerge from the central portion of the disc (loss of venous pulsations in pathologic conditions such as head trauma, meningitis, or mass lesions may be an early sign of elevated intracranial pressure)
- Inspect the retina. Distinguish arteries from veins based on the features listed below.

	Arteries	Veins
Color	Light red	Dark red
Size	Smaller (3/3 to 4/5 the diameter of veins)	Larger
Light Reflex (reflection)	Bright	Inconspicuous or absent

- Follow the vessels peripherally in each of four directions.
- Inspect the fovea and surrounding macula. Macular degeneration types include dry atrophic (more common but less severe) and wet exudative (neovascular). Undigested cellular debris, called drusen, may be hard or soft.
- Assess for any papilledema.

THE EARS

Examine on each side:

THE AURICLE

Inspect the auricle. Keloid, epidermoid cyst

If you suspect otitis:

 Move the auricle up and down, and press on the tragus. Pain in otitis externa ("the tug test")

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

• Press firmly behind the ear.

Possible tenderness in otitis media and mastoiditis

THE EAR CANAL AND EARDRUM


Pull the auricle up, back, and slightly out.

Inspect, through an otoscope speculum:

- The canal
- The eardrum

Cerumen; swelling and erythema in otitis externa

Red bulging drum in acute otitis media; serous otitis media, tympanosclerosis, perforations. See Table 7-7, Abnormal Eardrum, p. 134.



HEARING

Assess auditory acuity to whispered or spoken voice.

If hearing is diminished, use a 512-Hz tuning fork to:

These tests help distinguish between sensorineural and conduction hearing loss.

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

- Test lateralization (Weber test). Place vibrating and tuning fork on vertex of skull and check hearing.
- See Table 7-8, Patterns of Hearing Loss, p. 135.
- Compare air and bone conduction (Rinne test).
 Place vibrating and tuning fork on mastoid bone, then remove and check hearing.

THE NOSE AND SINUSES

Inspect the external nose.

Inspect, through a speculum, the

- Nasal mucosa that covers the septum and turbinates, noting its color and any swelling
- rhinitis, swollen and pale in allergic rhinitis; polyps; ulcer from cocaine use

Swollen and red in viral

 Nasal septum for position and integrity Deviation, perforation

Palpate the frontal and maxillary sinuses.

Tender in acute sinusitis

THE MOUTH AND PHARYNX

Inspect the

Lips
 Cyanosis, pallor, cheilosis

Oral mucosa
 Canker sores

Gums
 Gingivitis, periodontal disease

Teeth Dental caries, tooth loss

examination techniques	POSSIBLE FINDINGS
• Roof of the mouth	Torus palatinus
• Tongue, including	See Table 7-10, Abnormalities of the Tongue, pp. 137–138.
Papillae	Glossitis
Symmetry	Deviation to one side from paralysis of CN XII from CVA
Any lesions	Cancer
• Floor of the mouth	Cancer
• Pharynx, including	See Table 7-11, Abnormalities of the Pharynx, p. 139.
Color or any exudate	Pharyngitis
Presence and size of tonsils	Tonsillitis, peritonsillar abscess
Symmetry of the soft palate as patient says "ah"	Soft palate fails to rise in paralysis of CN X from CVA
THE NECK	
Inspect the neck.	Scars, masses, torticollis
Dalmaka kha kasanka na daa	Consider lymphadenenathy

Palpate the lymph nodes. Cervical lymphadenopathy from inflammation, malignancy, HIV

Deviated trachea from neck Inspect and palpate the position of the trachea. mass or pneumothorax

Inspect the thyroid gland:

Goiter, nodules. See At rest Table 7-12, Abnormalities of the Thyroid Gland, p. 140. • As patient swallows water

EXAMINATION TECHNIQUES

From behind patient, palpate the thyroid gland, including the isthmus and the lateral lobes:

- At rest
- As patient swallows water

POSSIBLE FINDINGS

Goiter, nodules, tenderness of thyroiditis



After examining the thyroid gland from behind the patient, you may proceed to musculoskeletal examination of the neck and upper back and check for costovertebral angle tenderness.

RECORDING YOUR FINDINGS

Recording the Physical Examination—The Head, Eyes, Ears, Nose, and Throat (HEENT)

HEENT: Head—Skull is normocephalic/atraumatic (NC/AT). Hair with average texture. Eyes—Visual acuity 20/20 bilaterally. Sclera white; conjunctiva pink. Pupils constrict 4 mm to 2 mm, equally round and reactive to light and accommodations. Disc margins sharp; no hemorrhages or exudates; no arteriolar narrowing. Ears—Acuity good to whispered voice. Tympanic membranes (TMs) with good cone of light. Weber midline. AC > BC. Nose—Nasal mucosa pink, septum midline; no sinus tenderness. Throat (or Mouth)—Oral mucosa pink; dentition good; pharynx without exudates. Neck—Trachea midline. Neck supple; thyroid isthmus palpable, lobes not felt. Lymph Nodes—No cervical, axillary, epitrochlear, inguinal adenopathy.

AIDS TO INTERPRETATION

TABLE	7-1	Prima

ıry Headaches

Problem	Common Characteristics	Associated Symptoms, With Provoking and Relieving Factors
Tension	Location: Variable Quality: Pressing or tightening pain; mild to moderate intensity Onset: Gradual Duration: Minutes to days	Sometimes photophobia, phonophobia; nausea absent ↑ by sustained muscle tension, as in driving or typing ↓ possibly by massage, relaxation
Migraine • With aura • Without aura • Variants	Location: Unilateral in ~70%; bifrontal or global in ~30% Quality: Throbbing or aching, variable in severity Onset: Fairly rapid, reaching a peak in 1–2 hrs Duration: 4–72 hrs	Nausea, vomiting, photophobia, visual auras (flickering zigzagging lines), motor auras affecting hand or arm, sensory auras (numbness, tingling usually precede headache) ↑ by alcohol, certain foods, or tension, noise, bright light. More common premenstrually. ↓ by quiet dark room, sleep
Cluster	Location: Unilateral, usually behind or around the eye Quality: Deep, continuous, severe Onset: Abrupt, peaks within minutes Duration: Up to 3 hrs	Lacrimation, rhinorrhea, miosis, ptosis, eyelid edema, conjunctival infection † sensitivity to alcohol during some episodes

TABLE 7-2 Secondary Headaches			
Problem	Common Characteristics	Associated Symptoms, With Provoking and Relieving Factors	
Analgesic Rebound	Location: Previous headache pattern Quality: Variable Onset: Variable Duration: Depends on prior headache pattern	Depends on prior headache pattern by fever, carbon monoxide, hypoxia, withdrawal of caffeine, other headache triggers —depends on cause	
Headaches From Eye Disorders Errors of Refraction (farsighted- ness and astigmatism, but not near- sightedness)	Location: Around and over the eyes; may radiate to the occipital area Quality: Steady, aching, dull Onset: Gradual Duration: Variable	Eye fatigue, "sandy" sensation in eyes, redness of the conjunctiva ↑ by prolonged use of the eyes, particularly for close work ↓ by rest of the eyes	
Acute Glaucoma	Location: In and around one eye Quality: Steady, aching, often severe Onset: Often rapid Duration: Variable, may depend on treatment	Diminished vision, sometimes nausea and vomiting —sometimes by drops that dilate the pupils	
Headache from Sinusitis	Location: Usually above eye (frontal sinus) or over maxillary sinus	Local tenderness, nasal congestion, discharge, and fever † by coughing, sneezing, or jarring the head	

TABLE 7-2 Secondary Headaches (continued)		
Problem	Common Characteristics	Associated Symptoms, With Provoking and Relieving Factors
	Quality: Aching or throbbing, vari- able in severity; consider possible migraine Onset: Variable Duration: Often several hours at a time, recurring over days or longer	↓ by nasal decongestants, antibiotics
Meningitis	Location: Generalized Quality: Steady or throbbing, very severe Onset: Fairly rapid Duration: Variable, usually days	Fever, stiff neck
Subarachnoid Hemorrhage	Location: Generalized Quality: Severe, "the worst of my life" Onset: Usually abrupt; prodromal symptoms may occur Duration: Variable, usually days	Nausea, vomiting, possibly loss of consciousness, neck pain
Brain Tumor	Location: Varies with the location of the tumor Quality: Aching, steady, variable in intensity	↑ by coughing, sneezing, or sudden movements of the head
		(table continues next pa

TABLE 7-2	Secondary Headaches (continued)		
Problem	Common Characteristics	Associated Symptoms, With Provoking and Relieving Factors	
	Onset: Variable Duration: Often brief		
Cranial Neuralgias: Trigeminal Neuralgia (CN V)	Location: Cheek, jaws, lips, or gums; trigeminal nerve divisions 2 and 3 >1 Quality: Shocklike, stabbing, burning, severe Onset: Abrupt, paroxysmal Duration: Each jab lasts seconds but recurs at intervals of seconds or minutes	Exhaustion from recurrent pain by touching certain areas of the lower face or mouth; chewing, talking, brushing teeth	
Giant Cell (Temporal) Arteritis	Location: Near the involved artery, often the temporal, also the occipital; age-related Quality: Throbbing, generalized, persistent, often severe Onset: Gradual or rapid Duration: Variable	Tenderness of the adjacent scalp; fever (in ~50%), fatigue, weight loss; new headache (~60%), jaw claudication (~50%), visual loss or blindness (~15%—20%), polymyalgia rheumatica (~50%) ↑ by movement of neck and shoulders	

TABLE /-Z Secondary Heddacnes (continued)		
Problem	Common Characteristics	Associated Symptoms, With Provoking and Relieving Factors
Post-traumatic Headache	Location: Injured area, but not necessarily Quality: Generalized, dull, aching, constant Onset: Within hours to 1–2 days of the injury Duration: Weeks, months, or even years	Poor concentration, problems with memory, vertigo, irritability, restlessness, fatigue by mental and physical exertion, straining, stooping, emotional excitement, alcohol by rest

Altitudinal (horizontal) defect, usually resulting from a vascular lesion of the retina Unilateral blindness, from a lesion of the retina or optic nerve Bitemporal hemianopsia, from a lesion of the optic chiasm Homonymous hemianopsia, from a lesion of the optic tract or optic radiation on the side opposite the blind area Homonymous quadrantic defect, from a partial lesion of the optic radiation on the side opposite the blind area Fight Figh

TABLE 7-4 Physical Findings in and Around the Eye

Eyelids



Ptosis. A drooping upper eyelid that narrows the palpebral fissure from a muscle or nerve disorder



Ectropion. Outward turning of the margin of the lower lid, exposing the palpebral conjunctiva




Entropion. Inward turning of the lid margin, causing irritation of the cornea or conjunctiva

In and Around the Eye



Pinguecula. Harmless yellowish nodule in the bulbar conjunctiva on either side of the iris; associated with aging



Episcleritis. A localized ocular redness from inflammation of the episcleral vessels



Physical Findings in and Around the Eye (continued)



Sty. A pimple-like infection around a hair follicle near the lid margin



Chalazion. A beady nodule in either eyelid caused by a chronically inflamed meibomian gland

TABLE 7-5 Abnormalities of the Optic Disc

Process

Normal



Tiny disc vessels give normal color to the disc.

Appearance

Disc is yellowish orange to creamy pink.
Disc vessels are tiny.
Disc margins are sharp (except perhaps nasally).

Papilledema



Venous stasis leads to engorgement and swelling. Disc is pink,
hyperemic.
Disc vessels are
more visible,
more numerous,
and curve over
the borders of
the disc.
Disc is swollen,
with margins
blurred.

Glaucomatous Cupping



Increased pressure within the eye leads to increased cupping (backward depression of the disc) and atrophy.

The base of the enlarged cup is pale.

Optic Atrophy



Death of optic nerve fibers leads to loss of the tiny disc vessels.

Disc is white. Disc vessels are absent.

Diabetic Retinopathy

Nonproliferative Retinopathy, Moderately Severe



Nonproliferative Retinopathy, Severe



Proliferative Retinopathy, With Neovascularization



Proliferative Retinopathy, Advanced



Note tiny red dots or microaneurysms, also the ring of hard exudates (white spots) located superotemporally. Retinal thickening or edema in the area of hard exudates can impair visual acuity if it extends to center of macula (detection requires specialized stereoscopic examination).

In superior temporal quadrant, note large retinal hemorrhage between two cotton-wool patches, beading of the retinal vein just above, and tiny tortuous retinal vessels above superior temporal artery, termed intraretinal microvascular abnormalities

Note new preretinal vessels arising on disc and extending across disc margins. Visual acuity is currently normal, but risk of severe visual loss is high. (Photocoagulation can reduce this risk ≥50%.)

Same eye as above, but 2 yrs later and without treatment. Neovascularization has increased, now with fibrous proliferations, distortion of macula, and reduced visual acuity.

(Source: Early Treatment Diabetic Retinopathy Study Research Group. Courtesy of M. F. Davis, MD, University of Wisconsin, Madison.)

TABLE 7-7

Abnormal Eardrum

Perforation



Hole in the eardrum that may be central or marginal Usually from *otitis media* or trauma

Tympanosclerosis



A chalky white patch Scar of an old otitis media; of little or no clinical consequence

Serous Effusion



Amber fluid behind the eardrum, with or without air bubbles Associated with viral upper respiratory infections or sudden changes in atmospheric pressure (diving, flying)

Acute Otitis Media



Red, bulging drum, loss of landmarks Associated with bacterial infection

TABLE 7-8 P	atterns of Hearing L	oss
	Conductive Loss	Sensorineural Loss
Impaired understanding of words	Minor	Often troublesome
Effect of noisy environment	May help	Increases the hearing difficulty
Usual age of onset	Childhood, young adulthood	Middle and old age
Ear canal and drum	Often a visible abnormality	Problem not visible
Weber test (in unilateral hearing loss)	Lateralizes to the impaired ear	Lateralizes to the good ear
Rinne test	BC > AC or $BC = AC$	AC > BC
Causes include	Plugged ear canal, otitis media, immobile or perforated drum, otosclerosis, foreign body	Sustained loud noise, drugs, inner ear infections, trauma, hereditary disorder, aging

TABLE 7-9 Abnormalities of the Lips



Angular cheilitis. Softening and cracking of the angles of the mouth



Herpes simplex. Painful vesicles, followed by crusting; also called cold sore or fever blister



Angioedema. Diffuse, tense, subcutaneous swelling, usually allergic in cause



Hereditary hemorrhagic telangiectasia. Red spots, significant because of associated bleeding from nose and GI tract



Peutz-Jeghers syndrome. Brown spots of the lips and buccal mucosa, significant because of their association with intestinal polyposis



Syphilitic chancre. A firm lesion that ulcerates and may crust



Carcinoma of the lip. A thickened plaque or irregular nodule that may ulcerate or crust; malignant

Abnormalities of the Tongue



Geographic tongue. Scattered areas in which the papillae are lost, giving a maplike appearance; harmless



Hairy tongue. Results from elongated papillae that may look yellowish, brown, or black; harmless



Fissured tongue. May appear with aging; harmless



Smooth tongue. Results from loss of papillae, caused by vitamin B or iron deficiency or possibly anticancer drugs



Candida infection. May show a thick, white coat, which, when scraped off, leaves a raw red surface; tongue may also be red; antibiotics, corticosteroids, AIDS may predispose



Hairy leukoplakia. White raised, feathery areas, usually on sides of tongue. Seen in HIV/AIDS

Abnormalities of the Tongue (continued)



Varicose veins. Dark round spots in the undersurface of the tongue, associated with aging; also called caviar lesions



Aphthous ulcer (canker sore). Painful, small, whitish ulcer with a red halo; heals in 7-10 days



Mucous patch of syphilis. Slightly raised, oval lesion, covered by a gravish membrane



Carcinoma of the tongue or floor of the mouth. A malignancy that should be considered in any nodule or nonhealing ulcer at the base or edges of the mouth

Abnormalities of the Pharynx



Pharyngitis, mild to moderate. Note redness and vascularity of the pillars and uvula.



Pharyngitis, diffuse. Note redness is diffuse and intense. Cause may be viral or, if patient has fever, bacterial. If patient has no fever, exudate, or cervical lymphadenopathy, viral infection is more likely.



Exudative pharyngitis. A sore red throat with patches of white exudate on the tonsils is associated with streptococcal pharyngitis and some viral illnesses.



Diphtheria. An acute infection caused by Corynebacterium diphtheriae. The throat is dull red, and a gray exudate appears on the uvula, pharynx, and tongue.



Koplik's spots. These small white specks that resemble grains of salt on a red background are an early sign of measles.

Abnormalities of the Thyroid Gland



Diffuse enlargement. May result from Graves' disease, Hashimoto's thyroiditis, endemic goiter (iodine deficiency), or sporadic goiter



Multinodular goiter. An enlargement with two or more identifiable nodules, usually metabolic in cause



Single nodule. May result from a cyst, a benign tumor, or cancer of the thyroid, or may be one palpable nodule in a clinically unrecognized multinodular goiter

The Thorax and Lungs

THE HEALTH HISTORY

Common or Concerning Symptoms

- Chest pain
- Shortness of breath (dyspnea)
- Wheezing
- Cough
- Blood-streaked sputum (hemoptysis)

Complaints of *chest pain* or *chest discomfort* raise the specter of heart disease but often arise from conditions in the thorax and lungs. For this important symptom, keep the possible causes below in mind. Also see Table 8-1, Chest Pain, pp. 152–153.

 The myocardium 	Angina pectoris, myocardial
•	infarction

- The pericardium Pericarditis
- The aorta Dissecting aortic aneurysm
- The trachea and large Bronchitis bronchi
- The parietal pleura
 Pericarditis, pneumonia
- The chest wall, including the musculoskeletal system and skin
 Costochondritis, herpes zoster

142 The Thorax and Lungs

The esophagus
 Reflux esophagitis, esophageal spasm

 Extrathoracic structures such as the neck, gallbladder, stomach
 Cervical arthritis, biliary colic, gastritis

For patients who are *short of breath*, focus on such *pulmonary complaints* as

• dyspnea and wheezing See Table 8-2, Dyspnea, pp. 154–155.

 cough and hemoptysis
 See Table 8-3, Cough and Hemoptysis, pp. 156–159.

HEALTH PROMOTION AND COUNSELING

Despite declines in smoking over the past several decades, ~23% of Americans still smoke.* Regularly counsel all adults, pregnant women, parents, and adolescents who smoke to stop. Include "the five A's":

- Ask about smoking at each visit,
- Advise patients to quit in regular, clear, personalized messages.
- Assess patient readiness to quit.
- Assist patients to set stop dates and provide educational materials for self-help.
- Arrange for follow-up visits to monitor and support progress.

Provide flu shots and pneumonia vaccine in target groups.


^{*}Centers for Disease Control (CDC). At A Glance. Targeting tobacco use: The nation's leading cause of death, 2004. Available at: www.cdc.gov/nccdphp/aag/pdf/aag_osh2004.pdf. Accessed September 11, 2007.

TECHNIQUES OF EXAMINATION

EXAMINATION TECHNIQUES



SURVEY OF THORAX



Inspect the thorax and its respiratory movements. Note:

- Rate, rhythm, depth, and effort of breathing
- Inspiratory retraction of the supraclavicular areas
- Inspiratory contraction of the sternomastoids

Observe shape of patient's chest.

Listen to patient's breathing for

 Rate and rhythm of breathing

Tachypnea, hyperpnea, Cheyne-Stokes breathing

Occurs in chronic obstructive pulmonary disease (COPD), asthma, upper airway obstruction

Indicates severe breathing difficulty

Normal or barrel chest (see Table 8-4, Deformities of the Thorax, pp. 159-160)

14-16 breaths/minute in adults (see Chapter 4, pp. 64–65)

POSSIBLE FINDINGS

Stridor

Stridor in upper airway obstruction from foreign body or epiglottitis

Wheezes

Expiratory wheezing in asthma and COPD

THE POSTERIOR CHEST

Inspect the chest for

- Deformities or asymmetry
- Abnormal inspiratory retraction of the interspaces
- Impairment or unilateral lag in respiratory movement

Palpate the chest for

- Tender areas
- Assessment of visible abnormalities
- Chest expansion



Kyphoscoliosis

Retraction in airway obstruction

Disease of the underlying lung or pleura, phrenic nerve palsy

Fractured ribs

Masses, sinus tracts

Impairment, both sides in COPD and restrictive lung disease


POSSIBLE FINDINGS

• Tactile fremitus as the patient says "aa" or "blue moon"

Local or generalized decrease or increase

Percuss the chest in the areas illustrated, comparing one side with the other at each level, using the side-toside "ladder pattern."

Dullness when fluid or solid tissue replaces normally airfilled lung; hyperresonance in emphysema or pneumothorax






Percussion Notes	;	
	Relative Intensity, Pitch, and Duration	Examples
Flatness	Soft/high/short	Large pleural effusion
Dullness	Medium/medium/ medium	Lobar pneumonia
Resonance	Loud/low/long	Normal lung, simple chronic bronchitis
Hyperresonance	Louder/lower/ longer	Emphysema, pneumothorax
Tympany	Loud/high (timbre is musical)	Large pneumothorax

POSSIBLE FINDINGS

Percuss level of diaphragmatic dullness on each side and **estimate** diaphragmatic descent after patient takes full inspiration.

Pleural effusion or a paralyzed diaphragm raises level of dullness.



Listen to chest with stethoscope in the "ladder" pattern, again comparing sides.

- Evaluate the breath sounds.
- Note any adventitious (added) sounds.

Observe their qualities, place in the respiratory cycle, and location on the chest wall. Do they clear with deep breathing or coughing?

See Table 8-5, Physical Findings in Selected Chest Disorders, p. 161.

Vesicular, bronchovesicular, or bronchial breath sounds; decreased breath sounds from decreased airflow

Crackles (fine and coarse) and continuous sounds (wheezes and rhonchi)

Breath Sounds			
	Duration	Intensity and Pitch of Expiratory Sound	Example Locations
Vesicular	Insp > exp	Soft/low	Most of the lungs
Bronchovesicular	Insp = exp	Medium/ medium	1st and 2nd interspaces, interscapular area
Bronchial	Exp > insp	Loud/high	Over the manubrium; lobar pneumonia
Tracheal	Insp = exp	Very loud/ high	Over the trachea

Duration is indicated by the length of the line, intensity by the width of the line, and pitch by the slope of the line.

Adventitious or Added Breath Sounds

Crackles (or Rales)


Wheezes and Rhonchi

- Discontinuous
- Intermittent, nonmusical,

 ≥250 msec, musical, prolonged and brief
- Like dots in time
- Fine crackles: Soft, highpitched, very brief (5-10 msec)

- Continuous
- (but not necessarily persisting throughout the respiratory cycle)
- Like dashes in time
- Wheezes: Relatively high-pitched (≥400 Hz) with hissing or shrill quality

- louder, lower in pitch, brief (20–30 msec)
- Coarse crackles: Somewhat Rhonchi: Relatively low-pitched (≤200 Hz) with snoring quality



POSSIBLE FINDINGS

Assess transmitted voice sounds if you have heard bronchial breath sounds in abnormal places. Ask patient to

• Say "99" and "ee."

Bronchophony and egophony

• Whisper "99" or "1, 2, 3."

Whispered pectoriloquy


 Transmitted Voice Sounds 	
Through Normally Air-Filled Lung	Through Airless Lung*
Spoken words muffled and indistinct	Spoken words louder, clearer (bronchophony)
Spoken "ee" heard as "ee"	Spoken "ee" heard as "ay" (egophony)
Whispered words faint and indistinct, if heard at all	Whispered words louder, clearer (whispered pectoriloquy)
Usually accompanied by vesicular breath sounds and normal tactile fremitus	Usually accompanied by bronchial or bronchovesicular breath sounds and increased tactile fremitus

^{*}As in lobar pneumonia and toward the top of a large pleural effusion


While the patient is still sitting, you may inspect the breasts and examine the axillary and epitrochlear lymph nodes, and examine the temporomandibular joint and the musculoskeletal system of the upper extremities.

POSSIBLE FINDINGS

THE ANTERIOR CHEST



ANTERIOR VIEW



RIGHT ANTERIOR **OBLIQUE VIEW**

Inspect the chest for

• Deformities or asymmetry

Intercostal retraction

 Impaired or lagging respiratory movement

Palpate the chest for

Tender areas

 Assessment of visible abnormalities

• Respiratory expansion

Tactile fremitus

Percuss the chest in the areas illustrated.

Pectus excavatum

From obstructed airways

Disease of the underlying lung or pleura, phrenic nerve palsy

Tender pectoral muscles, costochondritis

Flail chest

Normal cardiac dullness may disappear in emphysema.



Listen to the chest with stethoscope. Note:

- Breath sounds
- Adventitious sounds
- If indicated, transmitted voice sounds

SPECIAL TECHNIQUES

ASSESSMENT OF PULMONARY FUNCTION

Walk with patient down the hall or up a flight of stairs. Observe rate, effort, and sound of breathing, and inquire about symptoms.

Older adults walking 8 feet in <3 seconds are less likely to be disabled than those taking >5-6 seconds.

FORCED EXPIRATORY TIME

Ask patient to take a deep breath in and then breathe out as quickly and completely as possible, with mouth open. Listen over trachea with diaphragm of stethoscope, and time audible expiration. Try to get three consistent readings, allowing rests as needed.

If the patient understands and cooperates well, a forced expiratory time of 6 to 8 strongly suggests COPD.

RECORDING YOUR FINDINGS

Recording the Physical Examination— The Thorax and Lungs

"Thorax is symmetric with good expansion. Lungs resonant. Breath sounds vesicular; no rales, wheezes, or rhonchi. Diaphragms descend 4 cm bilaterally."

OR

"Thorax symmetric with moderate kyphosis and increased anteroposterior (AP) diameter, decreased expansion. Lungs are hyperresonant. Breath sounds distant with delayed expiratory phase and scattered expiratory wheezes. Fremitus decreased; no bronchophony, egophony, or whispered pectoriloguy. Diaphragms descend 2 cm bilaterally."

(Suggests COPD)

AIDS TO INTERPRETATION

TABLE 8-1

Chest Pain

Problem and Location

Quality, Severity, Timing, and Associated Symptoms

Cardiovascular

Angina Pectoris

Retrosternal or across the anterior chest, sometimes radiating to the shoulders, arms, neck, lower jaw, or upper abdomen

Myocardial Infarction
Same as in angina

Pericarditis

Precordial: May radiate to the tip of the shoulder and to the neck

Retrosternal

Dissecting Aortic Aneurysm Anterior chest, r

Anterior chest, radiating to the neck, back, or abdomen Quality: Pressing, squeezing, tight, heavy, occasionally burning Severity: Mild to moderate, sometimes perceived as discomfort rather than pain

Timing: Usually 1–3 min but up to 10 min; prolonged episodes up to 20 min

Associated Symptoms: Sometimes dyspnea, nausea, swelling

Quality: Same as in angina Severity: Often but not always a severe pain

Timing: 20 min to several hours **Associated Symptoms:** Nausea, vomiting, sweating, weakness

Quality: Sharp, knifelike Severity: Often severe Timing: Persistent

Associated Symptoms: Of the underlying illness; relieved by leaning forward

Quality: Crushing Severity: Severe Timing: Persistent

Associated Symptoms: Of the

underlying illness

Quality: Ripping, tearing **Severity:** Very severe

Timing: Abrupt onset, early peak, persistent for hours or more **Associated Symptoms:** Syncope,

hemiplegia, paraplegia

(table continues next page)

TABLE 8-

Chest Pain (continued)

Problem and Location

Quality, Severity, Timing, and Associated Symptoms

Pulmonary

Tracheobronchitis

Upper sternal or on either side of the sternum

Pleural Pain

Chest wall overlying the process

Gastrointestinal and Other

Reflex Esophagitis Retrosternal, may radiate to the back

Diffuse Esophageal Spasm Retrosternal, may radiate to the back, arms, and jaw

Chest Wall Pain Often below the left breast or along the costal cartilages;

Anxiety/Panic Attack

also elsewhere

Quality: Burning

Severity: Mild to moderate

Timing: Variable

Associated Symptoms: Cough

Quality: Sharp, knifelike Severity: Often severe Timing: Persistent

Associated Symptoms: Of the

underlying illness

Quality: Burning, may be squeezing

Severity: Mild to severe

Timing: Variable

Associated Symptoms: Sometimes regurgitation, dysphagia

Quality: Usually squeezing Severity: Mild to severe

Timing: Variable

Associated Symptoms: Dysphagia Quality: Stabbing, sticking, or dull

aching

Severity: Variable

Timing: Fleeting to hours or days Associated Symptoms: Often local

tenderness

TABLE 8-2 Dyspnea

Problem	Timing	Provoking and Relieving Factors
Left-sided Heart Failure (left ventricular failure or mitral stenosis)	Dyspnea may progress slowly or suddenly, as in acute pulmonary edema	↑ by exertion, lying down ↓ by rest, sitting up, though dyspnea may become persistent Associated Symptoms: Often cough, orthopnea, paroxys- mal nocturnal dyspnea, sometimes wheezing
Chronic Bronchitis (may be seen with COPD)	Chronic productive cough followed by slowly progressive dyspnea	↑ by exertion, inhaled irritants, respiratory infections ↓ by expectoration, rest though dyspnea may become persistent Associated Symptoms: Chronic productive cough, recurrent respiratory infections; wheezing possible
Chronic Obstructive Pulmonary Disease (COPD)	Slowly progressive; relatively mild cough later Acute episodes, then symptom- free periods; nocturnal episodes common	↑ by exertion ↓ by rest, though dyspnea may become persistent Associated Symptoms: Cough with scant mucoid sputum ↑ by allergens, irritants, respiratory infections, exercise, emotion ↓ by separation from aggra- vating factors Associated Symptoms: Wheezing, cough, tightness in chest

TABLE 8-2 Dyspnea (continued)

Problem	Timing	Provoking and Relieving Factors
Acute Pulmonary Embolism	Sudden onset of dyspnea	Associated Symptoms: Often none; retrosternal oppressive pain if occlusion is massive; pleuritic pain, cough, and hemoptysis may follow an embolism if pulmonary infarction ensues; symptoms of anxiety
Pneumonia	Acute illness; timing varies with causative agent	Associated Symptoms: Pleuritic pain, cough, sputum, fever, though not necessarily present
Diffuse Interstitial Lung Diseases (sarcoidosis, neoplasms, asbestosis, idiopathic pulmonary fibrosis)	Progressive; varies in rate of develop- ment depending on cause	↑ by exertion ↓ by rest, though dyspnea may become persistent Associated Symptoms: Often weakness, fatigue; cough less common than in other lung diseases
Spontaneous Pneumothorax	Sudden onset of dyspnea	Associated Symptoms: Pleuritic pain, cough

TABLE 8-3 Cough	and Hemoptysis
Problem	Cough, Sputum, Associated Symptoms, and Setting
Acute Inflammation	
Laryngitis	Cough and Sputum: Dry, or with variable amounts of sputum Associated Symptoms and Setting: Acute, fairly minor illness with hoarseness. May be associated with viral nasopharyngitis
Tracheobronchitis	Cough and Sputum: Dry or productive of sputum Associated Symptoms and Setting: An acute, often viral illness, with burning retrosternal discomfort
Mycoplasma and Viral Pneumonias	Cough: Dry and hacking Sputum: Often mucoid Associated Symptoms and Setting: An acute febrile illness, often with malaise, headache, and possibly dyspnea
Bacterial Pneumonias	Cough and Sputum: With pneumococcal infection, mucoid or purulent; may be blood streaked, diffusely pinkish, or rusty. With Klebsiella, similar to pneumococcal, or sticky red and jelly-like. Associated Symptoms and Setting: An acute illness with chills, high fever, dyspnea, and chest pain; often preceded by acute upper respiratory infection. Klebsiella often in older alcoholic men.

(table continues next page)

TABLE 8-3 Cough	and Hemoptysis (continued)
Problem	Cough, Sputum, Associated Symptoms, and Setting
Chronic Inflammation	
Postnasal Drip	Cough: Chronic Sputum: Mucoid or mucopurulent Associated Symptoms and Setting: Repeated attempts to clear the throat. Postnasal drip, discharge in posterior pharynx. Associated with chronic rhinitis, with or without sinusitis
Chronic Bronchitis	Cough: Chronic Sputum: Mucoid to purulent; may be blood streaked or even bloody Associated Symptoms and Setting: Often long-standing cigarette smoking. Recurrent superimposed infections; often wheezing and dyspnea.
Bronchiectasis	Cough: Chronic Sputum: Purulent, often copious and foul smelling; may be blood streaked or bloody Associated Symptoms and Setting: Recurrent bronchopulmonary infections common; sinusitis may coexist
Pulmonary Tuberculosis	Cough and Sputum: Dry, mucoid or purulent; may be blood streaked or bloody Associated Symptoms and Setting: Early, no symptoms. Later, anorexia, weight loss, fatigue, fever, and night sweats.
Lung Abscess	Cough and Sputum: Purulent and foul smelling; may be bloody Associated Symptoms and Setting: A febrile illness. Often poor dental hygiene and a prior episode of impaired consciousness

TABLE 8-3 Cough	and Hemoptysis (continued)
Problem	Cough, Sputum, Associated Symptoms, and Setting
Asthma	Cough and Sputum: Thick and mucoid, especially near end of an attack Associated Symptoms and Setting: Episodic wheezing and dyspnea, but cough may occur alone. Often a history of allergy
Gastroesophageal Reflux	Cough and Sputum: Chronic, especially at night or early morning Associated Symptoms and Setting: Wheezing, especially at night (often mistaken for asthma), early morning hoarseness, repeated attempts to clear throat. Often with history of heartburn and regurgitation
Neoplasm	Cough: Dry to productive
Lung Cancer	Sputum: May be blood streaked or bloody Associated Symptoms and Setting: Usually a long history of cigarette smoking
Cardiovascular	Cough: Often dry, especially on
Disorders Left Ventricular Failure or Mitral Stenosis	exertion or at night Sputum: May progress to pink and frothy, as in pulmonary edema, or to frank hemoptysis Associated Symptoms and Setting: Dyspnea, orthopnea, paroxysmal nocturnal dyspnea
Pulmonary Embolus	Cough: Dry to productive Sputum: May be dark, bright red, or mixed with blood


TABLE 8-3 Cough	and Hemoptysis (continued)
Problem	Cough, Sputum, Associated Symptoms, and Setting
	Associated Symptoms and Setting: Dyspnea, anxiety, chest pain, fever; factors that predispose to deep venous thrombosis
Irritating Particles, Chemicals, or Gases	Cough and Sputum: Variable. There may be a latent period between exposure and symptoms. Associated Symptoms and Setting: Exposure to irritants; eye, nose, and throat symptoms

TABLE 8-4 Deformities of the Thorax

Cross-Section of Thorax Normal Adult

The thorax is wider than it is deep; lateral diameter is greater than anteroposterior (A-P) diameter.

Barrel Chest Has increased A-P diameter, seen in normal infants and normal aging; also in COPD.




(table continues next page)

TABLE 8-4 Deformities of the Thorax (continued)

Traumatic Flail Chest

If multiple ribs are fractured, can see paradoxical movements of the thorax. Descent of the diaphragm decreases intrathoracic pressure on inspiration. The injured area may cave inward; on expiration, it moves outward.




Funnel Chest (Pectus Excavatum)

murmurs.

Depression in the lower portion of the sternum. Related compression of the heart and great vessels may cause



Cross-Section of Thorax Pigeon Chest (Pectus Carinatum) Sternum is displaced anteriorly, increasing the A-P diameter; costal cartilages adjacent to the protruding sternum are depressed.



Anteriorly displaced sternum

Thoracic Kyphoscoliosis Abnormal spinal curvatures and vertebral rotation deform the chest, making interpretation of lung findings difficult.




TABLE 8-5 Physical Findings in Selected Chest Disorders

	Trachea	Percussion Note	Breath Sounds	Transmitted Voice Sounds	Adventitious Sounds
Chronic Bronchitis Midline	Midline	Resonant	Normal	Normal	None, or wheezes, rhonchi, crackles
Left Heart Failure (early)	Midline	Resonant	Normal	Normal	Late inspiratory crackles in lower lungs; possible wheezes
Consolidation*	Midline	Dull	Bronchial	Increased†	Late inspiratory crackles
Atelectasis (Lobar)	May be shifted toward involved side	Dull	Usually absent	Usually absent	None
Pleural Effusion (large)	May be shifted away	Dull	Decreased to absent	Decreased to absent	Usually none, possible pleural rub
Pneumothorax	May be shifted away	Hyperresonant or Decreased to tympanitic absent	Decreased to absent	Decreased to absent	Possible pleural rub
COPD	Midline	Hyperresonant	Decreased to absent	Decreased	None unless bronchitis also
Asthma	Midline	Resonant to hyperresonant	May be obscured by wheezes	Decreased	Wheezes, perhaps crackles

*As in lobar pneumonia, pulmonary edema, or pulmonary hemorrhage With increased tactile fremitus, bronchophony, egophony, whispered pectoriloquy

The Cardiovascular System

THE HEALTH HISTORY

Common or Concerning Symptoms

- Chest pain
- Palpitations
- Shortness of breath: dyspnea, orthopnea, or paroxysmal nocturnal dyspnea
- Swelling or edema

As you assess reports of *chest pain or discomfort*, keep serious adverse events in mind, such as *angina pectoris*, *myocardial infarction*, or even a *dissecting aortic aneurysm*. Ask also about palpitations, orthopnea, paroxysmal nocturnal dyspnea (PND), and edema.

- Palpitations are an unpleasant awareness of the heart beat.
- Shortness of breath may represent dyspnea, orthopnea, or PND
 - *Dyspnea* is an uncomfortable awareness of breathing that is inappropriate for a given level of exertion.
 - Orthopnea is dyspnea that occurs when the patient is lying down and improves when the patient sits up. It suggests left ventricular heart failure or mitral stenosis; it also may accompany obstructive pulmonary disease.
 - *PND* describes episodes of sudden dyspnea and orthopnea that awaken the patient from sleep, usually 1 to 2 hours after going to bed, prompting the patient to sit up, stand up, or go to a window for air.

164 The Cardiovascular System

 Edema refers to the accumulation of excessive fluid in the interstitial tissue spaces; it appears as swelling. Dependent edema appears in the feet and lower legs when sitting or in the sacrum when bedridden.

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Screening for hypertension
- Screening for coronary heart disease and stroke
- Screening for dyslipidemias
- Promoting lifestyle modification and risk factor reduction

Cardiovascular disease is the leading cause of death for both men and women in the United States. *Primary prevention*, in those without evidence of cardiovascular disease, and *secondary prevention*, in those with known cardiovascular events (e.g., angina), remain important clinical priorities. Use education and counseling to help your patients maintain optimal levels of blood pressure, cholesterol, weight, and exercise and to reduce risk factors for cardiovascular disease and stroke

Screening for Hypertension. The U.S. Preventive Services Task Force recommends screening adults 18 years or older for high blood pressure. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7) has issued a simpler and more stringent blood pressure classification (see table on p. 63). Note that:

- Normal blood pressure is defined as less than 120/80 mm Hg. Lifestyle interventions should begin when blood pressure is ≥120–139/80–89, termed prehypertension.
- Adoption of healthy lifestyles by all people is considered indispensable.

Screening for Coronary Heart Disease and Stroke. The American Heart Association (AHA) recommends risk factor screening beginning at age 20 years and global absolute coronary heart disease (CHD) risk estimation for all adults 40 years or older.

Risk Assessment for Adults Beginning at Age 20 Years				
Risk Factor	Screening Frequency			
Family history of CHD	Update regularly			
Smoking status				
Diet	At each routine visit			
Alcohol intake				
Physical activity				
Blood pressure				
Body mass index (BMI)	At each routine visit (at least every 2 years)			
Waist circumference				
Pulse (to detect atrial fibrillation)	2 years)			
Fasting lipoprotein profile	At least every 5 years			
Fasting glucose	If risk factors for hyperlipidemia or diabetes present, every 2 years			

GLOBAL RISK ESTIMATION FOR 10-YEAR RISK OF CHD FOR Adults 40 Years OR Older

Establish multiple risk score for CHD based on age, sex, smoking status, systolic (and sometimes diastolic) blood pressure, total (and sometimes LDL) cholesterol, HDL cholesterol, and diabetes.

For calculation of global CHD risk, use the risk calculators found at either of the Web sites below (or other equations): www.americanheart.org/presenter.jhtml?identifier= 3003499

http://hin.nhlbi.nih.gov/atpiii/calculator.asp?usertype=prof

166 The Cardiovascular System

Screening for Dyslipidemias. LDL is the primary target of cholesterol-lowering therapy. Ten-year risk categories are as follows:

- *High risk* (10-year risk >20%): established CHD and CHD risk equivalents
- Moderately high risk (10-year risk 10% to 20%): multiple or 2+ risk factors
- Low risk (10-year risk <10%): 0 to 1 risk factor

For high-risk people, the recommended LDL goal is less than 70 mg/dL and intensive lipid therapy is a *therapeutic option*. A summary table of LDL goals based on risk is below.

Updated Adult Treatment Panel III Guidelines				
10-Year Risk Category	LDL Goal	Consider Drug Therapy		
High risk (>20%)	<100 mg/dL Optional goal: <70 mg/dL	≥100 mg/dL (<100 mg/dL: consider drug options, including further 30% to 40% reduction in LDL)		
Moderately high risk (10%–20%)	<130 mg/dL Optional goal: <100 mg/dL	≥130 mg/dL (100–129 mg/dL: consider drug options to achieve goal of <100 mg/dL)		
Moderate risk (<10%)	<130 mg/dL	≥160 mg/dL		
Lower risk (0 to 1 risk factor)	<160 mg/dL	>190 mg/dL (160–189 mg/dL: drug therapy <i>optional</i>)		

(Source: Adapted from National Cholesterol Education Panel Report. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III Guidelines. Grundy SM, Cleeman JI, Merz NB, et al., for the Coordinating Committee of the National Cholesterol Education Program. Circulation 119:227–239, 2004.)

For definitions of risk factors, CHD events, and CHD risk equivalents (such as diabetes or peripheral vascular disease), refer to Bates' Guide to Physical Examination and History Taking, 10th edition, Chapter 9.

Promoting Lifestyle Modification and Risk Factor Reduction.

The JNC 7 and AHA encourage well-studied effective lifestyle modification and risk interventions to prevent hypertension, CHD, and stroke.

LIFESTYLE MODIFICATIONS FOR CARDIOVASCULAR HEALTH

- Optimal weight (BMI of 18.5–24.9 kg/m²)
- Salt intake less than 6 g/day of sodium chloride or 2.4 g/day of sodium
- Regular aerobic exercise (e.g., brisk walking) for at least 30 min/day, most days of the week
- Moderate alcohol consumption of two or fewer drinks per day for men and one drink or fewer per day for women
- Diet rich in fruits, vegetables, and low-fat dairy products with reduced saturated and total fat
- Dietary intake of >3,500 mg of potassium
- Optimal blood pressure control (see pp. 56–57)
- Lipid management
- Diabetes management so that fasting glucose level is below 110 mg/dL and HgA1C is less than 7%
- Complete smoking cessation
- Conversion of atrial fibrillation to normal sinus rhythm or, if chronic, anticoagulation

Chapter 4 discusses healthy eating habits and dietary counseling in more detail. Remember to assess BMI and to evaluate the patient's eating habits and weight patterns in the family. To reduce risk for CHD, counsel patients to pursue aerobic exercise for at least 30 minutes on most days of the week. Spur motivation by emphasizing the immediate benefits to health and well-being.

EXAMINATION ECHNIQUES 0 F

EXAMINATION TECHNIQUES

HEART RATE AND BLOOD PRESSURE

If not already done, measure the radial or apical pulse.

Estimate systolic blood pressure by palpation and add 30 mm Hg. Use this sum as the target for further cuff inflations.

Measure blood pressure with a sphygmomanometer. If indicated, recheck it.

This step helps you to detect an auscultatory gap and avoid recording an inappropriately low systolic blood pressure.

Orthostatic (postural) hypotension with position change from supine to standing, SBP↓ ≥20 mm Hg; $HR\uparrow > 20$ beats/min

JUGULAR VEINS

Identify jugular venous pulsations and their highest point in the neck. Start with head of the bed at 30°; adjust angle of the bed as necessary.

Study the waves of venous pulsation. Note the α wave of atrial contraction and the p wave of venous filling.

Measure jugular venous pressure (JVP)—the vertical distance between this highest point and the sternal angle, normally less than 3-4 cm.

Absent a waves in atrial fibrillation; prominent v waves in tricuspid regurgitation.

Elevated IVP in right-sided heart failure; decreased JVP in hypovolemia from dehydration or gastrointestinal bleeding



POSSIBLE FINDINGS

CAROTID PULSE

Assess the amplitude and contour of the carotid upstroke.

Listen for bruits.

A delayed upstroke in aortic stenosis; a bounding upstroke in aortic insufficiency

Carotid bruits suggest atherosclerotic narrowing.

THE HEART

Sequence of the Cardiac Examination				
Patient Position	Examination			
Supine, with the head elevated 30°	Inspect and palpate the precordium: the 2nd interspaces; the right ventricle; and the left ventricle, including the apical impulse (diameter, location, amplitude, duration).			
Left lateral decubitus	Palpate the apical impulse if not previously detected. Listen at the apex with the <i>bell of the stethoscope for low-pitched extra sounds</i> (S ₃ , opening snap, diastolic rumble of mitral stenosis).			
Supine, with the head elevated 30°	Listen at the 2nd right and left interspaces, along the left sternal border, and across to the apex with the diaphragm. Listen with the bell at the right sternal border for tricuspid murmurs and sounds.			
Sitting, leaning forward, after full exhalation	Listen along the left sternal border and at the apex for the soft decrescendo diastolic murmur of aortic insufficiency.			

POSSIBLE FINDINGS

INSPECTION AND PALPATION

Inspect and **palpate** the anterior chest for heaves, lifts, or thrills.



Identify the *apical impulse*. Turn patient to left as necessary. **Note:**

- Location of impulse
- Diameter

- Amplitude—usually *tapping*
- Duration

Feel for a right ventricular impulse in left parasternal and epigastric areas.

Palpate left and right second interspaces close to sternum. Note any thrills in these areas.

Displaced to left in pregnancy

Increased diameter, amplitude, and duration in left ventricular dilatation from congestive heart failure (CHF) or ischemic cardiomyopathy

Sustained in left ventricular hypertrophy; diffuse in CHF


Prominent impulses suggest right ventricular enlargement.

Pulsations of great vessels; accentuated S₂; thrills of *aortic* or *pulmonic stenosis*

POSSIBLE FINDINGS

AUSCULTATION

Listen to heart by "inching" your stethoscope from the base to the apex (or apex to base) in the areas illustrated.



Use the *diaphragm* in the areas illustrated above for relatively *high-pitched sounds* like S_1 , S_2 .

Use the *bell* for *low-pitched sounds* at the lower left sternal border and apex.

Listen at each area for:

- \bullet S_1
- S₂. Is splitting normal in left 2nd and 3rd interspaces?
- Extra sounds in systole
- Extra sounds in diastole
- Systolic murmurs
- Diastolic murmurs

Also murmurs of aortic and mitral regurgitation; pericardial friction rubs

S₃, S₄, murmur of *mitral* stenosis

See Table 9-1, Heart Sounds, p. 176; Table 9-2, Variations in the First Heart Sound (S_1) , p. 177; Table 9-3, Variations in the Second Heart Sound (S_2) , pp. 178–179.

Physiologic (inspiratory) or pathologic (expiratory) splitting

Systolic clicks

S3, S4

Midsystolic, pansystolic, late systolic murmurs

Early, mid-, or late diastolic murmurs


EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

ASSESSING AND DESCRIBING MURMURS

Identify, if murmurs are present, their

- Timing in the cardiac cycle (systole, diastole)
- Shape



- Location of maximal intensity
- Radiation
- Pitch
- Quality
- Intensity on a 6-point scale

Listen at the apex with patient turned toward left side for low-pitched sounds.

See Table 9-4, Heart Murmurs, p. 180.

Plateau, crescendo, decrescendo

A crescendo–decrescendo murmur first rises in intensity, then falls.

A *plateau murmur* has the same intensity throughout.

A *crescendo murmur* grows louder.

A decrescendo murmur grows softer.

High, medium, low

Blowing, harsh, musical, rumbling

See "Gradations of Murmurs" on next page.

Left-sided S₃, S₄, and diastolic murmur of *mitral stenosis*

POSSIBLE FINDINGS



7 **Listen** down left sternal border to the apex as patient sits, leaning forward, with breath held after exhalation.

Diastolic decrescendo murmur of aortic regurgitation



Gradations of Murmurs		
Grade	Description	
Grade 1	Very faint, heard only after listener has "tuned in"; may not be heard in all positions	
Grade 2	Quiet, but heard immediately after placing the stethoscope on the chest	
Grade 3	Moderately loud	
Grade 4	Loud, with palpable thrill	
Grade 5	Very loud, with thrill. May be heard when the stethoscope is partly off the chest	
Grade 6	Very loud, with thrill. May be heard with stethoscope entirely off the chest	

SPECIAL TECHNIQUES

PULSUS ALTERNANS

Feel pulse for alternation in amplitude. Lower pressure of blood pressure cuff slowly to systolic level while you listen with stethoscope over brachial artery. Alternating amplitude of pulse or sudden doubling of Korotkoff sounds indicates pulsus alternans—a sign of left ventricular heart failure.

PARADOXICAL PULSE

Lower pressure of blood pressure cuff slowly and note two pressure levels: (1) where Korotkoff sounds are first heard and (2) where they first persist through the respiratory cycle. These levels are normally not more than 3–4 mm Hg apart.

A drop of greater than 10 mm Hg during inspiration signifies a paradoxical pulse. Consider obstructive pulmonary disease, pericardial tamponade, or constrictive pericarditis.

AIDS TO IDENTIFY SYSTOLIC MURMURS

Valsalva Maneuver

Ask patient to strain down.

In suspected *mitral valve prolapse (MVP)*, listen to the timing of click and murmur.

Ventricular filling decreases, the systolic click of MVP is earlier, and the murmur lengthens.

To distinguish aortic stenosis (AS) from hypertrophic cardiomyopathy (HC), listen to the intensity of the murmur.

In AS, the murmur decreases; in HC, it often increases.

Squatting and Standing

In suspected MVP, listen for the click and murmur in both positions.

Try to distinguish AS from HC by listening to the murmur in both positions.

Squatting increases ventricular filling and delays the click and murmur. Standing reverses the changes.

Squatting increases murmur of AS and decreases murmur of HC. Standing reverses the changes.

RECORDING YOUR FINDINGS

Recording the Physical Examination— The Cardiovascular Examination

"The jugular venous pulse (JVP) is 3 cm above the sternal angle with the head of the bed elevated to 30°. Carotid upstrokes are brisk, without bruits. The point of maximal impulse (PMI) is tapping, 7 cm lateral to the midsternal line in the 5th intercostal space. Crisp S₁ and S₂. At the base, S_2 is greater than S_1 and physiologically split, with $A_2 > P_2$. At the apex, S₁ is greater than S₂ and constant. No murmurs or extra sounds."

OR


"The JVP is 5 cm above the sternal angle with the head of the bed elevated to 50°. Carotid upstrokes are brisk; a bruit is heard over the left carotid artery. The PMI is diffuse, 3 cm in diameter, palpated at the anterior axillary line in the 5th and 6th intercostal spaces. S₁ and S₂ are soft. S₃ present at the apex. High-pitched, harsh 2/6 holosystolic murmur best heard at the apex, radiating to the axilla. No S₄ or diastolic murmurs." (Suggests CHF with possible left carotid occlusion and mitral regurgitation)

INTERPRETATION

TABLE 9-1


Finding

Heart Sounds



9		
S ₁ accentuated	Tachycardia, states of high cardiac output; mitral stenosis	
S ₁ diminished	First-degree heart block; reduced left ventricular contractility; immobile mitral valve, as in mitral regurgitation	
Systolic clicks(s)	Mitral valve prolapse	
S ₂ accentuated in right 2nd interspace	Systemic hypertension, dilated aortic root	
S ₂ diminished or absent in right 2nd interspace	Immobile aortic valve, as in calcific aortic stenosis	
P ₂ accentuated	Pulmonary hypertension, dilated pulmonary artery, atrial septal defect	
P ₂ diminished or absent	Aging, pulmonic stenosis	
Opening snap	Mitral stenosis	
S ₃	Physiologic (usually in children and young adults); volume overload of ventricle, as in mitral regurgitation or congestive heart failure	
S ₄	Excellent physical conditioning (trained athletes); resistance to ventricular filling because of decreased compliance, left ventricular hypertrophy from pressure overload, as in hypertensive heart disease, aortic stenosis	

Variations in the First Heart Sound (S₁)






Variations in the Second Heart Sound (S₂)

Inspiration

Expiration

Physiologic Splitting





Heard in the 2nd or 3rd left interspace: the pulmonic component of S_2 is usually too faint to be heard at the apex or aortic area, where S_2 is single and derived from aortic valve closure alone. Accentuated by inspiration; usually disappears on exertion.

Pathologic Splitting









Wide splitting of S₂ persists throughout respiration; from delayed closure of the pulmonic valve (e.g., by pulmonic stenosis or right bundle branch block); also from early closure of the aortic valve, as in mitral regurgitation.

Fixed Splitting









Does not vary with respiration, as in atrial septal defect, right ventricular failure.



Variations in the Second Heart Sound (S₂) (continued)

Inspiration

Expiration

Paradoxical or Reversed Splitting





Appears on expiration and disappears on inspiration. Closure of the aortic valve is abnormally delayed, so A_2 follows P_2 on expiration, as in left bundle branch block.

More on A₂ and P₂

Increased Intensity of A_2 , Second Right Interspace (where only A_2 can usually be heard) occurs in systemic hypertension because of the increased pressure. It also occurs when the aortic root is dilated, probably because the aortic valve is then closer to the chest wall.

Decreased or Absent A_2 , Second Right Interspace is noted in calcific aortic stenosis because of immobility of the valve. If A_2 is inaudible, no splitting is heard.

Increased Intensity of P₂. When P₂ is equal to or louder than A₂, pulmonary hypertension may be suspected. Other causes include a dilated pulmonary artery and an atrial septal defect. Splitting of the second heart sound that is heard widely, even at the apex and the right base, indicates an accentuated P₂.

Decreased or Absent P₂ is most commonly due to the increased anteroposterior diameter of the chest associated with aging. It can also result from pulmonic stenosis. If P₂ is inaudible, no splitting is heard.

Heart Murmurs

Likely Causes

Midsystolic



Innocent murmurs (no valve abnormality) Physiologic murmurs (from ↑ flow across a semilunar valve, as in pregnancy, fever, anemia)

Aortic stenosis

Murmurs that mimic aortic stenosis aortic sclerosis, bicuspid aortic valve, dilated aorta, and pathologically ↑ systolic flow across aortic valve Hypertrophic cardiomyopathy Pulmonic stenosis

Pansystolic



Mitral regurgitation Tricuspid regurgitation Ventricular septal defect

Late Systolic

Mitral valve prolapse, often with click (C)

Early Diastolic



Aortic regurgitation

Middiastolic

and Presystolic



Mitral stenosis—note opening snap (OS)

Continuous Murmurs and Sounds



Patent ductus arteriosus—harsh, machinery-like



Pericardial friction rub—a scratchy sound with 1–3 components



Venous hum—continuous, above midclavicles, loudest in diastole

10

The Breasts and Axillae

THE HEALTH HISTORY

Common or Concerning Symptoms

- Breast lump or mass
- Breast pain or discomfort
- Nipple discharge

Ask, "Do you examine your breasts?" . . . "How often?" Ask about any discomfort, pain, or lumps in the breasts. Also ask about any discharge from the nipples, change in breast contour, dimpling, swelling, or puckering of the skin over the breasts.

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Palpable masses of the breast
- Assessing risk factors for breast cancer
- Breast cancer screening
- Breast self-examination (BSE)

Palpable Masses of the Breast. Breast masses show marked variation in etiology, from fibroadenomas and cysts seen in younger women, to abscess or mastitis, to primary breast cancer. All breast masses warrant careful evaluation, and definitive diagnostic measures should be pursued.

Assessing Risk Factors for Breast Cancer. Although 70% of affected women have no known predisposing factors, definite

Palpable Masses of the Breast			
Age	Common Lesion	Characteristics	
15–25	Fibroadenoma	Usually fine, round, mobile, nontender	
25–50	Cysts	Usually soft to firm, round, mobile; often tender	
	Fibrocystic changes	Nodular, ropelike	
	Cancer	Irregular, stellate, firm, not clearly delineated from surrounding tissue	
50 or older	Cancer until proven otherwise	As above	
Pregnancy/ lactation	Lactating adenomas, cysts, mastitis, and cancer	As above	

(Adapted from Schultz MZ, Ward BA, Reiss M. Breast diseases. In: Noble J, Greene HL, Levinson W, et al., eds: Primary Care Medicine, 2nd ed. St. Louis: Mosby, 1996. See also Venet L, Strax P, Venet W, et al. Cancer 28(6):1546–1551, 1971.)

risk factors are well established. Use the Breast Cancer Risk Assessment Tool of the National Cancer Institute (http://brca.nci.nih/gov/brc/start/htm) or other available clinical models, such as the Gail model, to individualize risk factor assessment for your patients. Ask women beginning in their 20s about any family history of breast or ovarian cancer, or both, on the maternal or paternal side, to help assess risk of BRCA1 or BRCA2 gene mutation. See Table 10-1, Breast Cancer: Factors That Increase Relative Risk, p. 190.

Breast Cancer Screening. The American Cancer Society recommends:


- Clinical breast examination (CBE) by a health care professional every 3 years for women between 20 and 39 years of age, and annually after 40 years of age
- Yearly mammography for women 40 years of age and older.
 For women at increased risk, many clinicians advise

initiating screening mammography between ages 30 and 40, then every 2 to 3 years until 50 years of age.

• Regular breast self-examination (BSE) to help promote health awareness

ECHNIQUES OF EXAMINAT

EXAMINATION TECHNIQUES



THE FEMALE BREAST

Inspect the breasts in four positions.

Note:

Size and symmetry

Development, asymmetry

Contour

Flattening, dimpling

EXAMINATION TECHNIQUES

• Appearance of the skin

.,...



ARMS AT SIDES



HANDS PRESSED AGAINST HIPS

Inspect the nipples.

- Compare their size, shape, and direction of pointing.
- Note any rashes, ulcerations, or discharge.

O Palpate the breasts, including augmented breasts. Breast tissue should be flattened and the patient supine. Palpate a rectangular area extending from the clavicle to the inframammary fold or bra line, and from the midsternal line to the posterior axillary line and well into the axilla for the tail of Spence.

POSSIBLE FINDINGS

Edema (peau d'orange) in breast cancer



ARMS OVER HEAD



LEANING FORWARD

Inversion, retraction, deviation

Paget's disease of the nipple, galactorrhea

POSSIBLE FINDINGS

Note:

- Consistency
- Tenderness
- Nodules. If present, note location, size, shape, consistency, delimitation, tenderness, and mobility.

Use *vertical strip pattern* (currently the best validated technique) or a circular or wedge pattern. Palpate in *small, concentric circles.*

- For the lateral portion of the breast, ask the patient to roll onto the opposite hip, place her hand on her forehead, but keep shoulders pressed against the bed or examining table.
- For the medial portion of the breast, ask the patient to lie with her shoulders flat against the bed or examining table, place her hand at her neck, and lift up her elbow until it is even with her shoulder.

Palpate each nipple.

Palpate and inspect along the incision lines of mastectomy.

Physiologic nodularity

Infection, premenstrual tenderness

Cyst, fibroadenoma, cancer





Thickening in cancer

Local recurrences of breast cancer.

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

THE MALE BREAST

Inspect and palpate the nipple and areola.

Gynecomastia, mass suspicious for cancer, fat


AXILLAE

Inspect for rashes, infection, and pigmentation.

Palpate the axillary nodes, including the central, pectoral, lateral, and subscapular groups.

Hidradenitis suppurativa, acanthosis nigricans

Lymphadenopathy



ARROWS INDICATE DIRECTION OF LYMPH FLOW

SPECIAL TECHNIQUE

O BREAST DISCHARGE

Compress the areola in a spokelike pattern around the nipple. Watch for discharge. Type and source of discharge may be identified




BREAST SELE-EXAMINATION

PATIENT INSTRUCTIONS FOR THE BREAST SELF-EXAMINATION (BSE)

Supine



- 1. Lie down with a pillow under your right shoulder. Place your right arm behind your head.
- 2. Use the finger pads of the three middle fingers on vour left hand to feel for lumps in the right breast. The finger pads are the top third of each finger.
- 3. Press firmly enough to know how your breast feels. A firm ridge in the lower curve of each breast is normal. If vou're not sure how hard to press, talk with your health care provider, or try to copy the way the doctor or nurse does it.
- 4. Press firmly on the breast in an up-and-down or



"strip" pattern. You can also use a circular or wedge pattern, but be sure to use the same pattern every time. Check the entire breast area, and remember how your breast feels from month to month.

- 5. Repeat the examination on your left breast, using the finger pads of the right hand.
- 6. If you find any changes, see your doctor right away.

(continued)

PATIENT INSTRUCTIONS FOR THE BREAST SELF-EXAMINATION (BSE) (CONTINUED)

Standing



1. Repeat the examination of both breasts while standing, with one arm behind your head. The upright position makes it easier to check the upper outer part of the breasts (toward your armpit). This is where about half of breast cancers are found. You may want to do the upright part of the BSE while you are in the shower. Your soapy hands will make it easy to check how your breasts



feel as they glide over the wet skin.

- 2. For added safety, you might want to check your breasts by standing in front of a mirror right after your BSE each month. See if there are any changes in the way your breasts look, such as dimpling of the skin, changes in the nipple, redness, or swelling.
- 3. If you find any changes, see your doctor right away.

(Adapted from the American Cancer Society, www.cancer.org. Accessed October 24, 2007.)

RECORDING YOUR FINDINGS

Recording the Physical Examination— Breasts and Axillae

"Breasts symmetric and smooth, without masses. Nipples without discharge." (Axillary adenopathy usually included after Neck in section on Lymph Nodes; see p. 123.)

OR

TABLE 10

"Breasts pendulous with diffuse fibrocystic changes. Single firm 1 × 1 cm mass, mobile and nontender, with overlying peau d'orange appearance in right breast, upper outer quadrant at 11 o'clock, 2 cm from the nipple."

Breast Cancer in Women: Factors That

(table continues next page)

(Suggests possible breast cancer)

INTERPRETATION AIDS

Relative Risk	Factor
>4.0	 Female Age (65+ versus <65 years, although risk increases across all ages until age 80) Certain inherited genetic mutations for breast cancer (BRCA1 and/or BRCA2) Two or more first-degree relatives with breast cancer diagnosed at an early age Personal history of breast cancer High breast tissue density Biopsy-confirmed atypical hyperplasia
2.1–4.0	 One first-degree relative with breast cancer High-dose radiation to chest High bone density (postmenopausal)

Increase Relative Risk

TABLE 10-1 Increase Relative Risk (continued)		
Relative Risk	Factor	
1.1–2.0 Factors that affect circulating hormones	 Late age at first full-term pregnancy (>30 years) Early menarche (<12 years) Late menopause (>55 years) No full-term pregnancies Never breast-fed a child Recent oral contraceptive use Recent and long-term use of hormone replacement therapy Obesity (postmenopausal) 	
Other factors	 Personal history of endometrium, ovary, or colon cancer Alcohol consumption Height (tall) 	

Breast Cancer in Women: Factors That

(Source: American Cancer Society. Breast Cancer Facts and Figures 2007–2008, p. 10. Available at: www.cancer.org/downloads/STT/BCFF-Final.pdf. Accessed October 20, 2007.)

• Jewish heritage


• High socioeconomic status

TABLE 10-2 Visible Signs of Breast Cancer

Retraction Signs

Fibrosis from breast cancer produces retraction signs: dimpling, changes in contour, and retraction or deviation of the nipple. Other causes of retraction include fat necrosis and mammary duct ectasia.

Skin Dimpling





Abnormal Contours

Look for any variation in the normal convexity of each breast, and compare one side with the other.



Nipple Retraction and Deviation

A retracted nipple is flattened or pulled inward. It may also be broadened and feel thickened. The nipple may deviate, or point in a different direction, typically toward the underlying cancer.



(table continues next page)

Visible Signs of Breast Cancer (continued)


Edema of the Skin

From lymphatic blockade, appearing as thickened skin with enlarged pores—the socalled peau d'orange (orange peel) sign.



Paget's Disease of the Nipple

An uncommon form of breast cancer that usually starts as a scaly, eczema-like lesion. The skin may also weep, crust, or erode. A breast mass may be present. Suspect Paget's disease in any persisting dermatitis of the nipple and areola.



The Abdomen

THE HEALTH HISTORY

Common or Concerning Symptoms

Gastrointestinal Disorders

- Abdominal pain, acute and chronic
- Indigestion, nausea, vomiting including blood, loss of appetite, early satiety
- Dysphagia and/or odynophagia
- Change in bowel function
- Diarrhea, constipation
- laundice

Urinary and Renal Disorders

- Suprapubic pain
- Dysuria, urgency, or frequency
- Hesitancy, decreased stream in males
- Polyuria or nocturia
- Urinary incontinence
- Hematuria
- Kidney or flank pain
- Ureteral colic

PATTERNS AND MECHANISMS OF ABDOMINAL PAIN

Be familiar with three broad categories:

Visceral pain—occurs when hollow abdominal organs such as the intestine or biliary tree contract unusually forcefully or are distended or stretched.

Visceral pain in the right upper quadrant (RUQ) from liver distention against its capsule in alcoholic hepatitis

194 The Abdomen

- May be difficult to localize
- Varies in quality; may be gnawing, burning, cramping, or aching
- When severe, may be associated with sweating, pallor, nausea, vomiting, restlessness.

Parietal pain—from inflammation of the parietal peritoneum

- Steady, aching
- Usually more severe
- Usually more precisely localized over the involved structure than visceral pain

Referred pain—occurs in more distant sites innervated at approximately the same spinal levels as the disordered structure.

Pain from the chest, spine, or pelvis may be referred to the abdomen.

Visceral periumbilical pain in early acute appendicitis from distention of inflamed appendix gradually changes to parietal pain in the right lower quadrant (RLQ) from inflammation of the adjacent parietal peritoneum.

Pain of duodenal or pancreatic origin may be referred to the back; pain from the biliary tree—to the right shoulder or right posterior chest.

Pain from *pleurisy* or *acute* myocardial infarction may be referred to the upper abdomen.

THE GASTROINTESTINAL TRACT

Ask patients to describe the abdominal pain in their own words, especially timing of the pain (acute or chronic); then ask them to point to the pain.

Pursue important details: "Where does the pain start?"

"Does it radiate or travel?" "What is the pain like?" "How severe is it?" "How about on a scale of 1 to 10?" "What makes it better or worse?"

Elicit any symptoms associated with the pain, such as fever or chills; ask their sequence.

Upper Abdominal Pain or Discomfort. Ask about chronic or recurrent upper abdominal discomfort, or dyspepsia. Related symptoms include bloating, nausea, upper abdominal fullness, and heartburn.

Find out just what your patient means. Possibilities include:

- Bloating from excessive gas, especially with frequent belching, abdominal distention, or flatus, the passage of gas by rectum
- Nausea and vomiting
- Unpleasant abdominal fullness after normal meals or early satiety, the inability to eat a full meal
- Hearthurn

Consider diabetic gastroparesis, anticholinergic drugs, gastric outlet obstruction, gastric cancer. Early satiety may signify hepatitis.

Suggests gastroesophageal reflux disease (GERD)

Lower Abdominal Pain or Discomfort

If acute, is the pain sharp and continuous or intermittent and cramping?

Right lower quadrant (RLQ) pain, or pain migrating from periumbilical region in appendicitis; in women with RLQ pain, possible pelvic inflammatory disease, ectopic pregnancy

Left lower quadrant (LLQ) pain in *diverticulitis*

If chronic, is there a change in bowel habits? Alternating diarrhea and constipation? Colon cancer; irritable bowel syndrome

Other GI Symptoms

Anorexia

Liver disease, pregnancy, diabetic ketoacidosis, adrenal insufficiency, uremia, anorexia nervosa

Dysphagia or difficulty swallowing

Of solids and liquids: usually neuromuscular disorders affecting motility

If only solids, consider structural conditions like Zenker's diverticulum, Schatzki's ring, stricture, neoplasm

Odynophagia, or painful swallowing

Radiation; caustic ingestion, infection from *cytomegalovirus*, herpes simplex, HIV

 Diarrhea, acute (<2 weeks) and chronic Acute infection (viral, salmonella, shigella, etc.); chronic in *Crohn's disease*, ulcerative colitis; if oily (*steatorrhea*)—pancreatic insufficiency. See Table 11-1.

- Constipation
- Melena, or black tarry stools
- Jaundice from increased levels of bilirubin: intrahepatic jaundice can be hepatocellular, from damage to the hepatocytes, or cholestatic, from impaired excretion caused by damaged hepatocytes or intrahepatic bile ducts

Extrahepatic jaundice from obstructed extrahepatic bile ducts, commonly the cystic and common bile ducts

Ask about the color of the urine and stool.

Medications, especially anticholinergic agents and opioids; colon cancer

GI bleed

Impaired excretion of conjugated bilirubin in viral hepatitis, cirrhosis, primary biliary cirrhosis, drug-induced cholestasis

Dark urine from increased conjugated bilirubin in serum excreted in urine; acholic claycolored stool when excretion of bilirubin into intestine is obstructed

RISK FACTORS FOR LIVER DISEASE

- Hepatitis A: Travel or meals in areas with poor sanitation, ingestion of contaminated water or foodstuffs
- Hepatitis B: Parenteral or mucous membrane exposure to infectious body fluids such as blood, serum, semen, and saliva, especially through sexual contact with an infected partner or use of shared needles for injection drug use
- Hepatitis C: Illicit intravenous drug use or blood transfusion
- Alcoholic hepatitis or alcoholic cirrhosis: Interview the patient carefully about alcohol use

(continued)

RISK FACTORS FOR LIVER DISEASE (CONTINUED)

- Toxic liver damage from medications, industrial solvents, or environmental toxins
- Extrahepatic biliary obstruction: Resulting from gallbladder disease or surgery
- Hereditary disorders in the Family History

THE URINARY TRACT

Ask about *pain on urination*, usually a burning sensation, sometimes termed *dysuria* (also refers to difficulty voiding).

Other associated symptoms include

- Urgency, an unusually intense and immediate desire to void
- Urinary frequency, or abnormally frequent voiding
- Fever or chills; blood in the urine
- Any pain in the abdomen, flank, or back

In men *hesitancy* in starting the urine stream, *straining* to void, reduced caliber and force of the urine stream, or dribbling as they complete voiding

Bladder infection

Also, consider bladder stones, foreign bodies, tumors, and acute prostatitis. In women, internal burning in urethritis, external burning in vulvovaginitis

May lead to urge incontinence

Dull, steady pain in pyelonephritis; severe colicky pain in ureteral obstruction from renal stone

Prostatitis, urethritis

Assess any:

- *Polyuria*, a significant increase in 24-hour urine volume
- *Nocturia*, urinary frequency at night
- Urinary incontinence, involuntary loss of urine
 - From coughing, sneezing, lifting
 - From urge to void
 - From bladder fullness. with leaking but incomplete emptying

Diabetes mellitus, diabetes insipidus

Bladder obstruction

See Table 11-2, Urinary Incontinence, pp. 211–212.

Stress incontinence (poor urethral sphincter tone)

Urge incontinence (detrusor overactivity)

Overflow incontinence (anatomic obstruction, impaired neural innervation to bladder)

Important Topics for Health Promotion and Counseling

- Screening for alcohol abuse
- Risk factors for hepatitis A, B, and C
- Screening for colon cancer

Assessing use of alcohol is an important clinician responsibility. Focus on detection, counseling, and, for significant impairment, specific treatment recommendations. Use the four CAGE questions to screen for alcohol dependence or abuse in all adolescents and adults, including pregnant women (see Chap. 3, p. 41). Brief counseling interventions have been shown to reduce alcohol consumption by 13% to 34% over 6 to 12 months.

Protective measures against infectious hepatitis include counseling about transmission:

• Hepatitis A: Transmission is fecal-oral. Illness occurs approximately 30 days after exposure. Hepatitis A

vaccine is recommended for travelers to endemic areas; food handlers; military personnel; caretakers of children; Native Americans and Alaska Natives; selected health care, sanitation, and laboratory workers; homosexual men; and injection drug users.

- Hepatitis B: Transmission occurs during contact with infected body fluids, such as blood, semen, saliva, and vaginal secretions. Infection increases risk of fulminant hepatitis, chronic infection, and subsequent cirrhosis and hepatocellular carcinoma. Provide counseling and serologic screening for patients at risk. Hepatitis B vaccine is recommended for all young adults not previously immunized, injection drug users and their sexual partners, people at risk for sexually transmitted diseases, travelers to endemic areas, recipients of blood products as in hemodialysis, and health care workers with frequent exposure to blood products. Many of these groups also should be screened for HIV infection, especially pregnant women at their first prenatal visit.
- *Hepatitis C:* Hepatitis C, now the most common form, is spread by blood exposure and is associated with injection drug use. No vaccine is available.

For colorectal cancer, screen for risk factors, namely family history of colonic polyps, history of colorectal cancer or adenoma in a first-degree relative, and personal history of ulcerative colitis, adenomatous polyps, or prior diagnosis of endometrial, ovarian, or breast cancer. For men and women at average risk, the American Cancer Society recommends one of the following beginning at age 50:

- Fecal occult blood test (FOBT) annually
- Flexible sigmoidoscopy every 5 years
- Annual FOBT plus flexible sigmoidoscopy every 5 years
- Double contrast barium enema every 5 years
- Colonoscopy every 10 years

Detection rates for colorectal cancer and insertion depths of endoscopy are roughly as follows: 25% to 30% at 20 cm; 50% to 55% at 35 cm; 40% to 65% at 40 cm to 50 cm. Full colonoscopy or air contrast barium enema detects 80% to 95% of colorectal cancers.

TECHNIQUES OF EXAMINATION

EXAMINATION TECHNIQUES

THE ABDOMEN

• Inspect the abdomen, including

Skin Scars, striae, veins

 Umbilicus Hernia, inflammation

• Contours for shape, Bulging flanks, suprapubic symmetry, enlarged organs bulge, large liver or spleen, or masses tumors

 Any peristaltic waves GI obstruction

 Any pulsations Increased in aortic aneurysm

Auscultate the abdomen for

 Bowel sounds Increased or decreased motility

Bruit of renal artery stenosis Bruits

 Friction rubs Liver tumor, splenic infarct

Bowel Sounds and Bruits			
Change	Seen With		
Increased bowel sounds	Diarrhea Early intestinal obstruction		
Decreased, then absent bowel sounds	Adynamic ileus Peritonitis		
High-pitched tinkling bowel sounds	Intestinal fluid Air under tension in a dilated bowel		
	(continued)		

Bowel Sounds and Bruits (continued)				
Change	Seen With			
High-pitched rushing bowel sounds with cramping	Intestinal obstruction			
Hepatic bruit	Carcinoma of the liver Alcoholic hepatitis			
Arterial bruits	Partial obstruction of the aorta or renal arteries			
Aorta Renal artery Iliac artery Femoral artery				

Percuss the abdomen for patterns of tympany and dullness.

Palpate all quadrants of the abdomen

• Lightly for guarding, rebound, and tenderness



Ascites, GI obstruction, pregnant uterus, ovarian tumor

See Table 11-3.

Firm, boardlike abdominal wall suggests peritoneal inflammation. Guarding occurs when the patient flinches, grimaces, or reports pain during palpation.

Rebound tenderness from peritoneal inflammation; pain is greater when you withdraw

POSSIBLE FINDINGS

your hand than when you press down. Press slowly on a tender area, then quickly "let go."


 Deeply for masses or tenderness

Tumors, a distended viscus



THE LIVER

Percuss span of liver dullness Hepatomegaly in the midclavicular line (MCL).



Feel the liver edge, if possible, as patient breathes in.

Measure its distance from the costal margin in the MCL.

Firm edge of cirrhosis

Increased in hepatomegaly may be missed (as below) by starting palpation too high in the RUO





Note any tenderness or masses.

Tender liver of hepatitis or congestive heart failure; tumor mass

THE SPLEEN

Percuss across left lower anterior chest, noting change from tympany to dullness.

Try to **feel** spleen with the patient

- Supine
- Lying on the right side with legs flexed at hips and knees

Splenomegaly



POSSIBLE FINDINGS

THE KIDNEYS

— Try to palpate each kidney.



Enlargement from cysts, cancer, hydronephrosis

Check for costovertebral angle (CVA) tenderness.

Tender in kidney infection



THE AORTA

• Palpate the aorta's pulsations. In older people, estimate its width.





Periumbilical mass with expansile pulsations ≥3 cm in diameter in abdominal aortic aneurysm. Assess further due to risk of rupture.

ASSESSING ASCITES

•—/• Palpate for shifting dullness. Map areas of tympany and dullness with patient supine, then lying on side (see below).

Ascitic fluid usually shifts to dependent side, changing the margin of dullness (see below)





o— Check for a fluid wave. Ask patient or an assistant to press edges of both hands into midline of abdomen. Tap one side and feel for a wave transmitted to the other side.

A palpable wave suggests but does not prove ascites.



POSSIBLE FINDINGS

 Ballotte an organ or mass in an ascitic abdomen. Place your stiffened and straightened fingers on the abdomen, briefly jab them toward the structure, and try to touch its surface.

Your hand, quickly displacing the fluid, stops abruptly as it touches the solid surface.



ASSESSING POSSIBLE APPENDICITIS

Ask. In classic appendicitis:

"Where did the pain begin?" Near the umbilicus

"Where is it now?" Right lower quadrant (RLQ)

Ask patient to cough. **RLQ** "Where does it hurt?"

Palpate for local tenderness. **RLQ** tenderness

Palpate for muscular rigidity. **RLQ** rigidity

Perform a rectal examination and, in women, a pelvic examination (see Chaps. 14 and 15).

- Rovsing's sign: Press deeply and evenly in the left lower quadrant. Then quickly withdraw your fingers.
- Psoas sign: Place your hand just above the patient's right knee. Ask the patient to raise that thigh against your hand. Or, ask the

Local tenderness, especially if appendix is retrocecal

Pain in the right lower quadrant during leftsided pressure suggests appendicitis (a positive Rovsing's sign).

Pain from irritation of the psoas muscle suggests an inflamed appendix (a positive psoas sign).

patient to turn onto the left side. Then extend the patient's right leg at the hip to stretch the psoas muscle.

 Obturator sign: Flex the patient's right thigh at the hip, with the knee bent, and rotate the leg internally at the hip, which stretches the internal obturator muscle. Right hypogastric pain in a positive obturator sign, suggesting irritation of the obturator muscle by an inflamed appendix.

ASSESSING POSSIBLE ACUTE CHOLECYSTITIS

Auscultate, percuss, and palpate the abdomen for tenderness.

Assess for *Murphy's sign*. Hook your thumb under the right costal margin at edge of rectus muscle, and ask patient to take a deep breath.

Bowel sounds may be active or decreased; tympany may increase with an ileus; RUQ tenderness.

Sharp tenderness and a sudden stop in inspiratory effort constitute a *positive* Murphy's sign.

RECORDING YOUR FINDINGS

Recording the Physical Examination—The Abdomen

"Abdomen is protuberant with active bowel sounds. It is soft and nontender; no masses or hepatosplenomegaly. Liver span is 7 cm and in the right MCL; edge is smooth and palpable 1 cm below the right costal margin. Spleen and kidneys not felt. No CVA tenderness."

OR

"Abdomen is flat. No bowel sounds heard. It is firm and boardlike, with increased tenderness, guarding, and rebound in the right midquadrant. Liver percusses to 7 cm in the MCL; edge not felt. Spleen and kidneys not felt. No palpable mass. No. CVA tenderness." (Suggests peritonitis from possible appendicitis; see pp. 207–208.)

AIDS TO INTERPRETATION

Diarrhea

Problem/Process

Characteristics of Stool

Acute Diarrhea

Secretory Infections

Infection by viruses; preformed bacterial toxins such as Staphylococcus aureus, Clostridium perfringens, toxigenic Escherichia coli; Vibrio cholerae. Cryptosporidium, Giardia lamblia

Watery, without blood, pus, or mucus

Inflammatory Infections

Colonization or invasion of intestinal mucosa as in nontyphoid Salmonella, Shigella, Yersinia, Campylobacter, enteropathic E. coli, Entamoeba histolytica Loose to watery, often with blood, pus, or mucus

Drug-induced Diarrhea

Action of many drugs, such as magnesium-containing antacids, antibiotics, antineoplastic agents, and laxatives

Loose to watery

Chronic Diarrhea

Diarrheal Syndromes

- Irritable bowel syndrome: A disorder of bowel motility with alternating diarrhea and constipation
- Cancer of the sigmoid colon: Partial obstruction by a malignant neoplasm

Loose; may show mucus but no blood. Small, hard stools with constipation

May be blood streaked

TABLE 11-1

Diarrhea (continued)

Problem/Process

1 TODICITI/TTOCC33

- Inflammatory Bowel Disease
- Ulcerative colitis: inflammation and ulceration of the mucosa and submucosa of the rectum and colon
- Crolm's disease of the small bowel (regional enteritis) or colon (granulomatous colitis): chronic inflammation of the bowel wall, typically involving the terminal ileum, proximal colon, or both

Voluminous Diarrheas

- Malabsorption syndromes:
 Defective absorption of fat, including fat-soluble vitamins, with steatorrhea (excessive excretion of fat) as in pancreatic insufficiency, bile salt deficiency, bacterial overgrowth
- Osmotic diarrheas
 - Lactose intolerance: Deficiency in intestinal lactase
 - Abuse of osmotic purgatives: Laxative habit, often surreptitious
- Secretory diarrheas from bacterial infection, secreting villous adenoma, fat or bile salt malabsorption, hormonemediated conditions (gastrin in Zollinger–Ellison syndrome, vasoactive intestinal peptide [VIP]): Process is variable

Characteristics of Stool

- From soft to watery, often containing blood
- Small, soft to loose or watery, usually free of gross blood (enteritis) or with less bleeding than ulcerative colitis (colitis)
- Typically bulky, soft, light yellow to gray, mushy, greasy or oily, and sometimes frothy; particularly foul smelling; usually floats in the toilet
- Watery diarrhea of large
- Watery diarrhea of large volume
- Watery diarrhea of large volume

Urinary Incontinence

Problem

Stress Incontinence: Urethral sphincter weakened. Transient increases in intra-abdominal pressure raise bladder pressure to levels exceeding urethral resistance. Leads to voiding *small amounts* during laughing, coughing, and sneezing.

Urge Incontinence: Detrusor contractions are stronger than normal and overcome normal urethral resistance. Bladder is typically small. Results in voiding *moderate amounts*, urgency, frequency, and nocturia.

Overflow Incontinence: Detrusor contractions are insufficient to overcome urethral resistance. Bladder is typically large, even after an effort to void, leading to continuous dribbling.

Mechanisms

In women, weakness of the pelvic floor with inadequate muscular support of the bladder and proximal urethra and a change in the angle between the bladder and the urethra from childbirth, surgery, and local conditions affecting the internal urethral sphincter, such as postmenopausal atrophy of the mucosa and urethral infection

In men, prostatic surgery

- Decreased cortical inhibition of detrusor contractions, as in stroke, brain tumor, dementia, and lesions of the spinal cord above the sacral level
- Hyperexcitability of sensory pathways, as in bladder infection, tumor, and fecal impaction
- Deconditioning of voiding reflexes, caused by frequent voluntary voiding at low bladder volumes
- Obstruction of the bladder outlet, as by benign prostatic hyperplasia or tumor

TABLE 11-2 Urinary Incontinence (continued)

Problem

Mechanisms

Functional Incontinence:
Inability to get to the toilet in time because of impaired health or environmental conditions

Incontinence Secondary to Medications: Drugs may contribute to any type of incontinence listed.

- Weakness of detrusor muscle associated with peripheral nerve disease at the sacral level
- Impaired bladder sensation that interrupts the reflex arc, as in diabetic neuropathy


Problems in mobility from weakness, arthritis, poor vision, other conditions; environmental factors such as unfamiliar setting, distant bathroom facilities, bed rails, physical restraints

Sedatives, tranquilizers, anticholinergics, sympathetic blockers, potent diuretics




Visceral Tenderness

Peritoneal Tenderness



Tenderness From Disease in the Chest and Pelvis

Acute Pleurisy Acute Salpingitis



The Peripheral Vascular System

THE HEALTH HISTORY

Common or Concerning Symptoms

- Pain in the arms or legs
- Intermittent claudication
- Cold, numbness, pallor in the legs, hair loss
- Color change in fingertips or toes in cold weather
- Swelling in calves, legs, or feet
- Swelling with redness or tenderness

Ask about any pain in the arms and legs.

Is there *intermittent* claudication, exercise-induced pain that is absent at rest, makes the patient stop exertion, and abates within about 10 minutes? Ask "Have you ever had any pain or cramping in your legs when you walk or exercise?" "How far can you walk without stopping to rest?" and "Does pain improve with rest?"

Peripheral arterial disease (PAD) can cause symptomatic limb ischemia with exertion; distinguish this from spinal stenosis, which produces leg pain with exertion that may be reduced by leaning forward (stretching the spinal cord in the narrowed vertebral canal) and less readily relieved by rest.

216 The Peripheral Vascular System

Ask also about *coldness*, *numbness*, or *pallor* in legs or feet or *hair loss* over the anterior tibial surfaces.

Because patients have few symptoms, identify background risk factors tobacco abuse, hypertension, diabetes, hyperlipidemia, and history of myocardial infarction or stroke.

"Do your fingertips or toes ever change color in cold weather or when you handle cold objects?"

Ask about *swelling of feet* and *legs*, or any ulcers on lower legs, often near the ankles from peripheral vascular disease.

Hair loss over the anterior tibiae in PAD. "Dry" or brownblack ulcers from gangrene may ensue.

Only approximately 10% of affected patients have the classic symptoms of exertional calf pain relieved by rest.

Digital ischemic changes from arterial spasm cause blanching, followed by cyanosis and then rubor with cold exposure and rewarming in *Raynaud's* phenomenon or disease

Calf swelling in deep venous thrombosis; hyperpigmentation, edema, and possible cyanosis, especially when legs are dependent, in venous stasis ulcers; swelling with redness and tenderness in cellulitis

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Screening for peripheral arterial disease (PAD); the anklebrachial index
- Screening for renal artery disease
- Screening for abdominal aortic aneurysm

Screening for Peripheral Arterial Disease (PAD). PAD involves the femoral and popliteal arteries most commonly, followed by the tibial and peroneal arteries. PAD affects from 12% to 29% of community populations; despite significant association with cardiovascular and cerebrovascular disease, PAD often is underdiagnosed in office practices. Most patients with PAD have either no symptoms or a range of nonspecific leg symptoms, such as aching, cramping, numbness, or fatigue.

Screen patients for PAD risk factors, such as tobacco abuse, elevated cholesterol, diabetes, age older than 70 years, hypertension, or atherosclerotic coronary, carotid, or renal artery disease. Pursue aggressive risk factor intervention. Consider use of the ankle-brachial index (ABI), a highly accurate test for detecting stenoses of 50% or more in major vessels of the legs.

A wide range of interventions is available to reduce both onset and progression of PAD, including meticulous foot care and well-fitting shoes, tobacco cessation, treatment of hyperlipidemia, optimal control and treatment of diabetes and hypertension, use of antiplatelet agents, graded exercise, and, if needed, surgical revascularization. Patients with ABIs in the lowest category have a 20% to 25% annual risk for death.

Screening for Renal Artery Disease. The American College of Cardiology and the American Heart Association recommend diagnostic studies for renal artery disease, usually beginning with ultrasound, in patients with the following conditions: hypertension before 30 years; severe hypertension (see p. 63) after 55 years; accelerated, resistant, or malignant hypertension; new worsening of renal function or worsening after use of an angiotensin-converting enzyme inhibitor or an angiotensin-receptor blocking agent; an unexplained small kidney; or sudden unexplained pulmonary edema, especially in the setting of worsening renal function. Symptoms arise from these conditions rather than directly from atherosclerotic changes in the renal artery.

Screening for Abdominal Aortic Aneurysm (AAA). An AAA is present when the infrarenal aortic diameter exceeds 3.0 cm. Rupture and mortality rates dramatically increase for AAAs exceeding 5.5 cm in diameter. The strongest risk factor for

218 The Peripheral Vascular System

rupture is excess aortic diameter. Additional risk factors are smoking, age older than 65 years, family history, coronary artery disease, PAD, hypertension, and elevated cholesterol level. Because symptoms are rare, and screening is now shown to reduce mortality by approximately 40%, the U.S. Preventive Services Task Force recommends one-time screening by ultrasound in men between 65 and 75 years old with a history of "ever smoking," defined as more than 100 cigarettes in a lifetime.

TECHNIQUES OF EXAMINATION

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS



ADMS

Inspect for

Size and symmetry, any swelling

Venous pattern

Color and texture of skin and nails

Lymphedema, venous obstruction

Venous obstruction

Raynaud's disease

Palpate and grade the pulses:

Grading Arterial Pulses

- 3+ Bounding
- 2+ Brisk, expected (normal)
- 1+ Diminished, weaker than expected
- 0 Absent, unable to palpate

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

Radial



Bounding radial, carotid, and femoral pulses in aortic insufficiency

Lost in thromboangiitis obliterans or acute arterial occlusion

Brachial



Feel for the epitrochlear nodes.

Lymphadenopathy from local cut, infection

← LEGS

Inspect for

• Size and symmetry, any swelling in thigh or calf

Venous pattern

Color and texture of skin

Venous insufficiency, lymphedema; deep venous thrombosis

Varicose veins

Pallor, rubor, cyanosis; erythema, warmth in cellulitis, thrombophlebitis

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

• Hair distribution

Loss in arterial insufficiency

Check for pitting edema.

Dependent edema, congestive heart failure, hypoalbuminemia, nephrotic syndrome

Palpate the calves.

Tenderness in deep venous thrombosis (though tenderness often not present)

Palpate and grade the pulses:

Loss of pulses in acute arterial occlusion and arteriosclerosis obliterans

- Femoral
- Popliteal



• Dorsalis pedis



EXAMINATION TECHNIQUES

POSSIBLE FINDINGS


• Posterior tibial



Palpate the inguinal lymph nodes:

Lymphadenopathy in genital infections, lymphoma, AIDs

- Horizontal group
- Vertical group



Ask patient to stand, and reinspect the venous pattern.

Varicose veins

POSSIBLE FINDINGS

SPECIAL TECHNIQUES

EVALUATING ARTERIAL SUPPLY TO THE HAND

Feel ulnar pulse, if possible. Perform an Allen test.



1. Ask patient to make a tight fist, palm up. Occlude both radial and ulnar arteries with your thumb.



Persisting pallor of palm



2. Ask patient to open hand into a relaxed, slightly flexed position.



3. Release your pressure over one artery.



4. Palm should flush within 3 to 5 seconds. Repeat, releasing other artery.

POSTURAL COLOR **CHANGES OF CHRONIC** ARTERIAL INSUFFICIENCY

Raise both legs to 60° for about 1 minute. Then ask patient to sit up with legs dangling down. Note time required for (1) return of pinkness (normally 10 seconds) and (2) filling of veins on feet and ankles (normally about 15 seconds).

Marked pallor of feet on elevation, delayed color return and venous filling, and rubor of dependent feet suggest arterial insufficiency.

RECORDING YOUR FINDINGS

Recording the Physical Examination— The Peripheral Vascular System

"Extremities are warm and without edema. No varicosities or stasis changes. Calves are supple and nontender. No femoral or abdominal bruits. Brachial, radial, femoral, popliteal, dorsalis pedis (DP), and posterior tibial (PT) pulses are 2+ and symmetric."

OR

"Extremities are pale below the midcalf, with notable hair loss. Rubor noted when legs dependent but no edema or ulceration. Bilateral femoral bruits; no abdominal bruits heard. Brachial and radial pulses 2+; femoral, popliteal, DP, and PT pulses 1+." (Alternatively, pulses can be recorded as below.)

	Radial	Brachial	Femoral	Popliteal		Posterior Tibial
RT	2+	2+	1+	1+	1+	1+
LT	2+	2+	1+	1+	1+	1+
(Suggests atherosclerotic PAD)						

INTERPRETATION

Chronic Insufficiency of Arteries and Veins

Condition

Chronic Arterial Insufficiency



Chronic Venous Insufficiency



Characteristics

Intermittent claudication progressing to pain at rest. Decreased or absent pulses. Pale, especially on elevation; dusky red on dependency. Cool. No or mild edema. which may develop on lowering the leg to relieve pain. Thin, shiny, atrophic skin, with hair loss over foot and toes and thickened, ridged nails. Possible ulceration on toes or points of trauma on feet. Potential gangrene.

No pain to aching pain on dependency. Normal pulses, though may be hard to feel because of edema. Color normal or cyanotic on dependency; petechiae or brown pigment may develop. Often marked edema. Stasis dermatitis, possible thickening of skin, and narrowing of leg as scarring develops. Potential ulceration at sides of ankles. No gangrene.



Common Ulcers of the Feet and Ankles

Ulcer

Arterial Insufficiency



Chronic Venous Insufficiency



Neuropathic Ulcer



Characteristics

Located on toes, feet, or possible areas of trauma. No callus or excess pigment. May be atrophic. Pain often severe, unless masked by neuropathy. Possible gangrene. Decreased pulses, trophic changes, pallor of foot on elevation, dusky rubor on dependency.


Located on inner or outer ankle. Pigmented, sometimes fibrotic. Pain not severe. No gangrene. Edema, pigmentation, stasis dermatitis, and possibly cyanosis of feet on dependency.

Located on pressure points in areas with diminished sensation, as in diabetic neuropathy. Skin calloused. No pain (which may cause ulcer to go unnoticed). Usually no gangrene. Decreased sensation, absent ankle jerks.


TABLE 12-3 Using the Ankle-Brachial Index

Instructions for Measuring the Ankle-Brachial Index (ABI)*

1. Patient should rest supine in a warm room for at least 10 minutes before testing.



- Place blood pressure cuffs on both arms and ankles as illustrated, then apply ultrasound gel over brachial, dorsalis pedis, and posterior tibial arteries.
- 3. Measure systolic pressures in the arms
 - Use Doppler to locate brachial pulse
 - Inflate cuff 20 mm Hg above last audible pulse
 - Deflate cuff slowly and record pressure at which pulse becomes audible
 - Obtain 2 measures in each arm and record the average as the brachial pressure in that arm



- 4. Measure systolic pressures in ankles
 - Use Doppler to locate dorsalis pedis pulse
 - Inflate cuff 20 mm Hg above last audible pulse

(table continues next page)

Using the Ankle-Brachial Index (continued)

Instructions for Measuring the Ankle-Brachial Index (ABI)*

- Deflate cuff slowly and record pressure at which pulse becomes audible
- Obtain 2 measures in each ankle and record the average as the dorsalis pedis pressure in that leg
- Repeat above steps for posterior tibial arteries
- 5. Calculate ABI

 $Right \ ABI = \frac{highest \ right \ average \ ankle \ pressure \ (DP \ or \ PT)}{highest \ average \ arm \ pressure \ (right \ or \ left)}$


 $Left ABI = \frac{highest left average ankle pressure (DP or PT)}{highest average arm pressure (right or left)}$

Interpretation of Ankle-Brachial Index*

Ankle-Brachial Index Result >0.90 (with a range of 0.90 to 1.30)	Clinical Interpretation Normal lower extremity blood flow
<0.89 to >0.60	Mild PAD
<0.59 to >0.40	Moderate PAD
< 0.39	Severe PAD

^{(*}Source: Laine C, Goldman D, Wilson JF. In the clinic: peripheral arterial disease. Ann Int Med 146(5):ITC 3-1, 2007.)

Male Genitalia and Hernias



THE HEALTH HISTORY

Common or Concerning Symptoms

- Sexual preference and sexual response
- Penile discharge or lesions
- Scrotal pain, swelling, or lesions

230 Male Genitalia and Hernias

Explain your concern for the patient's sexual health. Pose questions in a neutral and nonjudgmental way.

- "What is your relationship status? Tell me about your sexual preference."
- "How is sexual function for you?" "Are you satisfied with your sexual life?" "What about your ability to perform sexually?"

To assess *libido*, or desire: "Have you maintained an interest in sex?"

For the *arousal* phase: "Can you achieve and maintain an erection?"

If *ejaculation* is premature or early: "About how long does intercourse last?" "Do you climax too soon?" For reduced or absent ejaculation: "Do you find that you cannot have orgasm even though you can have an erection?" "Does the problem involve the pleasurable sensation of *orgasm*, the ejaculation of seminal fluid, or both?"

Decreased interest from psychogenic causes such as depression, endocrine dysfunction, or side effects of medications

Erectile dysfunction from psychogenic causes, especially if early morning erection is preserved; also from decreased testosterone, decreased blood flow in hypogastric arterial system, impaired neural innervation

Premature ejaculation is common, especially in young men. Less common is reduced or absent ejaculation affecting middle-aged or older men. Possible causes are medications, surgery, neurologic deficits, or lack of androgen. Lack of orgasm with intact ejaculation is usually psychogenic.

To assess possible infection from sexually transmitted diseases (STDs), ask about any discharge from the penis.

Inquire about sores or growths on the penis and any pain or swelling in the scrotum.

STDs may involve other parts of the body. Ask about practices of oral and anal sex and any related sore throat, oral itching or pain, diarrhea, or rectal bleeding.

Penile discharge in gonococcal (usually yellow) and nongonococcal (clear or white) urethritis

See Table 13-1, Abnormalities of the Penis and Scrotum (p. 236), and Table 13-2, **Sexually Transmitted Diseases** of Male Genitalia (p. 237).

Rash in disseminated gonococcal infection

Important Topics for Health Promotion and Counseling

- Prevention of STDs and HIV
- Testicular self-examination

Prevention of STDs and HIV Infection. Focus on patient education about STDs and HIV, early detection of infection during history taking and physical examination, and identification and treatment of infected partners. Identify the patient's sexual orientation, the number of sexual partners in the past month, and any history of STDs. Also query use of alcohol and drugs, particularly injection drugs. Counsel patients at risk about limiting the number of partners, using condoms, and establishing regular medical care for treatment of STDs and HIV infection.

Counseling and testing for HIV are recommended for: all people at increased risk for infection with HIV, STDs, or

232 Male Genitalia and Hernias

both; men with male partners; past or present injection drug users; men and women having unprotected sex with multiple partners; sex workers; any past or present partners of people with HIV infection, bisexual practices, or injection drug use; and patients with a history of transfusion between 1978 and 1985.

Testicular Self-examination. Encourage men, especially those between 15 and 35 years of age, to perform monthly testicular self-examinations.

TECHNIQUES OF EXAMINATION

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

MALE GENITALIA

Wear gloves. The patient may be standing or supine.

THE PENIS



 Development of the penis and the skin and hair at its base

Sexual maturation, lice

Prepuce Phimosis

• Glans Balanitis, chancre, herpes,

warts, cancer

• Urethral meatus *Hypospadias,* discharge of

urethritis

Palpate

Any visible lesions Chancre, cancer

The shaft
 Urethral stricture or cancer

POSSIBLE FINDINGS

THE SCROTUM AND ITS CONTENTS

Inspect

- Contours of scrotum
- Skin of scrotum

Palpate each

- Testis, noting any
 - Lumps
 - Tenderness



- Epididymis
- Spermatic cord and adjacent areas



Hernia, hydrocele, cryptorchidism

Rashes

Testicular carcinoma

Orchitis, torsion of the spermatic cord, strangulated inguinal hernia

Epididymitis, cyst

Varicocele if multiple tortuous veins; if cystic structure, may be a hydrocele

HERNIAS

Patient is usually standing.

Inspect inguinal and femoral areas as patient strains down.

Inguinal and femoral hernias

Palpate external inguinal ring through scrotal skin and ask patient to strain down.

Indirect and direct inguinal hernias



SPECIAL TECHNIQUE

THE TESTICULAR SELF-EXAMINATION

PATIENT INSTRUCTIONS FOR THE TESTICULAR SELF-EXAMINATION

This examination is best performed after a warm bath or shower. The heat relaxes the scrotum and makes it easier to find anything unusual.

- Standing in front of a mirror, check for any swelling on the skin of the scrotum.
- Examine each testicle with both hands. Cup the index and middle fingers under the testicle and place the thumbs on top.
- Roll the testicle gently between the thumbs and fingers. One testicle may be larger than the other . . . that's normal, but be concerned about any lump or area of pain.

(continued)

PATIENT INSTRUCTIONS FOR THE TESTICULAR SELF-EXAMINATION (CONTINUED)

• Find the epididymis. This is a soft, tubelike structure at the back of the testicle that collects and carries sperm, not an abnormal lump.



 If you find any lump, don't wait. See your doctor. The lump may just be an infection, but if it is cancer, it will spread unless stopped by treatment.

(Source: Medline Plus. U.S. National Library of Medicine and National Institutes of Health. Medical Encyclopedia—Testicular self-examination. Available at: www.nlm.nih.gov/medlineplus/ency/article/003909.htm. Accessed June 8, 2007.)

RECORDING YOUR

Recording the Physical Examination— Male Genitalia and Hernias

"Circumcised male. No penile discharge or lesions. No scrotal swelling or discoloration. Testes descended bilaterally, smooth, without masses. Epididymis nontender. No inquinal or femoral hernias."

OR

"Uncircumcised male; prepuce easily retractible. No penile discharge or lesions. No scrotal swelling or discoloration. Testes descended bilaterally; right testicle smooth; 1×1 cm firm nodule on left lateral testicle. It is fixed and nontender. Epididymis nontender. No inquinal or femoral hernias."

(Suspicious for testicular carcinoma, the most common form of cancer in men between 15 and 35 years of age)

TERPRETATION

Abnormalities of the Penis and Scrotum



Hypospadias

A congenital displacement of the urethral meatus to the inferior surface of the penis. A groove extends from the actual urethral meatus to its normal location on the tip of the glans.



Peyronie's Disease

Palpable, nontender, hard plaques are found just beneath the skin, usually along the dorsum of the penis. The patient complains of crooked, painful erections.



Carcinoma of the Penis

An indurated nodule or ulcer that is usually nontender. Limited almost completely to men who are not circumcised, it may be masked by the prepuce. Any persistent penile sore is suspicious.




Scrotal Edema

Pitting edema may make the scrotal skin taut; seen in congestive heart failure or nephrotic syndrome.



Hydrocele

A nontender, fluid-filled mass within the tunica vaginalis. It transilluminates, and the examining fingers can get above the mass within the scrotum.



Scrotal Hernia

Usually an indirect inguinal hernia that comes through the external inguinal ring, so the examining fingers cannot get above it within the scrotum.

TABLE 13

Sexually Transmitted Diseases of Male Genitalia



Genital Warts (condylomata acuminata)

 Appearance: Single or multiple papules or plaques of variable shapes; may be round. acuminate (or pointed), or thin and slender.

May be raised, flat, or cauliflower-like (verrucous).

- Causative organism: Human papillomavirus (HPV), usually from subtypes 6, 11; carcinogenic subtypes rare, approximately 5-10% of all anogenital warts. Incubation: weeks to months; infected contact may have no visible warts.
- · Can arise on penis, scrotum, groin, thighs, anus; usually asymptomatic, occasionally cause itching and pain.
- May disappear without treatment.



Primary Syphilis

Appearance: Small red papule that becomes a chancre, or painless erosion up

to 2 cm in diameter. Base of chancre is clean, red, smooth, and glistening; borders are raised and indurated. Chancre heals within 3-8 weeks.

- Causative organism: Treponema pallidum, a spirochete. Incubation:
 - 9 to 90 days after exposure.
- May develop inguinal lymphadenopathy within 7 days; lymph nodes are rubbery, nontender, mobile.
- 20–30% of patients develop secondary syphilis while chancre still present (suggests co-infection with HIV).
- Distinguish from: genital herpes simplex, chancroid, granuloma inguinale from Klebsiella granulomatis (rare in U.S.; 4 variants, so difficult to identify).



Genital Herpes Simplex

- Appearance: Small scattered or grouped vesicles, 1-3 mm in size, on glans or shaft of penis. Appear as erosions if vesicular membrane breaks.
- Causative organism: Usually Herpes

simplex virus 2 (90%), a double-stranded DNA virus. Incubation: 2 to 7 days after exposure.

- Primary episode may be asymptomatic; recurrence usually less painful, of shorter duration.
- · Associated with fever, malaise, headache, arthralgias; local pain and edema, lymphadenopathy.
- Need to distinguish from genital herpes zoster (usually in older patients with dermatomal distribution); candidiasis.



Chancroid

 Appearance: Red papule or pustule initially, then forms a painful deep ulcer with ragged

non-indurated margins; contains necrotic exudate, has a friable base.

- Causative organism: Haemophilus ducreyi, an anaerobic bacillus. Incubation: 3 to 7 days after exposure.
- · Painful inguinal adenopathy; suppurative bobos in 25% of patients.
- Need to distinguish from: primary syphilis; genital herpes simplex; lymphomogranuloma venerium, granuloma inguinale from Klebsiella granulomatis (both rare in U.S.).

TABLE 13-3 AL

Abnormalities of the Testes



Cryptorchidism
Testis is atrophied and may lie in the inguinal canal or the abdomen, resulting in an unfilled scrotum.
As above, there is no palpable left testis or epididymis. Cryptorchidism markedly raises the risk for testicular cancer.



Small Testis In adults, testicular length is usually ≤3.5 cm. Small, firm testes seen in Klinefelter's *syndrome*, usually ≤2 cm. Small, soft testes suggesting atrophy seen in cirrhosis, myotonic dystrophy, use of estrogens, and hypopituitarism; may also follow orchitis.



Acute Orchitis
The testis is
acutely inflamed,
painful, tender,
and swollen. It
may be difficult
to distinguish
from the
epididymis. The
scrotum may be
reddened. Seen
in mumps and
other viral
infections; usually
unilateral.



Tumor of the Testis Usually appears as a painless nodule. Any nodule within the testis warrants investigation for malignancy.



As a testicular neoplasm grows and spreads, it may seem to replace the entire organ. The testicle characteristically feels heavier than normal.

Abnormalities of the Epididymis and Spermatic Cord



Acute Epididymitis

An acutely inflamed epididymis is tender and swollen and may be difficult to distinguish from the testis. The scrotum may be reddened and the vas deferens inflamed. It occurs chiefly in adults. Coexisting urinary tract infection or prostatitis supports the diagnosis.



Varicocele

Varicocele refers to varicose veins of the spermatic cord, usually found on the left. It feels like a soft "bag of worms" separate from the testis, and slowly collapses when the scrotum is elevated in the supine patient.



Spermatocele

and Cyst of the Epididymis A painless, movable cystic mass just above the testis suggests a spermatocele or an epididymal cyst. Both transilluminate. The former contains sperm, and the latter does not, but they are clinically indistinguishable.



Torsion of the Spermatic Cord Twisting of the testicle on its spermatic cord produces an acutely painful and swollen organ that is retracted upward in the scrotum, which becomes red and edematous. There is no associated urinary infection. It is a surgical emergency because of obstructed circulation.

Hernias in the Groin

Indirect Inquinal



Most common hernia at all ages, both sexes.

Originates above inguinal ligament and often passes into scrotum. May touch examiner's fingertip in inguinal canal.

Direct Inquinal




Less common than indirect hernia, usually occurs in men older than 40 years. Originates above inguinal ligament near external inguinal ring and rarely enters scrotum. May bulge anteriorly, touching side of examiner's finger.

Femoral



Least common hernia, more common in women than in men. Originates below inguinal ligament, more lateral than inguinal hernia. Never enters scrotum.

Female Genitalia



THE HEALTH HISTORY

Common Concerns

- Menarche, menstruation, menopause, postmenopausal bleeding
- Pregnancy
- Vulvovaginal symptoms
- Sexual preference and sexual response

242 Female Genitalia

For the *menstrual history*, ask the patient when her menstrual periods began (age at menarche).

When did her last period start, and the one before that? What is the interval between periods, from the first day of one to the first day of the next? Are menses regular or irregular? How long do they last? How heavy is the flow?

The interval from previous periods can signal possible pregnancy or menstrual irregularities.

In amenorrhea from pregnancy, common early symptoms are tenderness, tingling, or increased size of breasts; urinary frequency; nausea and vomiting; easy fatigability; and feelings that the baby is moving (usually noted at about 20 weeks).

Amenorrhea followed by heavy bleeding in threatened abortion or dysfunctional uterine bleeding

Dysmenorrhea, or painful menses, is common.

Primary dysmenorrhea from increased prostaglandin production; secondary dysmenorrhea from endometriosis, pelvic inflammatory disease, endometrial polyps

Amenorrhea is the absence of periods. Failure to begin periods is called *primary amenorrhea*, while cessation of established periods is termed *secondary amenorrhea*.

Secondary amenorrhea from low body weight from causes such as malnutrition, anorexia nervosa, stress, chronic illness, and hypothalamic–pituitary–ovarian dysfunction

Menopause, the absence of menses for 12 consecutive months, usually occurs between 48 and 55 years. Associated symptoms include hot flashes, flushing, sweating, and sleep disturbances.

For vaginal discharge and local itching, inquire about amount, color, consistency, and odor of discharge.

To assess sexual function, start with general nonjudgmental questions like "How is sex for you?" or "Are you having any problems with sex?"

Direct questions help you assess each phase of the sexual response: desire, arousal, and orgasm.

Ask also about dyspareunia, or discomfort or pain during intercourse.

For sexually transmitted diseases (STDs), identify sexual preference (male, female, or both) and the number of sexual partners in the previous month. Ask if the patient has concerns about HIV infection, desires HIV testing, or has current or past partners at risk.

Postmenopausal bleeding, or bleeding occurring 6 months after menses have stopped, raises the question of endometrial cancer, hormone replacement therapy, or uterine or cervical polyps.

See Table 14-1, Lesions of the Vulva, pp. 251-252; and Table 14-4, Vaginal Discharge, p. 255.

Superficial pain suggests local inflammation, atrophic vaginitis, or inadequate lubrication; deeper pain may result from pelvic disorders or pressure on a normal ovary.

In women, some STDs do not produce symptoms, but do pose as risk of infertility.

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Cervical cancer screening; Pap smear and HPV infection
- Options for family planning
- STDs and HIV
- Changes in menopause

Cervical Cancer Screening and HPV Infection. Observe the guidelines from the American College of Obstetricians and Gynecologists (ACOG) in 2003 for *Pap smear* screening in different age groups:

- **First screen:** 3 years after first sexual intercourse or by age 21, whichever comes first
- Women up to 30 years: Annual screening
- Women 30 years or older:
 - Screen every 2 to 3 years if three consecutive annual cervical cytology results are negative or if combined cervical cytology testing and high-risk human papillomavirus (HPV) testing are negative
 - Screen more frequently in patients with positive Pap or high-risk HPV test; HIV infection; immunosuppression; DES exposure in utero; prior history of cervical cancer
- Women with hysterectomy: Discontinue routine screening if the cervix was removed for benign reasons and there is no history of abnormal or cancerous cell growth. If the woman has such a history, screen annually, and discontinue screening if three consecutive vaginal cytology tests are negative.
- Older women: Base continued screening on clinical assessment of individual health and ability to monitor the patient.

The most important risk factor for cervical cancer is human papillomavirus (HPV) infection from HPV strains 16, 18, 6, or 11. The HPV vaccine prevents HPV 16 and 18 infection when given *before* sexual exposure.

Options for Family Planning. Counsel women, particularly adolescents, about the timing of ovulation, which occurs midway in the regular menstrual cycle. Discuss methods for contraception and their effectiveness: natural (periodic abstinence, withdrawal, lactation); barrier (condom, diaphragm, cervical cap); implantable (intrauterine device, subdermal implant); pharmacologic (spermicide, oral contraceptives, subdermal implant of levonorgestrel, estrogen/progesterone injectables and patch, vaginal ring); and surgical (tubal ligation).

STDs and HIV. For STDs and HIV, assess risk factors by taking a careful sexual history and counseling patients about spread of disease and ways to reduce high-risk practices. Test women younger than 24 years and pregnant women for chlamydia; in women at increased risk and pregnant women, test for gonorrhea, syphilis, and HIV.

Menopause. Be familiar with the psychological and physiologic changes of *menopause*. Help the patient to weigh the benefits and risks of treatment, taking into account the personal and family history of cardiovascular disease and osteoporosis.

ECHNIQUES OF EXAMINATION

TIPS FOR THE SUCCESSFUL PELVIC EXAMINATION

The Patient

- Avoids intercourse, douching, or use of vaginal suppositories for 24 to 48 hours before examination
- Empties bladder before examination
- Lies supine, with head and shoulders elevated, arms at sides or folded across chest to enhance eye contact and reduce tightening of abdominal muscles

The Examiner

- Obtains permission
- Explains each step of the examination in advance; selects chaperone
- Drapes patient from mid-abdomen to knees; depresses drape between knees to provide eye contact with patient
- Avoids unexpected or sudden movements
- Chooses a speculum that is the correct size
- Warms speculum with tap water
- Wears gloves
- Monitors comfort of the examination by watching the patient's face
- Uses excellent but gentle technique, especially when inserting the speculum

Male examiners should be accompanied by female assistants. Female examiners should be assisted whenever possible.

EXAMINATION TECHNIQUES

EXTERNAL GENITALIA

Observe pubic hair to assess sexual maturity.

Examine the external genitalia.

Labia

Normal or delayed puberty

Inflammation

Clitoris

Enlarged in masculinization

Urethral orifice

Urethral caruncle

Introitus

Imperforate hymen

Palpate for enlargement or tenderness of Bartholin's glands.

Bartholin's gland infection

Milk the urethra for discharge, if indicated. Discharge of urethritis

INTERNAL GENITALIA AND PAP SMEAR

Locate the cervix with a gloved and water-lubricated index finger.

Assess support of vaginal outlet by asking patient to strain down.

Cystocele, cystourethrocele, rectocele

Enlarge the introitus by pressing its posterior margin downward

Insert a water-lubricated speculum of suitable size, starting with speculum held obliquely.



ENTRY ANGLE



POSSIBLE FINDINGS

Rotate the speculum, open it, and inspect cervix.


Observe

- Position
- Color
- Epithelial surface

Cervix faces forward if uterus is retroverted.

Purplish in pregnancy

Squamous and columnar epithelium



- Any discharge or bleeding
- Any ulcers, nodules, or masses

Obtain specimens for cytology (Pap smears) with

- An endocervical spatula or brush (except in pregnant women), to scrape the ectocervix
- Or, if the woman is not pregnant, a cervical broom for a combined specimen (also used for Thin Prep)

Discharge from os in mucopurulent cervicitis from chlamydia or gonorrhea

Herpes, polyp, cancer

Early cancer before it is clinically evident

Inspect the vaginal mucosa as you withdraw the speculum.

POSSIBLE FINDINGS

Bluish color and deep rugae in pregnancy; vaginal cancer



Palpate, by means of a bimanual examination,

- The cervix and fornices
- The uterus
- Right and left adnexa (ovaries)

Assess strength of pelvic muscles. With your vaginal fingers clear of the cervix, ask patient to tighten her muscles around your fingers as hard and long as she can.

Pain on moving cervix in pelvic inflammatory disease


Pregnancy, myomas; soft isthmus in early pregnancy

Ovarian masses, salpingitis, tubal pregnancy

A firm squeeze that compresses your fingers, moves them up and inward, and lasts more than 3 seconds is full strength.

POSSIBLE FINDINGS

Heat Section When indicated, **perform** a rectovaginal examination (see p. 264).



SPECIAL TECHNIQUE

HERNIAS

Ask the woman to strain down, as you palpate for a bulge in

- The femoral canal
- The labia majora up to just lateral to the pubic tubercle

Femoral hernia

Indirect inquinal hernia

RECORDING YOUR FINDINGS

Recording the Physical Examination—Female Genitalia

"No inguinal adenopathy. External genitalia without erythema, lesions, or masses. Vaginal mucosa pink. Cervix parous, pink, and without discharge. Uterus anterior, midline, smooth, and not enlarged. No adnexal tenderness. Pap smear obtained. Rectovaginal wall intact. Rectal vault without masses. Stool brown and hemoccult negative."


OR

"Bilateral shotty inguinal adenopathy. External genitalia without erythema or lesions. Vaginal mucosa and cervix coated with thin, white, homogenous discharge with mild fishy odor. After swabbing cervix, no discharge visible in cervical os. Uterus midline; no adnexal masses. Rectal vault without masses. Stool brown and hemoccult negative." (Suggests bacterial vaginosis)

AIDS TO INTERPRETATION

Lesions of the Vulva

Epidermoid Cyst



A small, firm, round cystic nodule in the labia suggests an epidermoid cyst. They are sometimes yellowish in color. Look for the dark punctum marking the blocked opening of the gland.

Venereal Wart (Condyloma Acuminatum)



Warty lesions on the labia and within the vestibule suggest condylomata acuminata from infection with human papillomavirus.

Genital Herpes



Shallow ulcers on red bases Shallow, small, painful ulcers on red bases suggest a herpes infection. Initial infection may be extensive, as illustrated here. Recurrent infections are usually confined to a small local patch.

Lesions of the Vulva (continued)

Syphilitic Chancre



A firm, painless ulcer suggests the chancre of primary syphilis. Because most chancres in women develop internally, they often go undetected.

Secondary Syphilis (Condyloma Latum)



Slightly raised, round or oval flat-topped papules covered by a gray exudate suggest condylomata lata, a manifestation of secondary syphilis. They are contagious.

Carcinoma of the Vulva



An ulcerated or raised red vulvar lesion in an elderly woman may indicate vulvar carcinoma.

TABLE 14-2 **Shapes of the Cervical Os**

Normal Variations

Oval



Slitlike



Lacerations

Unilateral Transverse



Bilateral Transverse



Stellate



Abnormalities of the Cervix



Endocervical polyp. A bright red, smooth mass that protrudes from the os suggests a polyp. It bleeds easily.



Mucopurulent cervicitis. A vellowish exudate emerging from the cervical os suggests this diagnosis. Causes include Chlamydia and gonococcal infections.



Carcinoma of the cervix. An irregular, hard mass suggests cancer. Early lesions cannot be detected by physical examination alone.



Fetal exposure to diethylstilbestrol (DES). Several changes may be seen: a collar of tissue around the cervix, columnar epithelium that covers the cervix or extends to the vaginal wall (then termed vaginal adenosis), and, rarely, carcinoma of the vagina.



Note: Accurate diagnosis depends on laboratory assessment and cultures.

Trichomonas vaginitis

Discharge: Yellowish green, often profuse, may be malodorous

Other Symptoms: Itching, vaginal soreness, dyspareunia

Vulva: May be red

Vagina: May be normal or red, with red spots,

petechiae

Laboratory Assessment: Saline wet mount for

trichomonads

Candida vaginitis Discharge: White, curdy, often thick, not

malodorous

Other Symptoms: Itching, vaginal soreness, external dysuria, dyspareunia

Vulva: Often red and swollen

Vagina: Often red with white patches of

discharge

Laboratory Assessment: KOH preparation for

branching hyphae

Bacterial vaginosis **Discharge:** Gray or white, thin, homogeneous, scant, malodorous

Other Symptoms: Fishy genital odor

Vulva: Usually normal Vagina: Usually normal

Laboratory Assessment: Saline wet mount for "clue cells," "whiff test" with KOH for fishy

odor

Relaxations of the Pelvic Floor

When the pelvic floor is weakened, various structures may become displaced. These displacements are seen best when the patient strains down.



A cystocele is a bulge of the anterior wall of the upper part of the vagina, together with the urinary bladder above it.



A cystourethrocele involves both the bladder and the urethra as they bulge into the anterior vaginal wall throughout most of its extent.



A rectocele is a bulge of the posterior vaginal wall, together with a portion of the rectum.



A prolapsed uterus has descended down the vaginal canal. There are three degrees of severity: first, still within the vagina (as illustrated); second, with the cervix at the introitus; and third, with the cervix outside the introitus.

Positions of the Uterus and Uterine Myomas



An anteverted uterus lies in a forward position at roughly a right angle to the vagina. This is the most common position. Anteflexion—a forward flexion of the uterine body in relation to the cervix—often coexists.



A retroverted uterus is tilted posteriorly with its cervix facing anteriorly.




A retroflexed uterus has a posterior tilt that involves the uterine body but not the cervix. A uterus that is retroflexed or retroverted may be felt only through the rectal wall; some cannot be felt at all.



A myoma of the uterus is a very common, benign tumor that feels firm and often irregular. There may be more than one. A myoma on the posterior surface of the uterus may be mistaken for a retrodisplaced uterus; one on the anterior surface may be mistaken for an anteverted uterus.

The Anus, Rectum, and Prostate



THE HEALTH HISTORY

Common or Concerning Symptoms

- Change in bowel habits
- Blood in the stool
- Pain with defecation; rectal bleeding or tenderness
- Anal warts or fissures
- Weak stream of urine
- Burning with urination

260 The Anus, Rectum, and Prostate

Ask about any change in bowel habits or problems with diarrhea or constipation. Is there any blood in the stool, or dark tarry stools?

Any pain with defecation, or rectal bleeding or tenderness?

Any anal warts or fissures?

In men, is there difficulty starting the urine stream or holding back urine? Is the flow weak? What about frequent urination, especially at night? Or pain or burning when passing urine? Any blood in the urine or semen or pain with ejaculation? Is there frequent pain or stiffness in the lower back, hips, or upper thighs?

Pencil-like stool or blood in stool in colon cancer; dark tarry stools in gastrointestinal bleeding

Hemorrhoids; proctitis from STDs

Human papillomavirus (HPV), condylomata lata in secondary syphilis; fissures in proctitis, Crohn's disease

These symptoms suggest urethral obstruction as in benign prostatic hyperplasia (BPH) or prostate cancer, especially in men older than 70 years. The American Urological Association (AUA) Symptom Index helps quantify BPH severity (see Table 15-1, p. 267).

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Screening for prostate cancer
- Screening for polyps and colorectal cancer
- Counseling for STDs

Prostate Cancer. Prostate cancer is the leading cancer diagnosed in men in the United States and the third leading cause of death in men. Risk factors are age, family history of prostate cancer, and African-American ethnicity.

Screening methods such as the digital rectal examination (DRE) and the prostate-specific antigen (PSA) test are not highly accurate, which complicates decisions about screening of patients without symptoms. The DRE reaches only the posterior and lateral surfaces of the prostate, missing 25% to 35% of tumors in other areas. Sensitivity of the DRE for prostate cancer is low, ranging from 20% to 68%, and the rate of false positives is high. Recommendations about an annual DRE vary.

The benefits of PSA testing are also unclear. The PSA can be elevated in benign conditions like hyperplasia and prostatitis, and its detection rate for prostate cancer is low, about 28% to 35% in asymptomatic men. Several groups recommend annual combined screening with PSA and DRE for men older than 50 years and for African-Americans and men older than 40 years with a positive family history. Some experts now recommend baseline PSA testing at age 40, reducing the threshold for biopsy from 4.0 ng/ml to 2.5 ng/ml, and tracking PSA velocity, or the rate of rise in 1 year.

For men with symptoms of prostate disorders, the clinician's role is more straightforward. Men with incomplete emptying of the bladder, urinary frequency or urgency, weak or intermittent stream or straining to initiate flow, hematuria, nocturia, or even bony pains in the pelvis should be encouraged to seek evaluation and treatment early.

Colorectal Cancer (CRC). Screening recommendations have recently been revised to promote more aggressive surveillance by clinicians:

- Clinicians should first identify whether patients are at average or increased risk for CRC, ideally by approximately age 20 years, but earlier if the patient has a family history of familial adenomatous polyposis.
- Average-risk patients 50 years or older should be offered a range of screening options to increase compliance: annual fecal occult blood testing (FOBT); flexible sigmoidoscopy

262 The Anus, Rectum, and Prostate

every 5 years, either alone or combined with annual FOBT; double-contrast barium enema every 5 years; or colonoscopy every 10 years.

 People at increased risk should undergo colonoscopy at intervals ranging from 3 to 5 years.

Clinicians should also use the 6-sample fecal occult blood test (FOBT). Avoid single-sample FOBT and DRE, which have inadequate detection rates.

Counseling for STDs. Anal intercourse increases risk for HIV and STDs. Promote abstinence, use of condoms, and good hygiene.

TECHNIQUES OF EXAMINATION

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

→ Wear gloves.

MALE

Position the patient on his side, or standing leaning forward over the examining table and hips flexed.

POSSIBLE FINDINGS

Inspect the

- Sacrococcygeal area
- Perianal area

Palpate the anal canal and rectum with a lubricated and gloved finger. Feel the:

- Walls of the rectum
- Prostate gland, as shown below, including median sulcus



Hemorrhoids, warts, herpes, chancre, cancer, fissures from proctitis or Crohn's disease

Lax sphincter tone in some neurologic disorders; tightness in proctitis

Cancer of the rectum, polyps



POSSIBLE FINDINGS

Try to **feel** above the prostate for irregularities or tenderness, if indicated.

Benign hyperplasia, cancer, tenderness in acute prostatitis



→ /M FEMALE

The patient is usually in the lithotomy position or lying on her side.

Inspect the anus.

Palpate the anal canal and rectum.

Rectal shelf of peritoneal metastases; tenderness of inflammation

Hemorrhoids

Rectal cancer, normal uterine cervix or tampon (felt through the rectal wall)

RECORDING YOUR FINDINGS

Recording the Physical Examination— The Anus, Rectum, and Prostate

"No perirectal lesions or fissures. External sphincter tone intact. Rectal vault without masses. Prostate smooth and nontender with palpable median sulcus. (Or in a female, uterine cervix nontender.) Stool brown and hemoccult negative."

OR

"Perirectal area inflamed; no ulcerations, warts, or discharge. Cannot examine external sphincter, rectal vault, or prostate because of spasm of external sphincter and marked inflammation and tenderness of anal canal." (Raises concern of proctitis from infectious cause)

OR

"No perirectal lesions or fissures. External sphincter tone intact. Rectal vault without masses. Left lateral prostate lobe with 1×1 cm firm hard nodule; right lateral lobe smooth; medial sulcus is obscured. Stool brown and hemoccult negative." (Raises concern of prostate cancer)

TABLE 15-1

BPH Symptom Score Index: American Urological Association (AUA)

Score or ask the patient to score each of the questions below. Higher scores (maximum 35) indicate more severe symptoms; scores ≤7 are considered mild and generally do not warrant treatment.

Less
Than 1
Not at Time
All in 5

PART A

- Over the past month, how often have you had a sensation of not emptying your bladder completely after you finished urinating?
- 2. Over the past month, how often have you had to urinate again less than two hours after you finished urinating?
- 3. Over the past month, how often have you stopped and started again several times when you urinated?
- 4. Over the past month, how often have you found it difficult to postpone urination?
- 5. Over the past month, how often have you had a weak urinary stream?
- 6. Over the past month, how often have you had to push or strain to begin urination?

PART B None 1 Time

7. Over the past month, how many times did you most typically get up to urinate from the time you went to bed at night until the time you got up in the morning?

⁽Adapted from: Madsen FA, Bruskewitz RC. Clinical manifestations of benign prostatic hyperplasia. Urol Clin North AM 1995:22:291–298.)

Less				
Than	About	More		
Half the	Half the	Than Half	Almost	Total Points for
Time	Time	the Time	Always	Each Row

Points for 2 Times 3 Times 4 Times 5 Times Part B

TOTAL PARTS A and B (maximum 35)

TABLE 15-2 Abnormalities on Rectal Examination

External Hemorrhoids (Thrombosed). Dilated hemorrhoidal veins that originate below the pectinate line, covered with skin; a tender, swollen, bluish ovoid mass is visible at the anal margin.



Polyps of the Rectum. A soft mass that may or may not be on a stalk; may not be palpable



Benign Prostatic Hyperplasia. An enlarged, nontender, smooth, firm but slightly elastic prostate gland; can cause symptoms without palpable enlargement



Abnormalities on Rectal TABLE 15 **Examination** (continued)

Acute Prostatitis. A prostate that is very tender, swollen, and firm because of acute infection



Cancer of the Prostate. A hard area in the prostate that may or may not feel nodular



Cancer of the Rectum. Firm, nodular, rolled edge of an ulcerated cancer



16

The Musculoskeletal System

FUNDAMENTALS FOR ASSESSING JOINTS

Assessing joints requires knowledge of their structure and function. Learn the surface landmarks and underlying anatomy of each major joint. Be familiar with the following terms:


- Articular structures—include the joint capsule and articular cartilage, synovium and synovial fluid, intraarticular ligaments, and juxta-articular bone
- Extra-articular structures—include periarticular ligaments, tendons, bursae, muscle, fascia, bone, nerve, and overlying skin
 - Ligaments—the ropelike bundles of collagen fibrils that connect bone to bone
 - Tendons—the collagen fibers that connect muscle to bone
 - Bursae—the pouches of synovial fluid that cushion the movement of tendons and muscles over bone or other joint structures

Review the three primary types of joint articulation—synovial, cartilaginous, and fibrous—and the varying degrees of movement each type allows.

Joints

Synovial Joints


- Freely movable
- Separated by articular cartilage and a synovial cavity
- Lubricated by synovial fluid
- Surrounded by a joint capsule
- Example: knee, shoulder



Cartilaginous Joints

- Slightly movable
- Contain fibrocartilaginous discs that separate the bony surfaces
- Have a central nucleus pulposus of discs that cushions bony contact
- Example: vertebral bodies

SYNOVIAL



CARTILAGINOUS

Fibrous Joints

- No appreciable movement
- Consist of fibrous tissue or cartilage
- Lack a joint cavity
- Example: skull sutures



FIBROUS

Review the types of synovial joints and their associated features as well.

Note that joint structure determines joint function.

Synovial Joints				
Type of Joint	Articular Shape	Movement	Example	
Spheroidal (ball and socket)	Convex surface in concave cavity	Wide-ranging flexion, extension, abduction, adduction, rotation, circumduction	Shoulder, hip	
Hinge	Flat, planar	Motion in one plane; flexion, extension	Interphalangeal joints of hand and foot; elbow	
Condylar	Convex or concave	Movement of two articu- lating surfaces, not dissociable	Knee; temporo- mandibu- lar joint	

HE HEALTH HISTORY

Common or Concerning Symptoms

- Low back pain
- Neck pain
- Monoarticular or polyarticular joint pain
- Inflammatory or infectious joint pain
- Joint pain with systemic features such as fever, chills, rash, anorexia, weight loss, and weakness
- Joint pain with symptoms from other organ systems

Assess the seven features of any joint pain.

TIPS FOR ASSESSING JOINT PAIN

• Ask the patient to point to the pain. This may save considerable time, because the patient's verbal description is often imprecise. (continued)

TIPS FOR ASSESSING JOINT PAIN (CONTINUED)

- Clarify and record the mechanism of injury, particularly if there is a history of trauma.
- Determine whether the pain is localized or diffuse, acute or chronic, inflammatory or noninflammatory.

Low Back Pain. Ask "Any pains in your back?" Backache is the second most common reason for office visits. Ask if the pain is in the midline over the vertebrae, or off midline. If the pain radiates into the legs, ask about any associated numbness, tingling, or weakness.

Check for bladder or bowel dysfunction.

Neck Pain. Ask about location, radiation into the arms, arm or leg weakness, bladder or bowel dysfunction.

Joint Pain. Proceed with "Do you have any pain in your joints?"

Ask the patient to point to the pain. If localized and involving only one joint, it is monoarticular.

If polyarticular, does it migrate from joint to joint, or steadily spread from one joint to

See Table 16-1, Low Back Pain, pp. 301-302. Causes of midline back pain include vertebral collapse, disc herniation, epidural abscess, spinal cord compression, or spinal cord metastases. Pain off the midline in sacroiliitis. trochanteric bursitis, sciatica, or hip arthritis

Present in cauda equine syndrome from S2-S4 tumor or disc herniation

Often from C7 or C6 spinal nerve compression from foraminal impingement

Consider trauma, monoarticular arthritis, tendonitis, or bursitis. Hip pain near the greater trochanter suggests trochanteric bursitis.

Migratory pattern in rheumatic fever or gonococcal arthritis; progressive and symmetric in rheumatoid arthritis

multiple joint involvement? Is the involvement symmetric?

Ask if pain is extra-articular (bones, muscles, and tissues around the joint, such as the tendons, bursae, or even overlying skin). Are there generalized "aches and pains" (myalgia if in muscles, arthralgia if in joints with no evidence of arthritis)?

Bursitis if inflammation of bursae. tendonitis if in tendons. and tenosynovitis if in tendon sheaths; also sprains from stretching or tearing of ligaments

Assess the timing, quality, and severity of joint symptoms. If from trauma, what was the mechanism of injury or the series of events that caused the joint pain? Further, what aggravates or relieves the pain? What are the effects of exercise, rest, and treatment?

Severe pain of rapid onset in a red, swollen joint in acute septic arthritis or gout

Is the problem inflammatory or noninflammatory? Is there tenderness, warmth, or redness?

Fever, chills, warmth, redness in septic arthritis; also consider gout or rheumatic fever

Is the pain articular in origin, with swelling, stiffness, or decreased range of motion? Pain, swelling, loss of active and passive motion, "locking," deformity in articular joint pain; loss of active but not passive motion, tenderness outside the joint, no deformity in nonarticular pain

Assess any limitations of motion

Transient stiffness after limited activity in degenerative arthritis; prolonged stiffness in *rheumatoid* arthritis, fibromyalqia, polymyalgia rheumatica

276 The Musculoskeletal System

Ask about any *systemic* symptoms such as fever, chills, rash, anorexia, weight loss, and weakness.

Common in rheumatoid arthritis, systemic lupus erythematosus, polymyalgia rheumatica, and other inflammatory arthritides. High fever and chills suggest an infectious cause.

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Nutrition, exercise, and weight
- Low back: lifting and biomechanics
- Preventing falls
- Osteoporosis: screening and prevention

Nutrition and Exercise. Advise patients that a healthy lifestyle conveys direct benefits to the skeleton. Good nutrition supplies calcium for bone mineralization and bone density. Exercise appears to maintain and possibly supports bone mass, and improves outlook and stress management. Appropriate weight reduces excess mechanical wear on weight-bearing joints such as hips and knees.

Low Back. The low back is especially vulnerable, most notably at L5–S1, where the sacral vertebrae make a sharp posterior angle. Approximately 60% to 80% of the population experiences low back pain at least once. Exercises to strengthen the low back, especially in flexion and extension, and general fitness exercises appear equally effective. Education on lifting strategies, posture, and the biomechanics of injury is prudent for nurses, heavy-machinery operators, and construction workers.

Preventing Falls. Falls are the leading cause of nonfatal injuries and account for a dramatic rise in death rates after 65 years of age. Risk factors include unstable gait, imbalanced posture, reduced strength, cognitive loss and dementia, deficits in vision and proprioception, and osteoporosis. Urge patients to correct poor lighting, dark or steep stairs, chairs at awkward heights, slippery or irregular surfaces, and ill-fitting shoes. Scrutinize any medications affecting balance, especially benzodiazepines, vasodilators, and diuretics.

Osteoporosis. Osteoporosis is a major public health threat for postmenopausal women and some men. The World Health Organization defines osteoporosis as bone density equal to or greater than 2.5 standard deviations below the mean for young white adult women. A 10% drop in bone density, equivalent to one standard deviation, is associated with a 20% increase in risk of fracture.

The U.S. Preventive Services Task Force recommends routine bone density screening for women 65 years or older and for the risk factors below.

RISK FACTORS FOR OSTEOPOROSIS

- Postmenopausal status in white women
- Age older than 50 years
- Weight less than 70 kg
- Family history of fracture in a first-degree relative
- History of fracture
- Higher intakes of alcohol
- Women with delayed menarche or early menopause
- Current smokers
- Low levels of 25-hydroxyvitamin D
- Use of corticosteroids for more than 2 months
- Inflammatory disorders of the musculoskeletal, pulmonary, or gastrointestinal systems, including celiac sprue, chronic renal disease, organ transplantation, hypogonadism, anorexia nervosa

(Sources: NIH Consensus Development Panel on Osteoporosis Prevention, Diagnosis, and Therapy. Osteoporosis prevention, diagnosis, and therapy. JAMA 285(6):785-795, 2001; Raisz LG. Screening for osteoporosis. N Engl J Med 353(2):164-171, 2005.)

Several agents inhibit bone resorption: calcium, vitamin D, and antiresorptive agents such as bisphosphonates, selective estrogen-receptor modulators (SERMs), and calcitonin. Learn the therapeutic uses of the agents listed above and of exercise. The U.S. Preventive Services Task Force now recommends against the routine use of estrogen and progestin for the prevention of chronic conditions in postmenopausal women. Despite public interest, the natural estrogens, including the plant-derived phytoestrogens, have not been shown to reduce risk of fracture in humans.

ECHNIQUES OF EXAMINATION



GENERAL APPROACH

Inspect the joints and surrounding tissues as you examine the various body parts.

Identify joints with changes in structure and function, carefully assessing for:

- Symmetry of involvement—one or both sides of the body; one joint or several
- Deformity or malalignment of bones
- Changes in surrounding soft tissue—skin changes, subcutaneous nodules, muscle atrophy, crepitus
- Limitations in range of motion, ligamentous laxity
- Changes in muscle strength

Note signs of inflammation and arthritis: swelling, warmth, tenderness, redness.

EXAMINATION TECHNIQUES

TEMPOROMANDIBULAR JOINT (TMJ)

Inspect the TMJ for swelling or redness.

Palpate the TMJ as the patient opens and closes the mouth.

Palpate the muscles of mastication: the *masseters*, temporal muscles, and pterygoid muscles.



SHOULDERS

Inspect the contour of the shoulders and shoulder girdles from front and back.

PALPATE

- The clavicle from the sternoclavicular joint to the acromioclavicular joint
- The subacromial and subdeltoid bursae after lifting arm posteriorly



Muscle atrophy; anterior or posterior dislocation of humeral head; scoliosis if shoulder heights asymmetric

"Step-offs" if fracture from trauma

Subacromial or subdeltoid bursitis

Assess range of motion.

- Flexion—"Raise your arm in front of you and overhead "
- Extension—"Move your arms behind you."
- Abduction—"Raise your arms out to the side and overhead."

Intact glenohumeral motion if patient raises arms to shoulder level, palms facing down

Intact scapulothoracic motion if patient raises arms an additional 60 degrees, palms facing up

POSSIBLE FINDINGS

- Adduction—"Cross your arm in front of your body, keeping the arm straight."
- External and internal rotation



TESTS ABDUCTION AND EXTERNAL ROTATION

Perform maneuvers to assess rotator cuff—supraspinatus, infraspinatus, teres minor, and subscapularis muscles and tendon insertions.

• "Empty can test" for supraspinatus strength





TESTS ADDUCTION AND INTERNAL ROTATION

Weakness in rotator cuff tear



EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

• Infraspinatus strength




Weakness in rotator cuff tear or bicipital tendonitis

• Forearm supination



Pain in rotator cuff tear

• "Drop arm" test



If patient cannot hold arm fully abducted at shoulder level, possible rotator cuff tear

ELBOWS

Inspect and palpate

- Olecranon process
- Medial and lateral epicondyles
- Extensor surface of the ulna
- Grooves between the epicondyles and the olecranon

Ask patient to


- Flex and extend elbows
- Turn palms up and down (supination and pronation)

Olecranon bursitis; posterior dislocation from direct trauma or supracondylar fracture

Tenderness distal to epicondyle in epicondylitis (medial → "tennis elbow"; lateral → "pitcher's elbow")

Rheumatoid nodules

Tender in arthritis



POSSIBLE FINDINGS

WRISTS AND HANDS

Inspect

- Movement of the wrist (flexion, extension, ulnar and medial deviation), hands, and fingers
- Guarded movement in injury
- Contours of wrists, hands, and fingers

Deformities in rheumatoid and degenerative arthritis; swelling in arthritis, ganglia; impaired alignment of fingers in flexor tendon damage; flexion contractures in Dupuytren's contractures

Contours of palms

Thenar atrophy in median nerve compression (carpal tunnel syndrome); hypothenar atrophy in ulnar nerve compression

Palpate

Wrist joints



Swelling and tenderness in rheumatoid arthritis, gonococcal infection of joint or extensor tendon sheaths

Distal radius and ulna

Tenderness over ulnar styloid in Colles' fracture

EXAMINATION TECHNIQUES

 "Anatomic snuffbox," the hollow space distal to the radial styloid bone; thumb extensor and abductor tendons.



POSSIBLE FINDINGS

Tenderness suggests scaphoid fracture. Tenderness over extensor and abductor tendons in DeQuervain's tenosynovitis.

• Metacarpophalangeal joint



Swelling in rheumatoid arthritis

 Proximal and distal interphalangeal joint



Proximal nodules in rheumatoid arthritis (Bouchard's nodes), distal nodules in osteoarthritis (Heberden's nodes)

POSSIBLE FINDINGS

Assess range of motion.

- Wrists: Flexion, extension, adduction (radial deviation), abduction (lateral deviation)
- Fingers: Flexions, extension, abduction/ adduction (spread fingers apart and back)
- Thumbs

Arthritis, tenosynovitis

Trigger finger, Dupuytren's contracture



FLEXION



ABDUCTION AND ADDUCTION

Perform selected maneuvers.

• Hand grip strength



EXTENSION




OPPOSITION

Decreased grip strength if weakness of finger flexors or intrinsic hand muscles



• Thumb movement



Pain if DeQuervain's tenosynovitis

- Carpal tunnel
 - Thumb adduction



Weakness of abductor pollicis longus specific to median nerve

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

• Tinel's sign: Tap lightly over median nerve at volar wrist

Aching, tingling, and numbness in 2nd, 3rd, and 4th fingers



• Phalen's sign: Patient flexes wrists for 60 seconds

Aching, tingling, and numbness in 2nd, 3rd, and 4th volar fingers




SPINE

Inspect spine from the side and back, noting any abnormal curvatures.

Look for asymmetric heights of shoulders, iliac crests, or buttocks.

Kyphosis, scoliosis, lordosis, gibbus, list curvatures

Pelvic tilt



Identify and palpate

- Spinous processes of each vertebra
- Sacroiliac joints
- Paravertebral muscles, if painful
- Sciatic nerve (midway between greater trochanter and ischial tuberosity)

Tender if trauma, infection

"Step-offs" in spondylolisthesis, fracture

Sacroiliitis

Paravertebral muscle spasm in abnormal posture, degenerative and inflammatory muscle disorders

Herniated disc or nerve root compression



Test the range of motion in the neck and spine in: flexion, extension, rotation, and lateral bending.


Decreased mobility in arthritis

HIPS

Inspect gait for

• Stance (see below) and swing (foot moves forward, does not bear weight)

Most problems arise during the weight-bearing stance phase.



PHASES OF GAIT: STANCE (Right Leg) AND SWING (Left Leg)

• Width of base (usually 2 to 4 inches from heel to heel), shift of pelvis, flexion of knee

Cerebellar disease or foot problems if wide base: impaired shift of pelvis in arthritis, hip dislocation, abductor weakness; disrupted gait if poor knee flexion

Palpate

- Along the inguinal ligament
- The iliopectineal bursa, lateral to the femoral pulse
- The trochanteric bursa, on the greater trochanter of the femur

Bulges in inquinal hernia, aneurysm

Tender in synovitis, bursitis, iliopsoas abscess

Focal tenderness in trochanteric bursitis, often described by patients as "low back pain"

EXAMINATION TECHNIQUES

• The *ischiogluteal bursa*, superficial to the ischial tuberosity



TROCHANTERIC AND ISCHIOGLUTEAL BURSA

Check range of motion, including

 Flexion—"Bend your knee and pull it against your abdomen."



Tender in bursitis ("weaver's bottom")

Flexion of opposite leg suggests deformity of that hip.



Extension



Abduction

Painful in iliopsoas abscess

Restricted in hip arthritis

POSSIBLE FINDINGS

- Adduction
- Internal and external rotation



Restricted in hip arthritis

KNEES

Review the structures of the knee.



Inspect

• Gait for knee extension at heel strike, flexion during all other phases of swing and stance

Stumbling or pushing knee into extension in quadriceps weakness

• Push patella distally and ask patient to tighten knee against table.

examination techniques	POSSIBLE FINDINGS
• Alignment of knees	Bowlegs, knock-knees; flexion contractures in limb paralysis
 Contours of knees, including any atrophy of the quadriceps muscles 	Quadriceps atrophy with patellofemoral disorder
Inspect and palpate:	
The tibiofemoral joint—with knees flexed	
• Joint line—place thumbs on either side of the patellar tendon.	Irregular, bony ridges in osteoarthritis.
• Medial and lateral meniscus	Tenderness if meniscus tear
Medial and lateral collateral ligaments	Tenderness if MCL tear (LCL injuries less common)
The patellofemoral compartment	
• Patella	Swelling over the patella in prepatellar bursitis ("housemaid's knee")
• Palpate the patellar tendon and ask patient to extend the leg.	Tenderness or inability to extend the leg in partial or complete tear of the patellar tendon
• Press the patella against the underlying femur.	Pain, crepitus, and a history of knee pain in <i>patellofemoral</i> disorder

Pain during contraction of quadriceps in *chondromalacia*

Also:

- Suprapatellar pouch
- Infrapatellar spaces (hollow areas adjacent to patella)
- Medial tibial condyle
- Popliteal surface

Assess any effusions.

• Bulge Sign (minor effusions): Compress the suprapatellar pouch, stroke downward on medial surface, apply pressure to force fluid to lateral surface, and then tap knee behind lateral margin of patella.


Swelling in synovitis and arthritis

Swelling in arthritis

Swelling in pes anserine bursitis

Popliteal or Baker's cyst

A fluid wave returning to the medial surface after a lateral tap confirms an effusion—a positive "bulge sign."



EXAMINATION TECHNIQUES

Balloon Sign (major effusions): Compress suprapatellar pouch with one hand; with thumb and finger of other hand, feel for fluid entering the

spaces next to the patella.

POSSIBLE FINDINGS

A palpable fluid wave is a positive sign.



• Lateral collateral ligament: With knee slightly flexed, push laterally along medial surface of knee with one hand and pull medially at the ankle with the other hand (an adduction or varus stress).

Pain or a gap in the lateral joint line points to a partial or complete LCL tear.




• Anterior cruciate ligament (ACL): (1) With knee flexed, place thumbs on medial and lateral joint line and place fingers on hamstring insertions. Pull tibia forward, observe if tibia slides forward "like a drawer." Compare to opposite knee.

Forward slide of proximal tibia is a positive *anterior drawer sign* in ACL laxity or tear.



POSSIBLE FINDINGS

(2) Lachman test: Grasp the distal femur with one hand and the proximal tibia with the other (place the thumb on the joint line). Move the femur forward and the tibia back. Significant forward excursion of tibia in ACL tear



• Ballotte the patella (major effusion): Push the patella sharply against the femur; watch for fluid returning to the suprapatellar space.


Visible wave is a positive sign.



If painful, use maneuvers to assess ligaments.

- Medial meniscus and lateral meniscus—McMurray test: With the patient supine, grasp the heel and flex the knee. Cup your other hand over the knee joint with fingers and thumb along the medial and lateral joint line. From the heel, rotate the lower leg internally and externally. Then push on the lateral side to apply a valgus stress on the medial side of the joint and rotate leg externally and slowly extend it.
- Medial collateral ligament:
 With knee slightly flexed,
 push medially against
 lateral surface of knee with
 one hand and pull laterally
 at the ankle with the
 other hand (abduction
 or valgus stress).
- Posterior cruciate ligament (PCL): Position patient and hands as in the ACL test. Push the tibia posteriorly and observe for posterior movement, like a drawer sliding posteriorly.


Click or pop along the medial joint with valgus stress, external rotation, and leg extension in tear of posterior medial meniscus



Pain or a gap in the medial joint line points to a partial or complete MCL tear.



Isolated PCL tears are rare.



POSSIBLE FINDINGS

ANKLES AND FEET

Inspect ankles and feet.

Palpate

- Ankle joint
- Ankle ligaments: medialdeltoid; lateral-anterior and posterior talofibular, calcaneofibular
- Achilles tendon
- Compress the metatarsophalangeal joints; then palpate each joint between the thumb and forefinger.

Hallux valgus, corns, calluses


Tender joint in arthritis

Tenderness in sprain: lateral ligaments weaker, inversion injuries (ankle bows outward) more common

Rheumatoid nodules, tenderness in tendonitis

Tenderness in arthritis and other conditions





• Dorsiflex and plantar flex the ankle (tibiotalar joint).



Arthritic joint often painful when moved in any direction; sprain, when injured ligament is stretched

EXAMINATION TECHNIQUES

POSSIBLE FINDINGS

• Stabilize the ankle and invert and evert the heel (subtalar or talocalcaneal joint).

Ankle sprain



2

INVERSION

EVERSION

• Stabilize the heel and invert and evert the forefoot (*transverse tarsal joints*).

Trauma, arthritis





INVERSION

EVERSION

 Flex toes at metatarsophalangeal joints.


SPECIAL TECHNIQUES

Measuring Leg Length. Patient's legs should be aligned symmetrically. With a tape, measure distance from anterior superior iliac spine to medial malleolus. Tape should cross knee medially.

Unequal leg length may be the cause of scoliosis.

/ Measuring Range of Motion. To measure range of motion precisely, a simple pocket goniometer is needed. Estimates may be made visually. Movement in the elbow at the right is limited to range indicated by red lines.

A flexion deformity of 45° and further flexion to 90° $(45^{\circ} \rightarrow 90^{\circ})$



RECORDING YOUR FINDINGS

Recording the Physical Examination— The Musculoskeletal System

"Good range of motion in all joints. No evidence of swelling or deformity."

OR

"Good range of motion in all joints. Hand with degenerative changes of Heberden's nodes at the distal interphalangeal joints, Bouchard's nodes at proximal interphalangeal joints. Mild pain with flexion, extension, and rotation of both hips. Good range of motion in the knees, with moderate crepitus; no effusion but boggy synovium and osteophytes along the tibiofemoral joint line bilaterally. Both feet with hallux valgus at the first metatarsophalangeal joints." (Suggests osteoarthritis)

AIDS TO INTERPRETATION

TABLE 16-

Low Back Pain

Patterns

Mechanical Low Back Pain

Aching pain in the lumbosacral area; may radiate into lower leg, along L5 or S1 dermatomes. Usually acute, work-related, in age group 30 to 50 years; no underlying pathology

Sciatica (Radicular Low Back Pain)

Usually from disc herniation; more rarely from nerve root compression, primary or metastatic tumor

Lumbar Spinal Stenosis

Pseudoclaudication pain in the back or legs that improves with rest, forward lumbar flexion. Pain vague but usually bilateral, with paresthesias in one or both legs; usually from arthritic narrowing of spinal canal

Physical Signs

Paraspinal muscle or facet tenderness, muscle spasm or pain with back movement, loss of normal lumbar lordosis but no motor or sensory loss or reflex abnormalities. In osteoporosis, check for thoracic kyphosis, percussion tenderness over a spinous process, or fractures in the thoracic spine or hip.

Disc herniation most likely if calf wasting, weak ankle dorsiflexion, absent ankle jerk, positive crossed straight-leg raise (pain in affected leg when healthy leg tested); negative straight-leg raise makes diagnosis highly unlikely.

Posture may be flexed forward with lowerextremity weakness and hyporeflexia; straight-leg raise usually negative

TABLE 16-1

Low Back Pain (continued)

Patterns

Chronic Back Stiffness

Consider ankylosing spondylitis in inflammatory polyarthritis, most common in men <40 years. Diffuse idiopathic skeletal hyperostosis (DISH) affects men more than women, usually aged >50 years.

Nocturnal Back Pain Unrelieved by Rest

Consider metastatic malignancy to the spine from cancer of the prostate, breast, lung, thyroid, and kidney, and multiple myeloma.

Pain Referred from the Abdomen or Pelvis

Usually a deep, aching pain, the level of which varies with the source (1% of low back pain)

Physical Signs

Loss of the normal lumbar lordosis, muscle spasm, limited anterior and lateral flexion; improves with exercise. Lateral immobility of the spine, especially thoracic segment

Findings vary with the source. Local vertebral tenderness may be present.

Spinal movements are not painful and range of motion is not affected. Look for signs of the primary disorder, such as peptic ulcer, pancreatitis, dissecting aortic aneurysm.

Pains in the Neck

Patterns

Physical Signs

Mechanical Neck Pain

Aching pain in the cervical paraspinal muscles and ligaments with associated muscle spasm, with stiffness and tightness in the upper back and shoulder, lasting up to 6 weeks. No associated radiation, paresthesias, or weakness. Headache may be present.

Mechanical Neck Pain—Whiplash

Also mechanical neck pain with aching paracervical pain and stiffness, often beginning the day after injury. Occipital headache, dizziness, malaise, and fatigue may be present. Chronic whiplash syndrome if symptoms last more than 6 months, present in 20%-40% of injuries.

Cervical Radiculopathy from nerve root compression

Sharp burning or tingling pain in the neck and one arm, with associated paresthesias and weakness. Sensory symptoms often in myotomal pattern, deep in muscle, rather than dermatomal pattern.

Local muscle tenderness, pain on movement. No neurological deficits. Possible trigger points in fibromyalgia. Torticollis if prolonged abnormal neck posture and muscle spasm.

Localized paracervical tenderness, decreased neck range of motion, perceived weakness of the upper extremities. Causes of cervical cord compression such as fracture, herniation, head injury, or altered consciousness are excluded.

C7 nerve root affected most often (\sim 45%–60%), with weakness in triceps and finger flexors and extensors. C6 nerve root involvement also common, with weakness in biceps, brachioradialis, wrist extensors.

TABLE 16-2 Pains in the Neck (continued)

Patterns

Physical Signs

Cervical Myelopathy—from cervical cord compression

Neck pain with bilateral weakness and paresthesias in both upper and lower extremities, often with urinary frequency. Hand clumsiness, palmar paresthesias, and gait changes may be subtle. Neck flexion often exacerbates symptoms.

Hyperreflexia; clonus at the wrist, knee, or ankle; extensor plantar reflexes (positive Babinski signs); and gait disturbances. May also see *Lhermitte's sign:* neck flexion with resulting sensation of electrical shock radiating down the spine. Confirmation of cervical myelopathy warrants neck immobilization and neurosurgical evaluation.

Patterns of Pain in and **Around the Joints**

	Rheumatoid Arthritis	Osteoarthritis (Degenerative Joint Disease, or DJD)
Process	Chronic inflammation of synovial membranes with secondary erosion of adjacent cartilage and bone, and damage to ligaments and tendons	Degeneration and progressive loss of cartilage within the joints, damage to underlying bone, and formation of new bone at the margins of the cartilage
Common Locations	Hands (proximal interphalangeal and metacarpophalangeal joints), feet (metatarsophalangeal joints), wrists, knees, elbows, ankles	Knees, hips, hands (distal, sometimes proximal interphalangeal joints), cervical and lumbar spine, and wrists (first carpometacarpal joint); also joints previously injured or diseased
Pattern of Spread	Symmetrically additive: progresses to other joints while persisting in the initial ones	Additive; however, sometimes only one joint is involved
Onset	Usually insidious	Usually insidious
Progression and Duration	Often chronic, with remissions and exacerbations	Slowly progressive, with temporary exacerbations after periods of overuse

_____ Patterns of Pain in and

TABLE 16-3 Around the Joints (continued)		
	Rheumatoid Arthritis	Osteoarthritis (Degenerative Joint Disease, or DJD)
Associated Symptoms	Frequent swelling of synovial tissue in joints or tendon sheaths; also sub- cutaneous nodules Tender, often warm, but seldom red	Small effusions in the joints may be present, especially in the knees; also bony enlargement Possibly tender, seldom warm, and rarely red
	Prominent stiffness, often for 1 hour or more in the mornings, also after inactivity Limitation of motion common Weakness, fatigue, weight loss, and low fever	Frequent but brief stiffness (usually 5–10 min), in the morning and after inactivity Limitation of motion often develops

TABLE 16-4 Painful Tender Shoulders

Acromioclavicular Arthritis



Subacromial and Subdeltoid Bursitis



Rotator Cuff Tendinitis



Bicipital Tendinitis



Tenderness over the acromioclavicular joint, especially with adduction of the arm across the chest. Pain often increases with shrugging the shoulders, due to movement of scapula.

Pain over anterior superior aspect of shoulder, particularly when raising the arm overhead. Tenderness common anterolateral to the acromion, in hollow recess formed by the acromiohumeral sulcus. Often seen in overuse syndromes.

Tenderness over the rotator cuff, when elbow passively lifted posteriorly or with "drop-arm" maneuver.

Tenderness over the long head of the biceps when rolled in the bicipital groove or when flexed arm is supinated against resistance suggests bicipital tendinitis.

Painful Tender Knees







Arthritis. Degenerative arthritis usually occurs after age 50; associated with obesity. Often with medial joint line tenderness, palpable osteophytes, bowleg appearance, suprapatellar bursae and joint effusion. Systemic involvement, swelling, and subcutaneous nodules in rheumatoid arthritis

Bursitis, Inflammation and thickening of bursa seen in repetitive motion and overuse syndromes. Can involve prepatellar bursa ("housemaid's knee"), pes anserine bursa medially (runners, osteoarthritis), iliotibial band laterally (over lateral femoral condyle), especially in runners.

Painful Tender Knees (continued) TABLE







Patellofemoral instability. During flexion and extension of knee, due to subluxation and/or malalignment, patella tracks laterally instead of centrally in trochlear groove of femoral condyle. Inspect or palpate for lateral motion with leg extension. May lead to chondromalacia. osteoarthritis.

Meniscal tear. Commonly arises from twisting injury of knee; in older patients may be degenerative, often with clicking, popping, or locking sensation. Check for tenderness along joint line over medial or lateral meniscus and for effusion. May have associated tears of medial collateral of anterior cruciate ligaments.

Anterior cruciate tear or sprain. In twisting injuries of the knee, often with popping sensation, immediate swelling, pain with flexion/extension, difficulty walking, and sensation of knee "giving way." Check for anterior drawer sign, swelling of hemarthrosis, injuries to medial meniscus or medial collateral ligament. Consider evaluation by an orthopedic surgeon.

(table continues next page)

Painful Tender Knees (continued)





Posterior knee

Collateral ligament sprain or tear. From force applied to medial or lateral surface of knee (valgus or varus stress), producing localized swelling, pain, stiffness. Patients able to walk but may develop an effusion. Check for tenderness over affected ligament and ligamentous laxity during valgus or varus stress.

Baker's cyst. Cystic swelling palpable on the medial surface of the popliteal fossa, prompting complaints of aching or fullness behind the knee. Inspect, palpate for swelling adjacent to medial hamstring tendons. If present, suggests involvement of posterior horn of medial meniscus. In rheumatoid arthritis, cyst may expand into calf or ankle.

The Nervous System

FUNDAMENTALS FOR ASSESSING HE NERVOUS SYSTEM

The central nervous system (CNS) consists of the brain and spinal cord. The peripheral nervous system consists of the 12 pairs of cranial nerves and the spinal and peripheral nerves. Most peripheral nerves contain both motor and sensory fibers.



CENTRAL NERVOUS SYSTEM

The Brain

- Gray matter, or aggregations of neuronal cell bodies; rims the surfaces of the cerebral hemispheres, forming the cerebral cortex
- White matter, or neuronal axons coated with myelin, allowing nerve impulses to travel more rapidly
- Basal ganglia, which affect movement
- Thalamus, which processes and relays sensory impulses to the cerebral cortex
- Hypothalamus, which maintains homeostasis and regulates temperature, heart rate, and blood pressure; affects endocrine system, and governs emotional behaviors such as anger and sex drive; and contains hormones that act directly on the pituitary gland
- Brainstem, which connects the upper part of the brain with the spinal cord and has three sections: midbrain, pons, and medulla

312 The Nervous System

- Reticular activating (arousal) system, in the diencephalon and upper brainstem; activation linked to consciousness
- Cerebellum, at the base of the brain, which coordinates all movement and helps maintain the body upright in space

The Spinal Cord

- A cylindrical mass of nerve tissue encased within the bony vertebral column, extending from medulla to first or second lumbar vertebra
- Contains important motor and sensory nerve pathways that exit and enter the cord via anterior and posterior nerve roots and spinal and peripheral nerves
- Mediates reflex activity of the deep tendon (or spinal nerve) reflexes
- Divided into five segments: cervical (C1–8), thoracic (T1–12), lumbar (L1–5), sacral (S1–5), and coccygeal
- Roots fan out like a horse's tail at L1-2, the cauda equina



PERIPHERAL NERVOUS SYSTEM

The Cranial Nerves

- Cranial Nerves III through XII arise from the diencephalon and brainstem.
- Cranial Nerves I and II are actually fiber tracts emerging from the brain.

The Peripheral Nerves

- Thirty-one pairs of nerves carry impulses to and from the cord: 8 cervical, 12 thoracic, 5 lumbar, 5 sacral, and 1 coccygeal.
- Each nerve has an anterior (ventral) root containing motor fibers, and a posterior (dorsal) root containing sensory fibers.
- These merge to form a short (<5 mm) spinal nerve.
- Spinal nerve fibers commingle with similar fibers in plexuses outside the cord—from these emerge peripheral nerves.

THE HEALTH HISTORY

Common or Concerning Symptoms

- Headache
- Dizziness or vertigo
- Generalized, proximal, or distal weakness
- Numbness, abnormal or lost sensations
- Loss of consciousness, syncope, or near-syncope
- Seizures
- Tremors or involuntary movements

Headache: ask about location, severity, duration, and any associated symptoms, such as visual changes, weakness, or loss of sensation. Ask if coughing, sneezing, or sudden movements of the head affect the headache.

Dizziness can have many meanings. Is the patient lightheaded or feeling faint? Or is there *vertigo*, a perception that the room is spinning or rotating?

Are any medications contributing to dizziness? Are associated symptoms present, such as double vision (diplopia), difficulty forming words (dysarthria), or difficulty with gait or balance (ataxia)? Is there any weakness?

See Table 7-1, Primary
Headaches, p. 125, and
Table 7-2, Secondary
Headaches, pp. 126–129.
Subarachnoid hemorrhage may
evoke "the worst headache of
my life." Dull headache
affected by maneuvers,
especially on awakening and in
the same location, in mass
lesions such as brain tumors

Lightheadedness in palpitations; near-syncope from vasovagal stimulation, low blood pressure, febrile illness, and others; vertigo in inner ear conditions, brainstem tumor

Diplopia, dysarthria, ataxia in vertebrobasilar transient ischemic attack (TIA) or stroke

Weakness or paralysis in TIA or stroke

314 The Nervous System

Distinguish proximal from distal weakness. For proximal weakness, ask about combing hair, reaching for things on a high shelf, difficulty getting out of a chair or taking a high step up. For distal weakness ask about hand movements such as opening a jar or can or using hand tools (e.g., scissors, pliers, screwdriver). Ask about frequent tripping.

Bilateral proximal weakness in *myopathy;* bilateral, predominantly distal weakness in *polyneuropathy;* weakness worsened by repeated effort and improved by rest in *myasthenia gravis*

Is there any **loss of sensation**, difficulty moving a limb, or altered sensation such as tingling or pins and needles? Peculiar sensations without an obvious stimulus (*paresthesias*)? *Dysesthesias*, or disordered sensations in response to a stimulus, may last longer than the stimulus itself.

Loss of sensation, paresthesias, and dysesthesias in brain and spinal cord lesions; also in disorders of peripheral sensory roots and nerves; paresthesias in hands and around mouth in hyperventilation

"Have you ever fainted or passed out?" leads the discussion to any loss of consciousness.

Get a complete description of the event. What brought on the episode? Were there any warning symptoms? Was the patient standing, sitting, or lying down when it began? How long did it last? Could voices be heard while passing out and coming to? How rapid was recovery? Were onset and offset slow or fast?

Syncope if sudden but temporary loss of consciousness from decreased cerebral blood flow commonly called *fainting*.

Young people with emotional stress and warning symptoms of flushing, warmth, or nausea may have vasodepressor (or vasovagal) syncope of slow onset, slow offset. Cardiac syncope from dysrhythmias, more common in older patients, often with sudden onset, sudden offset.

Also ask if anyone observed the episode. What did the patient look like before, during, and after the episode? Was there any seizure-like movement of the arms or legs? Any incontinence of the bladder or bowel?

A seizure is a paroxysmal disorder caused by sudden excessive electrical discharge in the cerebral cortex or its underlying structures. Depending on the type, there may be loss of consciousness or abnormal feelings, thought processes, and sensations, including smells, as well as abnormal movements.

Tonic-clonic motor activity, incontinence, and postictal state in generalized seizures. Unlike syncope, injury such as tongue biting or bruising of limbs may occur.

PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling

- Preventing stroke or transient ischemic attack (TIA)
- Preventing risk of peripheral neuropathy

Preventing Stroke or TIA. Cerebrovascular disease is the third leading cause of death in the United States. Decreased vascular perfusion results in sudden focal but transient brain dysfunction in TIA, or in permanent neurological deficits in stroke.

Counsel patients about the warning signs of stroke attacks: sudden numbness or weakness of the face, arm, or leg; sudden confusion or trouble speaking or understanding; sudden difficulty walking, dizziness, or loss of balance or coordina-

316 The Nervous System

tion; sudden trouble seeing in one or both eyes; or sudden severe headache. Detecting TIAs is important—in the first 3 months after a TIA, subsequent stroke occurs in approximately 15% of patients.

Stroke prevention requires aggressive management of risk factors and patient education. Risk factors include smoking, excess weight, hypertension, dyslipidemia, heavy alcohol use, physical inactivity, obesity, and diabetes. Blood pressure should be $\leq 140/90$ mm Hg and $\leq 130/80$ mm Hg for those with diabetes or nondiabetic renal disease with proteinuria. Lipid-lowering agents may reduce risk of stroke. Urge patients to replace saturated and transunsaturated fats, found in dairy products, meat, and stick margarine, with polyunsaturated and unhydrogenated monosaturated fats, found in soybeans, liquid margarine, and fish oils. Or recommend increased intake of fruits, vegetables, and fiber. Counsel patients to maintain regular exercise and body weight and to control their intake of alcohol. Aim for optimal control of blood glucose levels, at approximately 100 mg/dL for patients with diabetes.

Preventing Risk of Peripheral Neuropathy. Promote optimal glucose control in diabetes to reduce risk of sensorimotor polyneuropathy, autonomic dysfunction, mononeuritis multiplex, or diabetic neuropathy.

TECHNIQUES OF EXAMINATION

• Cro	anial Nerves and Funct	ion
No.	Cranial Nerve	Function
I	Olfactory	Sense of smell
II	Optic	Vision
III	Oculomotor	Pupillary constriction, opening the eye (lid elevation), and most extraocular movements

• Cro	unial Nerves and Functio	ON (continued)
No.	Cranial Nerve	Function
IV	Trochlear	Downward, inward movement of the eye
VI	Abducens	Lateral deviation of the eye
V	Trigeminal	Motor—temporal and masseter muscles (jaw clenching), also lateral pterygoid's (lateral jaw movement) Sensory—facial. The nerve has three divisions: (1) ophthalmic, (2) maxillary, and (3) mandibular.
VII	Facial	Motor—facial movements, including those of facial expression, closing the eye, and closing the mouth Sensory—taste for salty, sweet, sour, and bitter substances on the anterior two thirds of the tongue
VIII	Acoustic	Hearing (cochlear division) and balance (vestibular division)
IX	Glossopharyngeal	Motor—pharynx Sensory—posterior portions of the eardrum and ear canal, the pharynx, and the posterior tongue, including taste (salty, sweet, sour, bitter)
X	Vagus	Motor—palate, pharynx, and larynx Sensory—pharynx and larynx
XI	Spinal accessory	<i>Motor</i> —the sternomastoid and upper portion of the trapezius
XII	Hypoglossal	Motor—tongue

POSSIBLE FINDINGS

CRANIAL NERVES

CN I (OLFACTORY)

Test sense of smell on each side.

Loss in frontal lobe lesions

CN II (OPTIC)

Assess visual acuity.

Blindness

Check visual fields.

Hemianopsia

Inspect optic discs.

Papilledema, optic atrophy

CN II, III (OPTIC AND OCULOMOTOR)

Test pupillary reactions to light. If abnormal, test reactions to near effort.

Blindness, CN III paralysis, tonic pupils; Horner's syndrome may affect light reactions

reactions


CN III, IV, VI (OCULOMOTOR, TROCHLEAR, AND ABDUCENS)

Assess extraocular movements.

Strabismus from paralysis of CN III, IV, or VI; nystagmus

CN V (TRIGEMINAL)

Test pain and light touch sensations on face in ophthalmic, maxillary, and mandibular zones.



CN V—SENSORY

Feel the contractions of temporal and masseter muscles.



TEMPORAL MUSCLES

POSSIBLE FINDINGS

Motor or sensory loss from lesions of CN V or its higher motor pathways



MASSETER MUSCLES

Check corneal reflexes.



CN VII (FACIAL)

Ask patient to raise both eyebrows, frown, close eyes tightly, show teeth, smile, and puff out cheeks.

CN VIII (ACOUSTIC)

Assess hearing. If decreased:

 Test for lateralization (Weber test, p. 122). Weakness from lesion of peripheral nerve, as in Bell's palsy, or of CNS, as in a stroke. See Table 17-3, Facial Paralysis, p. 339.

Sensorineural loss causes lateralization to affected ear where AC > BC. Conduction loss causes lateralization to affected ear and BC > AC.

POSSIBLE FINDINGS

• Compare air and bone conduction (**Rinne test**).

See p. 122.

CN IX, X (GLOSSOPHARYNGEAL AND VAGUS)

Observe any difficulty swallowing.

A weakened palate or pharynx impairs swallowing.

Listen to the voice.

Hoarseness or nasality

Watch soft palate rise with "ah."

Palatal paralysis in CVA

Test gag reflex on each side. Absent reflex

CN XI (SPINAL ACCESSORY)

Trapezius Muscles. Assess muscles for bulk, involuntary movements, and strength of shoulder shrug.

Atrophy, fasciculations, weakness



Sternomastoid Muscles. Assess strength as head turns against your hand.

Weakness of sternomastoid muscle when head turns to *opposite* side

CN XII (HYPOGLOSSAL)

Listen to patient's articulation.

Inspect the resting tongue.

Inspect the protruded tongue.

Dysarthria from damage to CN X or CN XII

Atrophy, fasciculations

Deviation to weak side



BODY POSITION

Observe the patient's body position during movement and at rest.

Hemiplegia in stroke

INVOLUNTARY MOVEMENTS

If present, observe location, quality, rate, rhythm, amplitude, and setting.

Tremors, fasciculations, tics, chorea, athetosis, oral-facial dvskinesias

MUSCLE BULK

Inspect muscle contours.

Atrophy

MUSCLE TONE

Assess resistance to passive stretch of arms and legs.

Spasticity, rigidity, flaccidity

MUSCLE STRENGTH

Test and grade the major muscle groups:

Gradi	ng Muscle Strength
Grade	Description
0	No muscular contraction detected
1	A barely detectable trace of contraction
2	Active movement with gravity eliminated
3	Active movement against gravity
4	Active movement against gravity and some resistance
5	Active movement against full resistance (normal)

Look for a pattern in any detectable weakness. It may suggest a lower motor neuron lesion affecting a peripheral nerve or nerve root. Weakness of one side of body suggests an upper

motor neuron lesion. A *polyneuropathy* causes symmetric distal weakness, and a *myopathy* usually causes proximal weakness. Weakness that worsens with repeated effort and improves with rest suggests *myasthenia gravis*.

- Elbow flexion (C5, C6)—biceps
- Elbow extension (C6, C7, C8)—triceps
- Wrist extension (C6, C7, C8)—radial nerve
- Grip (C7, C8, T1)

• Finger abduction (C8, T1)—ulnar nerve

Radial nerve damage: stroke or multiple sclerosis if hemiplesia

Weak grip in cervical radiculopathy, DeQuervain's tenosynovitis, carpal tunnel syndrome

Weak in ulnar nerve disorders



• Thumb opposition (C8, T1)—median nerve



Carpal tunnel syndrome

- Trunk—flexion, extension, lateral bending
- \ / \cup Hip flexion (L2, L3, L4)—iliopsoas
- Hip extension (S1) gluteus maximus
- Hip adduction (L2, L3, L4)—adductors
- Hip abduction (L4, L5, S1)—sulcus medius and minimus
- Knee extension (L2, L3, L4)—quadriceps
- Knee flexion (L4, L5, S1, S2) hamstrings
- Ankle dorsiflexion (L4, L5)
- Ankle plantar flexion (S1)

COORDINATION

Check:

Rapid alternating movements in arms and legs (tap foot)





Clumsy, slow movements in cerebellar disease

Point-to-point movements in arms and legs

Gait. Ask patient to

- Walk away, turn, and come back
- Walk heel to toe
- Walk on toes, then on heels
- Hop in place on each foot; do one-legged shallow knee bends. Substitute rising from a chair and climbing on a stool for hops and bends as indicated.

Stance

- Do a Romberg test (a sensory test of stance). Ask patient to stand with feet together and eyes open, then closed for 20 to 30 seconds. Mild swaying may occur. Stand close by to prevent falls.
- Look for a pronator drift. Watch as patient holds arms forward, with eyes closed, for 20 to 30 seconds.



POSSIBLE FINDINGS

Clumsy, unsteady movements in cerebellar disease

CVA, cerebellar ataxia, parkinsonism, or loss of position sense may all affect performance.

Corticospinal tract injury

Proximal hip girdle weakness increases risk of falls.

Loss of balance that appears when eyes are closed is a *positive* Romberg test, suggesting poor position sense.

Flexion and pronation at elbow and downward drift of arm from contralateral corticospinal tract lesion



Ask patient to keep arms up and tap them downward. A smooth return to position is normal.

Weakness, incoordination, poor position sense

THE SENSORY SYSTEM

/o— Using an object such as a broken cotton swab to test sharp and dull sensation, compare symmetric areas on the two sides of the body. Do not reuse the object on another patient.

Hemisensory deficits

Also **compare** proximal and distal areas of arms and legs for pain, temperature, and touch sensation. Scatter stimuli to sample most dermatomes and major peripheral nerves.

"Glove-and-stocking" loss of peripheral neuropathy

Map any area of abnormal response, including dermatomes, if present.

Dermatomal sensory loss in herpes zoster, nerve root compression. See Table 17-7, pp. 344-345, for dermatomes.

Assess response to the following stimuli. Except when you are explaining the tests, patient's eyes should be closed.

• Pain. Use the sharp end of a pin or other suitable tool. The dull end serves as a control.

Analgesia, hypalgesia, hyperalgesia

- *Temperature* (if indicated). Use test tubes with hot and ice-cold water, or other objects of suitable temperature.
- *Light touch*. Use a fine wisp of cotton.

Check for *vibration and position senses*. If responses are abnormal, test more proximally.

- Vibration and position.
 Vibration: Use a 128-Hz or 256-Hz tuning fork, held on a bony prominence.
 Vibration and position senses, both carried in the posterior columns, often correlate.
- *Position*. Holding patient's finger or big toe by its sides, move it up or down.

POSSIBLE FINDINGS

Temperature and pain senses usually correlate.

Anesthesia, hyperesthesia

Loss of vibration and position senses in peripheral neuropathy from diabetes or alcoholism and in posterior column disease from syphilis or B₁₂ deficiency





Assess one or more of the *discriminative* sensations:

• Stereognosis. Ask for identification of a common

Lesions in the posterior columns or sensory cortex

object placed in patient's hand.

• Number identification. Ask for identification of a number drawn on patient's palm with blunt end of a pen.

- Two-point discrimination. Find minimal distance on pad of patient's finger at which the sides of two points can be distinguished from one (normally <5 mm).
- Point localization. Touch skin briefly, and ask patient to open both eyes and identify the place touched.
- Extinction. Simultaneously touch opposite, corresponding areas of the body, and ask patient where the touch is felt.

POSSIBLE FINDINGS

impair stereognosis, number identification, and two-point discrimination.





A lesion in the sensory cortex may impair point localization on the opposite side and cause extinction of the touch sensation on that side.



Grading Re	eflexes
Grade	Description
4+	Hyperactive (with clonus)
3+	Brisker than average, not necessarily abnormal
2+	Average, normal
1+	Diminished, low normal
0	No response

Biceps (C5, C6)



Hyperactive deep tendon reflexes, absent abdominal reflexes, and a Babinski response in *upper motor* neuron lesions

Triceps (C6, C7)



POSSIBLE FINDINGS

Supinator (brachioradialis) (C5, C6)





Ankle (S1)



Ankle jerks symmetrically; decreased or absent in peripheral polyneuropathy **Check** for clonus if reflexes seem hyperactive.



CUTANEOUS STIMULATION REFLEXES

• Abdominals (upper T8, T9, T10; lower T10, T11, T12)

May be absent with upper or lower neuron lesions



Plantar (L5, S1), normally flexor



Babinski response from corticospinal tract lesion



POSSIBLE FINDINGS

Anal Reflex. With a dull object, stroke outward from anus in four quadrants. Watch for anal contraction.

Loss of reflex suggests lesion at \$2-3-4 level.

SPECIAL TECHNIQUES

- Meningeal Signs. With patient supine, flex head and neck toward chest. Note resistance or pain, and watch for flexion of hips and knees (Brudzinski's sign).

Meningeal irritation may cause resistance or pain on flexion during both maneuvers.

Flex one of patient's legs at hip and knee, then straighten knee. Note resistance or pain (Kernig's sign).

A compressed lumbosacral nerve root also causes pain on straightening the knee of the raised leg.



- Lumbosacral Radicu**lopathy.** Straight leg raise.

With patient supine, raise relaxed and straightened leg, flexing the leg at the hip. Then dorsiflex the foot.

Pain and muscle weakness if herniated disc; may also see calf wasting and weak ankle dorsiflexion



Asterixis. Ask patient to hold both arms forward, with hands cocked up and fingers spread. Watch for 1 to 2 minutes.

Winging of the Scapula. Ask patient to push against the wall of your hand with a partially straightened arm. Inspect scapula. It should stay

close to the chest wall.

POSSIBLE FINDINGS

Sudden brief flexions in liver disease, uremia hypercapnia.

Winging of scapula away from chest wall suggests weakness of the serratus anterior muscle.



The Stuporous or Comatose Patient. — Assess ABCs (airway, breathing, and circulation).

Take pulse, blood pressure, and rectal temperature.

Establish level of consciousness with escalating stimuli.

Don't dilate pupils, and **don't flex patient's neck** if cervical cord may have been injured.

Lethargy, obtundation, stupor, coma

Levels of Consc	iousness
Alertness	Patient is awake and aware of self and environment. When spoken to in a normal voice, patient looks at you and responds fully and appropriately to stimuli.
Lethargy	When spoken to in a loud voice, patient appears drowsy but opens eyes and looks at you, responds to questions, and then falls asleep.
Obtundation	When shaken gently, patient opens eyes and looks at you but responds slowly and is somewhat confused. Alertness and interest in environment are decreased.
Stupor	Patient arouses from sleep only after painful stimuli. Verbal responses are slow or absent. Patient lapses into unresponsiveness when stimulus stops. Patient has minimal awareness of self or environment.
Coma	Despite repeated painful stimuli, patient remains unarousable with eyes closed. No evident response to inner need or external stimuli is shown.

Observe	
Breathing pattern	Cheyne-Stokes, ataxic breathing
• Pupils	Asymmetric if structural lesions or possible brain herniation
Ocular movements	Deviation to affected side in hemispheric stroke
Note posture of body.	Decorticate rigidity, decerebrate rigidity, flaccid hemiplegia

Check for the oculocephalic reflex (doll's eye movements). Holding upper eyelids open, turn head quickly to each side, and then flex and extend patient's neck. This patient's head will be turned to her right.

POSSIBLE FINDINGS

In a comatose patient with an intact brainstem, the eyes move in the opposite direction, in this case to her left (doll's eye movements) as below.





Test for flaccid paralysis.

- Hold forearms vertically; note wrist positions.
- From 12 to 18 inches above bed, drop each arm.
- Support both knees in a somewhat flexed position, and then extend each knee and let lower leg drop to the bed.
- From a similar starting position, release both legs.

Complete the neurologic and general physical examination.

Very deep coma or a lesion in midbrain or pons abolishes this reflex, so eyes do not move.

A flaccid hand droops to the horizontal.

A flaccid arm drops more rapidly.

The flaccid leg drops more rapidly.

A flaccid leg falls into extension and external rotation.

RECORDING YOUR FINDINGS

Recording the Physical Examination— The Nervous System

"Mental Status: Alert, relaxed, and cooperative. Thought process coherent. Oriented to person, place, and time. Detailed cognitive testing deferred. Cranial Nerves: I—not tested; II through XII intact. Motor: Good muscle bulk and tone. Strength 5/5 throughout. Cerebellar: Rapid alternating movements (RAMs), finger-to-nose (F \(\times N\)), heelto-shin (H→S) intact. Gait with normal base. Romberg maintains balance with eyes closed. No pronator drift. Sensory: Pinprick, light touch, position, and vibration intact. Reflexes: 2+ and symmetric with plantar reflexes downgoing."

Alternatively, record reflexes using table or stick figure:

	Biceps	Triceps	Brach	Knee	Ankle	PI	+ + + +
RT LT	2+ 2+	2+ 2+	2+ 2+	2+ 2+	2+ 1+	$\downarrow \\ \downarrow$	OR # #
							# # #

AIDS TO INTERPRETATION

TABLE 17-1 Types of Stroke

Assessing patients with stroke involves three fundamental questions based on a careful history and detailed physical examination: What brain area and related vascular territory explain the patient's findings? Is the stroke ischemic or hemorrhagic? If ischemic, is the mechanism thrombus or embolus? Stroke is a medical emergency, and timing is of the essence. Answers to these questions are critical to patient outcomes and use of antithrombotic therapies in acute ischemic stroke.

In *acute ischemic stroke*, ischemic brain injury begins with a central core of very low perfusion and often irreversible cell death. This core is surrounded by an *ischemic penumbra* of metabolically disturbed cells that are still potentially viable, depending on restoration of blood flow and duration of ischemia. Because most irreversible damage occurs in the first 3 to 6 hours after onset of symptoms, therapies targeted to the 3-hour window achieve the best outcomes, with recovery in up to 50% of patients in some studies.

Clinician performance in diagnosing stroke improves with training. Understanding the pathophysiology of stroke takes dedication, expert supervision to improve techniques of neurological examination, and perseverance. *This brief overview is intended to prompt further study and practice*. Accuracy in clinical examination is achievable, and more important than ever in determining patient therapy.

TABLE 17Types of Stroke (continued)




TABLE 17-2 Clinical Features and Vascular Territories of Stroke


Major Clinical Features	Vascular Territory	Additional Comments
Contralateral leg weakness	Anterior circulation—anterior cerebral artery (ACA)	Includes stem of circle of Willis connecting internal carotid artery to ACA, and the segment distal to ACA and its anterior choroidal branch
Contralateral face, arm > leg weakness, sensory loss, field cut, aphasia (left MCA) or neglect, apraxia (right MCA)	Anterior circulation—middle cerebral artery (MCA)	Largest vascular bed for stroke
Contralateral motor or sensory deficit without cortical signs	Subcortical circulation—lenticulostriate deep penetrating branches of MCA	Small vessel subcortical <i>lacunar infarets</i> in internal capsule, thalamus, or brainstem. Four common syndromes: pure motor hemiparesis; pure sensory hemianesthesia; ataxic hemiparesis; clumsy hand-dysarthria syndrome
Contralateral field cut	Posterior circulation—posterior cerebral artery (PCA)	Includes paired vertebral arteries, the basilar artery, paired posterior cerebral arteries. Bilateral PCA infarction causes cortical blindness but preserved pupillary light reaction.
Dysphagia, dysarthria, tongue/palate deviation and/or ataxia with crossed sensory/motor deficits (= ipsilateral face with contralateral body)	Posterior circulation—brainstem, vertebral, or basilar artery branches	
Oculomotor deficits and/or ataxia with crossed sensory/motor deficits	Posterior circulation—basilar artery	Complete basilar artery occlusion—"locked-in syndrome" with intact consciousness but inability to speak and quadriplegia

(Source: Adapted from American College of Physicians. Stroke, in Neurology. Medical Knowledge Self-Assessment Program (MKSAP) 14. Philadelphia: American College of Physicians, 2006. pp. 52-68.)

TABLE 17-3**Facial Paralysis**

Distinguish peripheral from central lesions of CN VII by closely observing movements of the upper face. Because of innervation from both hemispheres, the movements are preserved in central lesions

	Lesion of Peripheral Nervous System	Lesion of Central Nervous System
Side of face affected	Same side as the lesion	Side opposite the lesion
Upper face	Unable to wrinkle forehead, raise eyebrow, close eye	Movements normal or slightly weak
Lower face	Unable to smile, show teeth	Same



Bell's palsy Common cause **CVA** (injury to ČŃ VII)

	2	
	e	٩
	2	2
-	2	5
	L	J
	E	•
-	2	2
		7
_		r
1	ú	n
	N	
ightharpoonup		
u	S	
		ī

	Peripheral Nervous System Disorder	Central Nervous System Disorder*	Parkinsonism (Basal Ganglia Disorder)	Cerebellar Disorder
Involuntary movements	Often fasciculations	No fasciculations	Resting tremors	Intention tremors
Muscle bulk	Atrophy	Normal or mild atrophy (disuse)	Normal	Normal
Muscle tone	Decreased or absent	Increased, spastic	Increased, rigid	Decreased
Muscle strength	Decreased or lost	Decreased or lost	Normal or slightly decreased	Normal or slightly decreased
Coordination	Unimpaired, though limited by weakness	Slowed and limited by weakness	Good, though slowed and often tremulous	Impaired, ataxic
Reflexes Deep tendon Plantar Abdominals	Decreased or absent Flexor or absent Absent	Increased Extensor Absent	Normal or decreased Flexor Normal	Normal or decreased Flexor Normal

*Upper motor neuron

Involuntary Movements



Resting static tremors. Seen at rest, usually disappear with movement: seen in Parkinsonism



Postural tremor. Seen when maintaining active posture; in anxiety, hyperthyroidism; also familial



Intention tremor. Seen with intentional movement, absent at rest; in cerebellar disorders, including multiple sclerosis



Fasciculations. Fine, rapid flickering of muscle bundles



Chorea. Brief, rapid, irregular, jerky; face, head, arms, or hands



Athetosis. Slow, twisting, writhing; face, distal limbs

TABLE 17-5 Involuntary Movements (continued)



Oral-facial dyskinesias.

Rhythmic, repetitive, bizarre movements of face, mouth.

Tardive dyskinesias with prolonged use of psychotropic drugs such as phenothiazines



Tics. Brief, irregular, repetitive, coordinated movements (e.g., winking, shrugging); in Tourette's syndrome, users of phenothiazines, amphetamines



Dystonia. Grotesque, twisted postures, often in trunk or, as shown, in neck (spasmodic torticollis)

Disorders of Muscle Tone

Spasticity

Location. Upper motor neuron of the corticospinal tract at any point from the cortex to the spinal cord

Description. Increased muscle tone (hypertonia) that is ratedependent. Tone is greater when passive movement is rapid, and less when passive movement is slow. Tone is also greater at the extremes of the movement arc. During rapid passive movement, initial hypertonia may give way suddenly as the limb relaxes. This spastic "catch" and relaxation is known as "clasp-knife" resistance.

Common Cause. Stroke, especially late or chronic stage

Flaccidity

Location, Lower motor neuron at any point from the anterior horn cell to the peripheral nerves

Description. Loss of muscle tone (hypotonia), causing the limb to be loose or floppy. The affected limbs may be hyperextensible or even flaillike.

Common Cause. Guillain-Barré syndrome; also initial phase of spinal cord injury (spinal shock) or stroke

Rigidity

Location. Basal ganglia system

Description. Increased resistance that persists throughout the movement arc, independent of rate of movement, is called lead-pipe rigidity. With flexion and extension of the wrist or forearm. a superimposed rachetlike jerkiness is called cogwheel rigidity.

Common Cause. Parkinsonism


Paratonia

Location. Both hemispheres, usually in the frontal lobes

Description. Sudden changes in tone with passive range of motion. Sudden loss of tone that increases the ease of motion is called mitgehen (moving with). Sudden increase in tone making motion more difficult is called gegenhalten (holding against).

Common Cause. Dementia

TABLE 17-7 Dermatomes




DERMATOMES INNERVATED BY POSTERIOR ROOTS

(table continues next page)

TABLE 17-7

Dermatomes (continued)



DERMATOMES INNERVATED BY POSTERIOR ROOTS

	Toxic-Metabolic	Structural
Pathophysiology	Arousal centers poisoned or critical substrates depleted	Lesion destroys or compresses brainstem arousal areas, either directly or secondary to more distant expanding mass lesions.
• Respiratory pattern	If regular, may be normal or hyperventilation. If irregular, usually Cheyne-Stokes	Irregular, especially Cheyne-Stokes or ataxic breathing. Also with selected stereotypical patterns like "apneustic" respiration (peak inspiratory arrest) or central hyperventilation.
Pupillary size and reaction	Equal, reactive to light. If pinpoint from opiates or cholinergies, you may need a magnifying glass to see the reaction. May be unreactive if fixed and dilated from anticholinergies or hypothermia	Unequal or unreactive to light (fixed) Midposition, fixed—suggests midbrain compression Dilated, fixed—suggests compression of CN III from herniation
• Level of consciousness Examples of Cause	Changes after pupils change Uremia, hyperglycemia Alcohol, drugs, liver failure Hypothyroidism, hypoglycemia Anoxia, ischemia Meningitis, encephalitis Hyperthermia, hypothermia	Changes before pupils change Epidural, subdural, or intracerebral hemorrhage Cerebral infarct or embolus Tumor, abscess Brainstem infarct, tumor, or hemorrhage Cerebellar infarct, hemorrhage, tumor, or abscess

Pupils in Comatose Patients

Small or Pinpoint Pupils



Bilaterally small pupils (1-2.5 mm) suggest (1) damage to the sympathetic pathways in the hypothalamus or (2) metabolic encephalopathy (a diffuse failure of cerebral function that has many causes, including drugs). Light reactions are usually normal.

Pinpoint pupils (<1 mm) suggest (1) a hemorrhage in the pons or (2) the effects of morphine, heroin, or other narcotics. The light reactions may be seen with a magnifying glass.

Midposition Fixed Pupils



Midposition or slightly dilated pupils (4-6 mm) and fixed to light suggest damage in the midbrain.

Large Pupils



Bilaterally fixed and dilated pupils in severe anoxia with sympathomimetic effects, may be seen with cardiac arrest. They also result from atropine-like agents, phenothiazines, or tricyclic antidepressants.

One Large Pupil



One fixed and dilated pupil warns of herniation of the temporal lobe, causing compression of the oculomotor nerve and midbrain.

18

Assessing Children: Infancy Through Adolescence

CHILD DEVELOPMENT

Children display tremendous variations in physical, cognitive, and social development compared with adults.

Key Principles of Child Development

- Child development proceeds along a predictable pathway marked by developmental milestones.
- The range of normal development is wide. Children mature at different rates.
- Various physical, psychological, social, and environmental factors, as well as diseases, can affect child development and health. For example, chronic diseases, child abuse, and poverty can contribute to detectable physical abnormalities and influence the rate and course of developmental advancement.
- The child's developmental level affects how you conduct the medical history and physical examination.

THE HEALTH HISTORY

The child's history follows the same outline as the adult's history, with certain *additions* presented here.

Identifying Data. Record date and place of birth, nickname, and first names of parents (and last name of each, if different).

Chief Complaints. Determine if they are the concerns of the child, the parent(s), a schoolteacher, or some other person.

Present Illness. Determine how each family member responds to the child's symptoms, why he or she is concerned, and whether the illness may provide for the child any secondary gain.

Past History

Birth History. This is especially important when neurologic or developmental problems are present. Get hospital records if necessary.

- Prenatal—maternal health: medications; tobacco, drug, and alcohol use; weight gain; duration of pregnancy
- Natal—nature of labor and delivery, birth weight, Apgar scores at 1 and 5 minutes
- Neonatal—resuscitation efforts, cyanosis, jaundice, infections, bonding

Feeding History. This is particularly important with either undernutrition or obesity.

- Breast-feeding—frequency and duration of feeds, difficulties, timing and method of weaning
- Bottle-feeding—type; amount; frequency; vomiting; colic; diarrhea
- Vitamins, iron, and fluoride supplements; introduction of solid foods
- Eating habits—types and amounts of food eaten, parental attitudes and responses to feeding problems

Growth and Developmental History. This is particularly important with delayed growth or development and behavioral disturbances.

- Physical growth—weight and height at all ages; head circumference at birth and less than 2 years; periods of slow or rapid growth
- Developmental milestones—ages child held head up, rolled over, sat, stood, walked, and talked
- Speech development, performance in preschool and school
- Social development—day and night sleeping patterns; toilet training; habitual behaviors; discipline problems; school behavior; relationships with family and peers

Current Health Status

Allergies. Pay particular attention to history of eczema, urticaria, perennial allergic rhinitis, asthma, food intolerance, insect hypersensitivity, and recurrent wheezing.

Immunizations. Include dates given and any untoward reactions.

Screening Tests. These are likely to vary according to the child's medical and social conditions. Include newborn screening results, anemia screening, blood lead, sickle cell disease, vision, hearing, developmental screening, and others (e.g., tuberculosis).

PROMOTION AND COUNSELING H E A L I H

- 1. Age-appropriate developmental achievement of the child
 - Physical (maturation, growth, puberty)
 - Motor (gross and fine motor skills)
 - Cognitive (achievement of milestones, language, school performance)
 - Emotional (self-efficacy, self-esteem, independence, morality)
 - Social (social competence, self-responsibility, integration with family and community)
- 2. Health supervision visits (per health supervision schedule)
 - Periodic assessment of medical and oral health
 - Adjustment of frequency for children or families with special needs
- 3. Integration of physical examination findings
- 4. Immunizations
- 5. Screening procedures
- 6. Anticipatory guidance
 - Healthy habits
 - Nutrition and healthy eating
 - Emotional and mental health
 - Oral health
 - Safety and prevention of injury
 - Sexual development and sexuality
 - Self-responsibility and efficacy
 - Family relationships (interactions, strengths, supports)
 - Prevention or recognition of illness
 - Prevention of risky behaviors and addictions
 - School and vocation
 - Peer relationships
 - Community interactions
- 7. Partnership between health care provider and child, adolescent, and family

ECHNIQUES OF EXAMINATION

Sequence of Examination

The sequence of examination varies according to the child's age and comfort level.

- For infants and young children, perform nondisturbing maneuvers early and potentially distressing maneuvers toward the end. For example, palpate the head and neck and auscultate the heart and lungs early; examine the ears and mouth and palpate the abdomen near the end. If the child reports pain in an area, examine that part last.
- For older children and adolescents, use the same sequence as with adults, except examine the most painful areas last.

ASSESSING NEWBORNS

EXAMINATION TECHNIQUES

IMMEDIATE ASSESSMENT AT BIRTH

Listen to the anterior thorax with your stethoscope. Palpate the abdomen. Inspect the head, face, oral cavity, extremities, genitalia, and perineum.

Apgar Score. Score each newborn according to the following table, at 1 and 5 minutes after birth, according to the 3-point scale (0, 1, or 2) for each component.

If the 5-minute score is 8 or more, proceed to a more complete examination.

• The	Apgar Sc	oring System			
Assign	ed Score	2			
Clinical	Sign	0		1	2
Heart 1	ate	Absent	<100)	>100
Respira effort		Absent	Slow irre	and gular	Good; strong
Muscle	tone	Flaccid	of t		Active movement
Reflex irrital	bility*	No responses	Grim	ace	Crying vigorously, sneeze, or cough
Color		Blue, pale	blu	body, e remities	Pink all over
1-Minu	ite Apga	nr Score	5-Minu	ıte Apgar	Score
8-10	Norma	ıl	8-10	Normal	
5–7		nervous system ession	0–7	seque	sk for sub- nt central
0–4	requi imm	depression, iring ediate citation			us system and organ system action

^{*}Reaction to suction of nares with bulb syringe.

Gestational Age and Birth Weight. Classify newborns according to their gestational age and birth weight.

CLASSIFICATION BY GESTATION AND BIRTH WEIGHT	NAL AGE
Gestational Age	
Classification	Gestational Age
PretermTermPostterm	<37 wks (<259th day) 37–42 wks >42 wks (>294th day)
Birth Weight	
Classification	Weight
Extremely low birth weightVery low birth weightLow birth weightNormal birth weight	<1,000 grams <1,500 grams <2,500 grams ≥2,500 grams

Newborn Classifications		
Category	Abbreviation	Percentile
Small for gestational age	SGA	<10th
Appropriate for gestational age	AGA	10-90th
Large for gestational age	LGA	>90th

EXAMINATION SEVERAL HOURS AFTER BIRTH

During the first day of life, newborns should have a comprehensive examination following the technique outlined under "Infants." Wait until 1 or 2 hours after a feeding, when the newborn is more responsive. Ask parents to remain.

Observe the baby's color, size, body proportions, nutritional status, posture, respirations, and movements of the head and extremities.

Normally findings are spontaneous motor activity, flexion and extension of the extremities, and brief body tremors.

Most newborns are bowlegged, reflecting their curled up intrauterine position.

Inspect the newborn's umbilical cord to detect abnormalities. Normally, there are two thick-walled umbilical arteries and one larger but thin-walled umbilical vein, which is usually located at the 12 o'clock position.

A single umbilical artery may be associated with congenital anomalies. Umbilical hernias in infants are from a defect in the abdominal wall.

The neurologic screening examination of all newborns should include assessment of mental status, gross and fine motor function, tone, cry, deep tendon reflexes, and primitive reflexes.

Signs of severe neurologic disease include extreme irritability; persistent asymmetry of posture or extension of extremities; constant turning of head to one side; marked extension of head, neck, and extremities (opisthotonus): severe flaccidity; and limited pain response.

ASSESSING INFANTS

MENTAL AND PHYSICAL STATUS

Observe the parents' affect when talking about the baby; their manner of holding,

Common causes of developmental delay include abnormalities in embryonic

POSSIBLE FINDINGS

moving, and dressing the baby; and their response to situations that may produce discomfort for the baby. **Observe** a breast or bottle feeding. **Determine** attainment of developmental milestones using the Denver Developmental Screening Test before conducting the physical examination.

development, hereditary and genetic disorders, environmental and social problems, other pregnancy or perinatal problems, childhood diseases such as infection (e.g., meningitis), trauma, and severe chronic disease.

GENERAL SURVEY

Growth, reflected in increases in height and weight within expected limits, is an excellent indicator of health during infancy and childhood. Deviations from normal may be early indications of an underlying problem. To assess growth, compare a child's parameters with respect to

Causes of *failure to thrive* can be environmental or psychosocial, or various gastrointestinal, neurologic, cardiac, endocrine, renal, and other diseases.

- Normal values according to age and sex
- Prior readings to assess trends

Height and Weight. Plot each child's height and weight on standard growth charts to determine progress. Measures above the 97th or below the 3rd percentile, or recent rises or falls from prior levels, require investigation.

Reduced growth in height may indicate *endocrine disease*, other *causes of short stature*, or, if weight is also low, other *chronic diseases*.

POSSIBLE FINDINGS

Head Circumference.

Determine head circumference at every physical examination during the first 2 years.



Premature closure of the sutures or microcephaly may cause small head size. Hydrocephalus, subdural hematoma, or, rarely, brain tumor or inherited syndromes may cause abnormally large head size.

VITAL SIGNS

Blood Pressure. Measure blood pressure at least once during infancy. Although the hand-held method is shown here, the most easily used measure of systolic blood pressure in infants and young children is obtained with the *Doppler method*.



Newborn	Middle Childhood
Renal artery disease (stenosis, thrombosis)	Renal parenchymal or arteria disease
Congenital renal malformations	Primary hypertension
Coarctation of the aorta	Coarctation of the aorta
Infancy and Early Childhood	Adolescence
Renal parenchymal or artery disease	Primary hypertension Renal parenchymal disease
Coarctation of the aorta	Drug induced

POSSIBLE FINDINGS

Pulse. The heart rate is quite variable.

Tachycardia (>180–200/min) usually indicates *paroxysmal* supraventricular tachycardia. Bradycardia may result from serious underlying disease.

Respiratory Rate. The respiratory rate has a greater range and is more responsive to illness, exercise, and emotion than in adults.

Respiratory diseases such as bronchiolitis or pneumonia may cause rapid respirations (up to 80–90/min), but also increased work of breathing.

THE SKIN

Assess

Texture and appearance

• Texture and appearance

• Vasomotor changes

 Pigmentation (e.g., Mongolian spots)

• Hair (e.g., lanugo)

 Common skin conditions (e.g., milia, erythema toxicum)

Jaundice

Turgor

Cutis marmorata

Acrocyanosis; cyanotic congenital heart disease

Café-au-lait spots

Midline hair tuft on back

Herpes simplex


Hemolytic disease

Dehydration

THE HEAD

Examine *sutures* and *fontanelles* carefully.

Head small with microcephaly, enlarged with hydrocephaly; fontanelles full and tense with meningitis, closed with microcephaly, separated with increased intracranial pressure



(hydrocephaly, subdural hematoma, and brain tumor)

Swelling from subperiosteal hemorrhage (cephalohematoma) does not cross suture lines; swelling from bleeding associated with a fracture does.

Check the *face* for symmetry. Examine for an overall impression of the facies; comparing with the faces of the parents is helpful.

PEARLS TO EVALUATE POTENTIALLY ABNORMAL FACIES

Carefully review the history, especially the family history, pregnancy, and perinatal history.

Note abnormalities, especially of growth, development, or dysmorphic somatic features.

Measure and plot percentiles, especially of head circumference, height, and weight.

Consider the three mechanisms of facial dysmorphogenesis:

- Deformations from intrauterine constraint
- Disruptions from amniotic bands or fetal tissue
- Malformations from an intrinsic abnormality (either face/head or brain)

Examine parents and siblings (similarity may be reassuring but might point to a familial disorder).

Determine whether facial features fit a recognizable syndrome. Compare against references, pictures, tables, and databases.

THE EYES

Newborns and young infants may look at your face and follow a bright light if you catch them while alert.

Normal visual milestones are as follows:

Nystagmus, strabismus

Visual Milestones of Infancy	
Birth	Blinks, may regard face
1 month	Fixes on objects
$1^{1}/_{2}$ –2 months	Coordinated eye movements
3 months	Eyes converge, baby reaches
12 months	Acuity around 20/50

THE EARS

Check position, shape, and features.

Small, deformed or low-set auricles may indicate associated *congenital defects*, especially renal disease.

Signs That a	n Infant Can Hear
Age	Signs
0–2 months	Startle response and blink to a sudden noise Calming down with soothing voice or music
2–3 months	Change in body movements in response to sound Change in facial expression to familiar sounds
3–4 months	Turning eyes and head to sound
6–7 months	Turning to listen to voices and conversation

THE NOSE

Test patency of the nasal passages by occluding alternately each nostril while holding the infant's mouth closed.

With choanal atresia, the baby cannot breathe if one nostril is occluded.

THE MOUTH AND PHARYNX

Inspect (with a tongue blade and flashlight) and palpate.


Supernumerary teeth, Epstein's pearls

You may see a whitish covering on the tongue. If this coating is from milk, you can easily remove it by scraping or wiping it away.

Oral candidiasis (thrush)

THE NECK

Palpate the lymph nodes, and assess for any additional masses (e.g., congenital cysts).



Lymphadenopathy is usually from viral or bacterial infections. Other neck masses include malignancy, branchial cleft or thyroalossal duct cysts. periauricular cysts and sinuses.

THE THORAX AND LUNGS

Carefully assess respirations and breathing pattern.

Apnea

Don't rush to the stethoscope, but observe the patient carefully first.

Upper respiratory infections may cause nasal flaring.

Examination of	f the Lungs in Infants—Befo	re You Touch the Child!
Assessment	Possible Findings	Explanation
General appearance	Inability to feed or smile Lack of consolability	Lower respiratory infections below the vocal cords (e.g., bronchiolitis, pneumonia) are common in infants.
Respiratory rate	Tachypnea	Cardiac or respiratory disease
Color	Pallor or cyanosis	Cardiac or pulmonary disease
Nasal component of breathing	Nasal flaring (enlargement of both nasal openings during inspiration)	Upper or lower respiratory infection
Audible breath sounds	Grunting (repetitive, short expiratory sound) Wheezing (musical expiratory sound) Stridor (high-pitched, inspiratory noise) Obstruction (lack of breath sounds)	Acute stridor is a potentially serious condition with causes such as laryngotracheobronchitis (croup), epiglottitis, bacterial tracheitis, foreign body, vascular ring
Work of breathing	Nasal flaring Grunting Retractions (or chest indrawing): Supraclavicular (motion of soft tissue above clavicles) Intercostal (indrawing of the skin between ribs) Subcostal (just below the costal margin)	In infants, abnormal work of breathing combined with abnormal findings on auscultation is the best finding for ruling in pneumonia.

Distinguishing Upper	er Airway From Lower Ai	irway Sounds
Technique	Upper Airway	Lower Airway
Compare sounds from nose/stethoscope	Same sounds	Often different sounds
Listen to harshness of sounds	Harsh and loud	Variable
Note symmetry (left/right)	Symmetric	Often asymmetric
Compare sounds at different locations (higher or lower)	Sounds louder as stethoscope is moved up chest	Sounds louder lower in chest
Inspiratory vs. expiratory	Almost always inspiratory	Often has expiratory phase

THE HEART

Inspection. Observe carefully for any cyanosis.

At birth: *Transposition of the* great arteries; pulmonary valve atresia or stenosis

Within a few days of birth:

The above: also total anomalous pulmonary venous return, hypoplastic left heart

Weeks, months, or years of life: The above; also pulmonary vascular disease with atrial, ventricular, or great vessel shunting

Palpation. Palpate the peripheral pulses. The point of maximal impulse (PMI) is not always palpable in

No or diminished femoral pulses suggest coarctation of the aorta. Weak or thready, difficult-to-feel pulses may

POSSIBLE FINDINGS

infants. *Thrills* are palpable when enough turbulence is within the heart or great vessels.

reflect myocardial dysfunction and congestive heart failure.

Auscultation. Heart rhythm is evaluated more easily in infants by listening to the heart than by feeling the peripheral pulses.

The most common dysrhythmia in children is paroxysmal supraventricular tachycardia.

Heart Sounds. Evaluate S_1 and S_2 carefully. They are normally crisp.

A louder-than-normal pulmonic component, particularly louder than the aortic sound, suggests pulmonary hypertension.
Persistent splitting of S₂ may indicate a right ventricular volume load, such as atrial septal defect.

THE BREASTS

The breasts of males and females may be enlarged for months after birth as a result of maternal estrogen, and even engorged for 1 to 2 weeks with a white liquid.

THE ABDOMEN

You will find it easy to palpate an infant's abdomen, because infants like being touched.

Abnormal abdominal masses can be associated with kidney, bladder, or bowel tumors. In *pyloric stenosis*, deep palpation in the right upper quadrant or midline can reveal an "olive," or a 2-cm firm pyloric mass.

MALE GENITALIA

Inspect with the infant supine.

In 3% of infants, one or both testes cannot be felt in the scrotum or inguinal canal. Try to milk the testes into the scrotum.

Common scrotal masses are hydroceles and inquinal hernias.

FEMALE GENITALIA

In females, genitalia may be prominent for several months after birth from the effects of maternal estrogen. Ambiguous genitalia involves masculinization of the female external genitalia.

THE MUSCULOSKELETAL SYSTEM

The focus is detecting congenital abnormalities, particularly in the hands, spine, hips, legs, and feet.

Examine the *hips* carefully at each visit for signs of dislocation. There are two major techniques: one to test for a posteriorly dislocated hip (Ortolani test) and the other to test for the ability to sublux or dislocate an intact but unstable hip (Barlow test). Skin tags, remnants of digits, polydactyly (extra fingers), or syndactyly (webbed fingers) are congenital defects. Fracture of the clavicle can occur during a difficult delivery.

Congenital hip dysplasia may have a positive Ortolani or Barlow test, particularly during the first 3 months of age.

With a hip dysplasia, you feel a "clunk."



ORTOLANI TEST



BARLOW TEST

POSSIBLE FINDINGS

Some normal infants exhibit twisting or *torsion of the tibia* inwardly or outwardly on its longitudinal axis.

Pathologic tibial torsion occurs only in association with deformities of the feet or hips.

THE NERVOUS SYSTEM

Evaluate the developing central nervous system by assessing *infantile automatisms*, called *primitive reflexes*.

Suspect a *neurologic* or *developmental abnormality* if primitive reflexes are

- Absent at appropriate age
- Present longer than normal
- Asymmetric
- Associated with posturing or twitching

Some neurologic abnormalities produce deficits or slow cognitive and social development. Infants with developmental delay may have abnormal findings on neurologic examination because much of it is based on age-specific norms.

Children with *spastic diplegias* will often have hypotonia as infants.

ASSESSING CHILDREN (1 TO 10 YEARS)

Tips for Interviewing Children

 Establish rapport. Refer to children by name and meet them on their own level. Maintain eye contact at their level (e.g., sit on the floor if needed). Participate in play and talk about their interests.

- Work with families. Ask simple, open-ended questions such as "Are you sick? Tell me about it," followed by more specific questions. Once the parent has started the conversation, direct questions back to the child. Also observe how parents interact with the child.
- Identify multiple agendas. Your job is to discover as many perspectives and agendas as possible.
- Use the family as the key resource. View parents as experts in the care of their child and you as their consultant.
- Note hidden agendas. As with adults, the chief complaint may not relate to the real reason the parent has brought the child to see you.

The following discussion focuses on those areas of the comprehensive physical examination that are different for children than for infants and for adults.

MENTAL AND PHYSICAL STATUS

In children 1 to 5 years, **observe** the degree of sickness or wellness, mood, nutritional state, speech, cry, facial expression, and developmental skills. Note parent-child interaction, including separation tolerance, affection, and response to discipline.

In children 6 to 10 years, determine orientation to time and place, factual knowledge, and language and number skills. Observe motor skills used in writing, tying laces, buttoning, cutting, and drawing.

This overall examination can uncover evidence of chronic disease, developmental delay, social or environmental disorders, and family problems.

POSSIBLE FINDINGS

Body Mass Index for Age.

Age- and sex-specific charts are now available to assess body mass index (BMI) in children.

Underweight is <5th percentile, at risk of over-weight is ≥85th percentile, and overweight is ≥95th percentile.

BLOOD PRESSURE

Hypertension during childhood is more common than previously thought. Recognizing, confirming, and appropriately managing it is important. Blood pressure readings should be part of the physical examination of every child older than 2 years. Proper cuff size is essential for accurate determination of blood pressure in children.

The most frequent "cause" of elevated blood pressure in children is probably an improperly performed examination, often from an incorrect cuff size.

Causes of sustained hypertension in childhood include renal disease, coarctation of the aorta, and primary hypertension.

THE EYES

The two most important aspects of the eye examination for young children are to test visual acuity in each eye and to determine whether the gaze is conjugate or symmetric.

Strabismus; amblyopia; myopia; or hyperopia

SPECIAL TECHNIQUES

The corneal light reflex test (*left*) and the cover-uncover test (*right*) are particularly useful in young children.

Any difference in visual acuity between eyes is abnormal.

POSSIBLE FINDINGS





Visual Acuity	
Age	Visual Acuity
3 months 12 months Younger than 4 years 4 years and older	Eyes converge, baby reaches ~20/200 20/40 20/30

THE EARS

Examine the ear canal and drum. There are two positions for the child (lying down or sitting), and also two ways to hold the otoscope, as illustrated.





SPECIAL TECHNIOUE

Pneumatic Otoscope.

Learn to use a *pneumatic* otoscope to improve your accuracy of diagnosis of otitis media in children.

- Check for leaks by placing your finger over the tip of the speculum and squeezing the bulb.
- Insert the speculum, obtaining a proper seal.
- When air is introduced into the normal ear canal, the tympanic membrane and its light reflex move inward. When air is removed, the tympanic membrane moves outward toward you.

Acute otitis media involves a red and bulging tympanic membrane.

Diminished movement of tympanic membrane with acute otitis media; no movement with otitis media with effusion. Pain on movement of the pinna with otitis externa.

THE MOUTH AND PHARYNX

For anxious or young children, you may want to leave this examination until toward the end. If you need to use the tongue blade, the best technique is to push down and pull slightly forward toward you while the child says "ah." Be careful not to place the blade too far posteriorly, eliciting a gag reflex.

Examine the teeth for the timing and sequence of eruption, number, character, condition, and position.

Abnormalities of the enamel may reflect local or general disease.

Carefully **inspect** the inside of the upper teeth, as shown.

Nursing bottle caries; dental caries; staining of the teeth, which may be intrinsic or extrinsic



Dental caries are the most common health problem of children and are particularly prevalent in impoverished children.

Look for abnormalities of tooth position.

Malocclusion

Note the size, position, symmetry, and appearance of the tonsils.


Peritonsillar abscess

THE HEART

One of the most challenging aspects to cardiac examination of children is evaluation of heart murmurs. In addition to trying to listen to a squirming, perhaps uncooperative child, a major challenge is to

See Table 18-4, Characteristics of Pathologic Heart Murmurs, pp. 381-382.





distinguish common benign murmurs from unusual or pathologic ones. Most (indeed, some say nearly all) children will have one or more functional, or benign, heart murmurs before adulthood (see above).

THE ABDOMEN

Most children are ticklish when you first place your hand on their abdomens for palpation. This reaction tends to disappear, particularly if you distract the child.

A pathologically enlarged liver in children usually is palpable more than 2 cm below the costal margin, has a round, firm edge, and often is tender.

MALE GENITALIA

There is an art to palpation of the young boy's scrotum and testes, because many have an extremely active cremasteric reflex that may cause the testes to retract upward into the inguinal canal and thereby appear undescended. A useful

In *precocious puberty,* the penis and testes are enlarged, with signs of pubertal changes.

A painful testicle requires rapid treatment and may indicate *torsion*.

technique is to have the boy sit cross-legged on the examining table.

Inquinal hernias in older boys present as they do in adult men.

FEMALE GENITALIA

For the genital examination, use a calm, gentle approach, including a developmentally appropriate explanation.

Examine the genitalia in an efficient and systematic manner. The normal hymen can have various configurations.



Vaginal discharge in early childhood can result from perineal irritation (e.g., from bubble baths, soaps), foreign body, vaginitis, or sexually transmitted diseases from sexual abuse. Vaginal bleeding is always concerning. Abrasions or signs of trauma to the external genitalia can result from sexual abuse.

THE MUSCULOSKELETAL SYSTEM

In older children, abnormalities of the upper extremities are rare in the absence of injury. Observe the child standing and walking barefoot. You also can ask the child to touch the toes, rise from sitting, run a short distance, and pick up objects. You will detect most abnormalities by watching carefully.

A screening musculoskeletal examination for children participating in sports can detect injuries or abnormalities that may result in problems during athletics.

THE NERVOUS SYSTEM

Beyond infancy (when the primitive reflexes have disappeared), the neurologic examination includes the components evaluated in adults. Again, combine the neurologic and developmental assessments. You will need to turn this into a game with the child. The goal is to assess optimal development and neurologic performance, which requires the child's cooperation.

Delayed language or cognitive skills can be due to neurologic disease as well as developmental disorders.

ASSESSING ADOLESCENTS

The key to successfully examining teens is a comfortable, confidential environment that makes the examination relaxed and informative. Adolescents are more likely to open up when the interview focuses on them rather than on their problems.

Consider the patient's cognitive and social development when deciding issues of privacy, parental involvement, and confidentiality. Explain to both teen and parent that the purpose of confidentiality is to improve health care, not keep secrets. Your goal is to help adolescents bring their concerns or questions to their parents. Never make confidentiality unlimited, however. Always state to teens explicitly that you may need to act on information that makes you concerned about safety.

The physical examination of the adolescent is similar to that of the adult. Keep in mind issues particularly relevant to teens, such as puberty, growth, development, family and peer relationships, sexuality, decision making, and risk behaviors. For more details on specific techniques of examination, the reader should refer to the corresponding chapter for the regional examination of interest or concern. Following are special areas to highlight when examining adolescents.

THE BREASTS

Assess normal maturational development.

See Table 18-5, Sex Maturity Ratings in Girls: Breasts, p. 383

SPECIAL TECHNIQUE

Testing for Scoliosis.

Inspect any child who can stand for scoliosis. Make sure the child bends forward with the knees straight (Adams bend test). Evaluate any asymmetry in positioning or gait. If you detect scoliosis, use a scoliometer to test for the degree of scoliosis.



MALE AND FEMALE GENITALIA

An important goal when examining adolescent males and females is to assign a sexual maturity rating, regardless of chronologic age.

See Table 18-6, Sex Maturity Ratings in Boys, pp. 384-385, and Table 18-7, Sex Maturity Ratings in Girls: Pubic Hair, pp. 386-387.

RECORDING YOUR FINDINGS

The format of the pediatric medical record is the same as that of the adult. Thus, although the sequence of the physical examination may vary, convert your written findings back to the traditional format.

Recording the Physical Examination— The Pediatric Patient

Brian is a chubby, active, and energetic toddler. He plays with the reflex hammer, pretending it is a truck. He appears closely bonded with his mother, looking at her occasionally for comfort. She seems concerned that Brian will break something. His clothes are clean.

Vital Signs. Ht 90 cm (90th percentile). Wt 16 kg (>95th percentile). BMI 19.8 (>95th percentile). Head circumference 50 cm (75th percentile). BP 108/58. Heart rate 90 and regular. Respiratory rate 30; varies with activity. Temperature (ear) 37.5°C. Obviously no pain. Skin: normal except for bruises on legs, and patchy, dry skin over external surface of elbows.

HEENT. *Head:* Normocephalic; no lesions. *Eyes:* Difficult to examine because he won't sit still. Symmetric with normal extraocular movements. Pupils 4 to 5 mm constricting. Discs difficult to visualize; no hemorrhages noted. *Ears:* Normal pinna; no external abnormalities. Normal external canals and tympanic membranes (TMs). *Nose:* Normal nares; septum midline. *Mouth:* Several darkened teeth on inside surface of upper incisors. One clear cavity on upper right incisor. Tongue normal. Cobblestoning of posterior pharynx; no exudates. Tonsils large but adequate gap (1.5 cm) between them.

Neck. Supple, midline trachea, no thyroid palpable.

Lymph Nodes. Easily palpable (1.5 to 2 cm) tonsillar lymph nodes bilaterally. Small (0.5 cm) nodes in inguinal canal bilaterally. All lymph nodes mobile and nontender.

Lungs. Good expansion. No tachypnea or dyspnea. Congestion audible, but seems to be upper airway (louder near mouth, symmetric). No rhonchi, rales, or wheezes. Clear to auscultation.

Cardiovascular. PMI in 4th or 5th interspace and midsternal line. Normal S1 and S2. No murmurs or

abnormal heart sounds. Normal femoral pulses; dorsalis pedis pulses palpable bilaterally.

Breasts. Normal, with some fat under both.


Abdomen. Protuberant but soft; no masses or tenderness. Liver span 2 cm below right costal margin (RCM) and not tender. Spleen and kidneys not palpable.

Genitalia. Tanner I circumcised penis; no pubic hair, lesions, or discharge. Testes descended, difficult to palpate because of active cremasteric reflex. Normal scrotum both sides.

Musculoskeletal. Normal range of motion of upper and lower extremities and all joints. Spine straight. Gait normal.

Neurologic. *Mental Status:* Happy, cooperative child. Developmental (DDST): Gross motor—Jumps and throws objects. Fine motor—Imitates vertical line. Language— Does not combine words; single words only, three to four noted during examination. Personal-social—Washes face, brushes teeth, and puts on shirt. Overall—Normal, except for language, which appears delayed. Cranial Nerves: Intact, although several difficult to elicit. Cerebellar: Normal gait; good balance. Deep tendon reflexes (DTRs): Normal and symmetric throughout with downgoing toes. Sensory: Deferred.

Classification of Newborn's **Level of Maturity**



Weight Small for Gestational Age (SGA) = Birth weight <10th percentile on the intrauterine growth curve

Weight Appropriate for Gestational Age (AGA) = Birth weight within the 10th and 90th percentiles on the intrauterine growth curve

Weight Large for Gestational Age (LGA) = Birth weight >90th percentile on the intrauterine growth curve

Level of intrauterine growth based on birth weight and gestational age of liveborn, single, white infants. Point A represents a premature infant, while point B indicates an infant of similar birth weight who is mature but small for gestational age; the growth curves are representative of the 10th and 90th percentiles for all of the newborns in the sampling.

Classification of the low-birth-weight infant. In: Klaus MH, Fanaroff AA. Care of the High-Risk Neonate, 3rd ed. Philadelphia: WB Saunders, 1986.

TABLE 18-2 Recommendations for Preventive Pediatric Health Care

Each child and family is unique; therefore, these recommendations are designed for the care of children who are receiving competent parenting, have no manifestation of any important health problems, and are growing and developing in satisfactory fashion. Additional visits may become necessary if circumstances suggest variation from normal.

	÷	•		ဟ ဟ	
	y 20	_	• •		
	- 19			တ တ	• •
	8	•	• •	00	• •
OE	17 y	•	• •	တ တ	• •
ADOLESCENCE	16 y	•	• •	တ တ	
OLES	15 y	•	• •	00	
AD	14 y	•	• •	တ တ	
	13 y	•	• •	တ တ	
	12 y	•	• •	00	
	> =	•	• •	တ တ	
	10y 11y 12y 13y 14y 15y 16y 17y 18y 19y 20y+	•		00	
MIDDLE	× 8	•	• •	00	
MIDDLE	6 y	•	• •	00	
Ö	> 10	•	• •	00	
	4 Y	٠	• •	00	
8	e ک	•	• •	0 0	
EARLY CHILDHOOD	4mo	•	• •	တ တ	
뿔	8 mo	•	• •	တ တ	
	5 mo	•	• •	တ တ	
	2 mo	•	• •	တ တ	
	9 mo	•	• •	တ တ	
ζÇ	9 mo	•	• •	တ တ	
INFANCY	4 mo	•	• •	တ တ	
	, mo	٠	• •	တ တ	
	- A	•	• •	တ တ	
	2-4 By add 2 mo 4 mo 6 mo 9 mo 12 mo15 mo 18 mo24mo	•	• •	တ တ	
	AGE	HISTORY Initial / Interval	MEASUREMENTS Height and Weight Head Circumference Blood Pressure	SENSORY SCREENING Vision Hearing	DEVELOPMENTAL/ BEHAVIORAL ASSESSMENT ² PHYSICAL EXAMINATION ³

Adapted from Recommendations For Preventive Pediatric Health Care promulgated by the American Academy of Pediatrics Committee on Practice and Ambulatory Medicine, 1999

By history and appropriate physical examination: if suspicious, by

^{3.} At each visit, a complete physical examination is essential, with infant totally unclothed, older child undressed and suitably draped specific objective development testing


to be performed S = subjective, by history O = objective, by a standard testing method


TABLE 18-3 Hypertension in Childhood

Hypertension can start in childhood. While young children with elevated blood pressure are more likely to have a renal, cardiac, or endocrine cause, adolescents with hypertension are most likely to have primary or essential hypertension.

This child developed hypertension during adolescence, and it "tracked" into adulthood. Children tend to remain in the same percentile for blood pressure as they grow. This tracking of blood pressure continues into adulthood, supporting the concept that adult essential hypertension begins during childhood.

The consequences of untreated hypertension can be severe.





Characteristics of Pathologic Heart Murmurs

Congenital Defect

Pulmonary Valve Stenosis Mild



Moderate



Severe



Aortic Valve Stenosis



Tetralogy of Fallot With pulmonic stenosis



Characteristics of Murmur

Location. Upper left sternal border

Radiation. In mild degrees of stenosis, the murmur may be heard over the course of the pulmonary arteries in the lung fields.

Intensity. Increases in intensity and duration as the degree of obstruction increases

Quality. Ejection, peaking later in systole as the obstruction increases

Location. Midsternum, upper right sternal border

Radiation. To the carotid arteries and suprasternal notch; may also be a thrill

Intensity. Varies, louder with increasingly severe obstruction

Quality. An ejection, often harsh, systolic murmur

General. Variable cyanosis, increasing with activity

Location. Mid to upper left sternal border. If pulmonary atresia, there is no systolic murmur but the continuous murmur of ductus arteriosus flow at upper left sternal border or in the back.

(table continues next page)



Characteristics of Pathologic Heart Murmurs (continued)

Congenital Defect

With pulmonic atresia



Transposition of the **Great Arteries**

Ventricular Septal Defect Small to moderate



Characteristics of Murmur

Radiation. Little, to upper left sternal border, occasionally to lung fields

Intensity. Usually Grade III-IV Quality. Midpeaking, systolic ejection murmur

General. Intense generalized cvanosis

Location. No characteristic murmur. If a murmur is present, it may reflect an associated defect such as VSD or patent ductus arteriosus.

Radiation. Depends on associated abnormalities Quality. Depends on associated abnormalities

Location. Lower left sternal border

Radiation Little

Intensity. Variable, only partially determined by the size of the shunt. Small shunts with a high pressure gradient may have very loud murmurs. Large defects with elevated pulmonary vascular resistance may have no murmur. Grade II-IV/VI with a thrill if Grade IV/VI or higher.

Stage 1

Preadolescent—elevation of nipple only

Stage 2



Breast bud stage. Elevation of breast and nipple as a small mound; enlargement of areolar diameter

Stage 3

Stage 5



Further enlargement and elevation of breast and areola, with no separation of the contours

Stage 4





Projection of areola and nipple to form a secondary mound above the level of the breast



Mature stage; projection of nipple only. Areola has receded to general contour of the breast (although in some normal individuals areola continues to form a secondary mound).

⁽Photos reprinted, with permission from the American Academy of Pediatrics.)

TABLE 18-6 Sex Maturity Ratings in Boys

In assigning SMRs in boys, observe each of the three characteristics separately. Record two separate ratings: pubic hair and genital. If the penis and testes differ in their stages, average the two into a single figure for the genital rating

		Genitalia	lia
	Pubic Hair	Penis	Testes and Scrotum
Stage 1	Preadolescent—no pubic hair except for the fine body hair (vellus hair) similar to that on the abdomen	Preadolescent—same size and proportions as in childhood	Preadolescent—same size and proportions as in childhood
Stage 2	Sparse growth of long, slightly pigmented, downy hair, straight or only slightly curled, chiefly at the base of the penis	Slight to no enlargement	Testes larger; scrotum larger, somewhat reddened, and altered in texture
Stage 3	Darker, coarser, curlier hair spreading sparsely over the pubic symphysis	Larger, especially in length	Further enlarged

Stage 4



Coarse and curly hair, as in the adult; area covered greater than in stage 3 but not as great as in the adult and not yet including the thighs

Further enlarged in length and breadth, with development of the glans

Further enlarged; scrotal

skin darkened

Adult in size and shape

Adult in size and shape

Hair adult quantity and quality, spread to the medial surfaces of the thighs but not up over the abdomen

Stage 5

(Photos reprinted from Pediatric Endocrinology and Growth, 2nd ed., Wales & Wit, 2003, with permission from Elsevier.)

TABLE 18-7

Sex Maturity Ratings in Girls: Pubic Hair

Stage 1

Preadolescent—no pubic hair except for the fine body hair (vellus hair) similar to that on the abdomen

Stage 2



Sparse growth of long, slightly pigmented, downy hair, straight or only slightly curled, chiefly along the labia

Stage 3



Darker, coarser, curlier hair, spreading sparsely over the pubic symphysis

Stage 4



Coarse and curly hair as in adults; area covered greater than in stage 3 but not as great as in the adult and not yet including the thighs

TABLE 18

Sex Maturity Ratings in Girls: Pubic Hair (continued)

Stage 5



Hair adult in quantity and quality, spread on the medial surfaces of the thighs but not up over the abdomen

Photos reprinted, with permission from the American Academy of Pediatrics.)

TABLE 18-8 Physical Signs of Sexual Abuse

Physical Signs that May Indicate Sexual Abuse in Children*

- Marked and immediate dilatation of the anus in knee-chest position, with no constipation, stool in the vault, or neurologic disorders
- 2. Hymenal notch or cleft that extends greater than 50% of the inferior hymenal rim (confirmed in knee–chest position)
- 3. Condyloma acuminata in a child older than 3 years
- 4. Bruising, abrasions, lacerations, or bite marks of labia or perihymenal tissue
- 5. Herpes of the anogenital area beyond the neonatal period
- 6. Purulent or malodorous vaginal discharge in a young girl (all discharges should be cultured and viewed under a microscope for evidence of a sexually transmitted disease)

Physical Signs that Strongly Suggest Sexual Abuse in Children*

- 1. Lacerations, ecchymoses, and newly healed scars of the hymen or the posterior fourchette
- 2. No hymenal tissue from 3 to 9 o'clock (confirmed in various positions)
- 3. Healed hymenal transections, especially between 3 and 9 o'clock (complete cleft)
- 4. Perianal lacerations extending to external sphincter

A sexual-abuse expert must evaluate a child with concerning physical signs for a complete history and sexual abuse examination.

^{*}Any physical sign must be evaluated in light of the entire history, other parts of the physical examination, and laboratory data.

The Pregnant Woman

THE HEALTH HISTORY

Common Concerns

- Symptoms of pregnancy
- Smoking, alcohol, use of illicit drugs, domestic violence
- Prior complications of pregnancy
- Chronic illnesses and family history
- Determining weeks of gestation by date and expected date of delivery

Focus the *initial prenatal visit* on confirming the pregnancy, assessing the health status of the mother and any risks for complications, and counseling to ensure a healthy pregnancy. Ask about the following topics:

- Symptoms of pregnancy: absence of menses, breast fullness or tenderness, nausea or vomiting, fatigue, and urinary frequency. Explain that serum or urine testing for beta human chorionic gonadotropin (HCG) offers the best confirmation of pregnancy.
- Maternal concerns and attitudes: Review the mother's feelings about the pregnancy and whether she plans to continue to term. Ask about any fears and about support from the father.
- Current state of health: risk factors that could adversely affect the mother or fetus. Ask about eating patterns and assess the quality of her nutrition. Does she smoke, drink alcohol, or use any illicit drugs? Does she take any medications? Or have any exposures to toxic substances? What about her income and her social support network? Are

390 The Pregnant Woman

there any sources of unusual stress at home or in the workplace? Is there any history of physical abuse or domestic violence?

- Past obstetric history: Has she had any complications during past pregnancies, including labor and delivery? Has she had a premature or growth-retarded infant, or a baby large for gestational age? Has there been a prior fetal demise?
- Past medical history: Review any history of acute or chronic illnesses, especially hypertension, diabetes, cardiac conditions, asthma, systemic lupus erythematosus (SLE), and seizures, and any history of sexually transmitted diseases, exposure to diethylstilbestrol in utero, or HIV infection.
- Family history of chronic illnesses or genetically transmitted diseases: sickle cell anemia, cystic fibrosis, muscular dystrophy, and others.

Expected weeks of gestation by dates and the expected date of delivery (EDD):

- To establish expected weeks of gestation, count in weeks from either (1) the first day of the last menstrual period (LMP), known as menstrual age, or (2) the date of conception if this is known, termed conception age. The expected weeks of gestation can help with later assessment of uterine size, assuming that the LMP was normal, the dates were remembered accurately, and conception actually occurred. During the examination, compare your estimate of expected uterine size with the palpable size of the uterus if still within the pelvic cavity, or with the height of the fundus if above the symphysis pubis. If there are discrepancies, investigate possible causes.
- The first day of the LMP is also used to calculate the EDD, or the time projected to term labor and delivery assuming regular 28- to 30-day menstrual cycles. Using *Naegele's rule*, estimate the EDD by adding 7 days to the first day of the LMP, subtracting 3 months, and adding 1 year. This date may be one of the first questions the mother asks you.
- If the patient cannot remember her LMP or has irregular menstrual cycles, or if the dating is uncertain, vaginal

- probe ultrasound is used to confirm dating in the first trimester.
- Establish the desired frequency of *follow-up visits*. These visits will include measurement of blood pressure and weight, palpation of the uterine fundus to assess fetal growth, verification of fetal heart tones, and determination of fetal presentation and activity.

Important Topics for Health Promotion and Counseling

- Nutrition
- Weight gain
- Exercise
- Smoking cessation, alcohol, and illicit drugs
- Screening for domestic violence
- Immunizations

Nutrition and Weight Gain. Evaluate nutritional status during the first prenatal visit, including: diet history; measurement of height, weight, and body mass index (BMI); and a hematocrit. Check use of needed vitamin and mineral supplements. Develop a nutrition plan appropriate to cultural preferences, typically three balanced meals each day, with increased intake as follows: 300 additional kcal; 5 to 6 g of protein; 15 mg of iron; 250 mg of calcium; and 400 to 800 mcg of folic acid, plus prenatal supplements. Caution against excess amounts of vitamin A, which can become toxic; fish with mercury exposure such as sharks, swordfish, or even canned tuna; unpasteurized dairy products; and undercooked meats.

Ideal weight gain in pregnancy follows a pattern: very little gain the first trimester, rapid increase in the second trimester, and mild slowing of the increase in the third trimester. Average weight gain is approximately 28 lbs, or ~10 kg. Weigh the woman at each visit, with the results plotted on a graph. Recommended weight gain ranges from the Institute of Medicine (1992), displayed next, are still current.

Recommended Total Weight Gain Ranges for Pregnant Women		
Prepregnancy Weight-for-Height Category	Recommend lbs	ded Total Gain kg
Low BMI <19.8	28-40	12.5–18
Normal BMI 19.8–26.0	25-35	11.5–16
High BMI 26.0-29.0	15–25	7.0-11.5
Obese BMI >29.0	~15	~7.0

Figures are for single pregnancies. The range for women carrying twins is 35–45 lb (16–20 kg). Young adolescents (<2 years after menarche) should strive for gains at the upper end of the range. Short women (<62 in. or <157 cm) should strive for gains at the lower end of the range.

Exercise. The American College of Obstetricians and Gynecologists (2002) recommends that in conjunction with their physician and in the absence of contraindications, pregnant women should engage in moderate exercise for ~30 minutes most days of the week. Women initiating exercise during pregnancy should consider programs developed specifically for pregnant women. After the first trimester, women should avoid exercise in the supine position, which can compress the inferior vena cava and decrease blood flow to the placenta. The pregnant woman should stop exercise when she feels fatigued or uncomfortable and avoid overheating and dehydration.

Smoking Cessation. Any smoking should be discontinued. Smoking has been linked to *complications of labor* such as placental abruption, placenta previa, preterm deliveries, low-birthweight babies, and even perinatal death.

Screening for Domestic Violence. Prevalence during pregnancy ranges from 7% to 20% and may result in femicide, or murder of the mother and child. The American College of Obstetricians and Gynecologists recommends assessment of all women for any history of domestic violence that may escalate during the pregnancy. Clues include frequent changes in appointments at the last minute, behavior during the interview, chronic headache or abdominal pain, and bruises or other signs of injury.

Clinicians should ask direct questions in a nonjudgmental manner in a private setting at each prenatal visit. For example, ask "Since you've been pregnant, have you been slapped or otherwise physically hurt by anyone?" Women may need multiple opportunities to discuss abuse because of fears about safety and reprisal. Validate positive responses and mark the area of injury on a body diagram. Above all, in situations of admitted abuse, ask how you might help. Offer information on safe shelters, counseling centers, hotline telephone numbers, and other sources of assistance when she is ready to pursue these.* Assess the patient's safety and make any necessary referrals. Learn about requirements for mandatory reporting.

Tetanus and influenza vaccines can be given in any trimester. Pneumococcal, meningococcal, and hepatitis B vaccines are also safe.

^{*}National Domestic Violence Hotline: Web site: www.ncvh.org/index.html; 1-800-799-SAFE (7233); TTY for hearing impaired, 1-800-787-3224.

TECHNIQUES OF EXAMINATION

PREPARING FOR THE EXAMINATION

Show respect for the woman's comfort and privacy, as well as for her individual needs and sensitivities. Ask her to wear her gown with the opening in front to ease the examination of both breasts and the pregnant abdomen.

Positioning

- The semisitting position with the knees bent (see p. 396) affords the most comfort and protects abdominal organs and vessels from the weight of the gravid uterus.
- Avoid prolonged periods of lying on the back. Make your abdominal palpation efficient and accurate.
- The pelvic examination also should be relatively quick.

Equipment

- The examiner's hands should be warm and firm yet gentle in palpation. The fingers should be together and flat. Make palpation smooth and continuous rather than kneading, using the more sensitive palmar surfaces of the ends of the fingers.
- You may need a speculum larger than the usual size.
 Because of the increased vascularity of the vaginal and cervical structures, insert and open the speculum gently to avoid tissue trauma and bleeding.
- Avoid the cervical brush for Pap smears, because it often causes bleeding. Use the Ayre wooden spatula or "broom" sampling device.

VITAL SIGNS, HEIGHT, AND WEIGHT

Measure the blood pressure. In midpregnancy, it may be lower than in the nonpregnant state.

HYPERTENSION IN PREGNANCY

- Gestational hypertension if systolic blood pressure (SBP) ≥140 mm Hg and diastolic blood pressure (DBP) ≥90 mm Hg, first occurring after week 20 and without proteinuria
- Chronic hypertension if SBP ≥140 mm Hg and DBP ≥90 mm Hg prior to pregnancy, before week 20, and after 12 weeks postpartum
- Preeclampsia if SBP ≥140 mm Hg and DBP ≥90 mm Hg after week 20 and with proteinuria

Measure the height and weight. Calculate BMI. First-trimester weight loss should not exceed 5% of prepartum weight.

Weight loss of more than 5% in excessive vomiting, or hyperemesis

HEAD AND NECK

• Face. Check for the mask of pregnancy, chloasma, or irregular brownish patches around the forehead and cheeks, across the bridge of the nose, or along the jaw.

Facial edema after 24 weeks in gestational hypertension

Hair

Hair loss should not be attributed to pregnancy.

• Eyes. Note the conjunctival color.

Anemia of pregnancy may cause conjunctival pallor.

 Nose, including nasal congestion

Nosebleeds are more common.

Mouth

Gingival enlargement common

POSSIBLE FINDINGS

 Thyroid gland. Inspect and palpate. Modest symmetric enlargement is common. Significant enlargement is abnormal and should be investigated.

THORAX AND LUNGS

Inspect the thorax for contours. Observe the pattern of breathing.

Respiratory alkalosis in later trimesters

HEART

Palpate the apical impulse.

Impulse may be higher than normal in the 4th intercostal space because of transverse and leftward rotation of the heart from the higher diaphragm.

Auscultate the heart. A venous hum and systolic or continuous mammary souffle (see p. 180) are common.

New diastolic murmurs should be investigated.

BREASTS

Inspect the breasts and nipples for symmetry and color.

Palpate for masses.

Compress each nipple between your index finger and thumb.

The venous pattern may be marked, the nipples and areolae are dark, and Montgomery's glands are prominent.

During pregnancy, breasts are tender and nodular.

This may express colostrum from the nipples.

ABDOMEN

Place the pregnant woman in a semisitting position with her knees flexed.



- Inspect any scars or striae, the shape and contour of the abdomen, and the fundal height.
- Assess the shape and contour to estimate pregnancy size.

• Palpate for:

Organs or masses

Fetal movements, usually detected after 24 weeks


Uterine contractility

If woman is >20 weeks pregnant, measure fundal height with a tape measure from the top of the symphysis pubis to the top of the uterine fundus. After 20 weeks, measurement in centimeters should roughly equal the weeks of gestation.

Auscultate the fetal heart, noting its rate (FHR), location, and rhythm. A Doptone will detect the

POSSIBLE FINDINGS

Purplish striae and linea nigra are normal.



EXPECTED HEIGHT OF UTERINE FUNDUS BY WEEKS OF PREGNANCY

Irregular contractions after 12 weeks or after palpation during the third trimester

Prior to 37 weeks, regular uterine contractions or bleeding are abnormal, suggesting *preterm labor*.

If fundal height is more than 2 cm higher than expected, consider multiple gestation, a big baby, extra amniotic fluid, or uterine leiomyoma. If more than 4 cm lower, consider missed abortion, transverse lie, growth retardation, or false pregnancy.

Lack of an audible FHR may indicate pregnancy of fewer weeks than expected, fetal demise, or false pregnancy. FHR after 10 weeks. The FHR is audible with a fetoscope after 18 weeks.

- From 12 to 18 weeks, listen in the midline of the lower abdomen.
- After 28 weeks, listen over the fetal back or chest. Use modified *Leopold's* maneuvers to palpate the fetal head and back and identify where to listen.

The *rate* usually is in the 160s during early pregnancy, and then slows to the 120s to 140s near term. After 32 to 34 weeks, the FHR should increase with fetal movement.

An FHR that drops noticeably near term with fetal movement could indicate poor placental circulation.

Rhythm. In the third trimester, expect a variance of 10 to 15 beats per minute (BPM) over 1 to 2 minutes.

Lack of beat-to-beat variability late in pregnancy warrants investigation with an FHR monitor.

GENITALIA, ANUS, AND RECTUM

Inspect the *external* genitalia.

Parous relaxation of the introitus, enlargement of the labia and clitoris, scars from an *episiotomy* or perineal lacerations

Inspect the anus.

Hemorrhoids may engorge later in pregnancy.

Palpate *Bartholin's* and *Skene's glands*.

Check for a cystocele or rectocele.

POSSIBLE FINDINGS

Speculum Examination.

Inspect the *cervix* for color, shape, and healed lacerations.

Purplish color of pregnancy; lacerations from prior deliveries

Take Pap smears, if indicated.

Specimens may be needed for diagnosis of vaginal or cervical infection

Inspect the vaginal walls.

Bluish or violet color, deep rugae, leukorrhea in normal pregnancy; vaginal irritation, itching, and discharge in infection

Bimanual Examination.

Insert two lubricated fingers into introitus, palmar side down, with slight pressure downward on the perineum. Slide fingers into the posterior vaginal vault. Maintaining downward pressure, gently turn fingers palmar side up.

Closed os if nulliparous; os open to size of fingertip if multiparous

- Assess cervical os and degree of effacement. Place your finger gently in the os, and then sweep it around the surface of the cervix.
- Estimate the *length of the cervix*. Palpate the lateral surface from the cervical tip to the lateral fornix.
- Palpate the *uterus* for size, shape, consistency, and position.

Prior to 34 to 36 weeks, cervix should retain normal length of ~1.5 to 2 cm.

Hegar's sign, or early softening of the isthmus; pear-shaped uterus up to 8 weeks, then globular

- Estimate uterine size. With your internal fingers placed at either side of cervix, palmar surfaces upward, gently lift the uterus toward the abdominal hand. Capture the fundal portion of the uterus
- Palpate the left and right adnexa.

between your two hands and gently estimate size.

 Perform a rectovaginal examination to confirm uterine size or the integrity of the rectovaginal septum.

POSSIBLE FINDINGS

An irregularly shaped uterus suggests uterine myomata or a bicornuate uterus, two distinct uterine cavities separated by a septum.

Early in pregnancy, it is important to rule out tubal (*ectopic*) pregnancy.

EXTREMITIES

Inspect the legs for *varicose* veins.

Inspect the hands and legs for *edema*.

Check knee and ankle *reflexes*.

Pathologic edema is often 3+ or more pretibially.

After 24 weeks, reflexes greater than 2+ may indicate pregnancy-induced hypertension.

SPECIAL TECHNIQUES

LEOPOLD'S MANEUVERS

Leopold's maneuvers help determine:

 Fetal lie, or where the fetus lies in relation to the woman's back (longitudinal or transverse) Common deviations include breech presentation (fetal buttocks present at the outlet of the maternal pelvis) and absence of the presenting part well down into the maternal pelvis at term.

POSSIBLE FINDINGS

- Presentation, or the end of the fetus that is presenting at the pelvic inlet (head or buttocks)
- Location of the fetal back
- Engagement, or how far the presenting part of the fetus has descended into the maternal pelvis
- Estimated fetal weight

First Maneuver (Upper Pole)

Stand at the woman's side, facing her head. Keep the fingers of both examining hands together. Palpate gently with the fingertips to determine what part of the fetus is in the upper pole of the uterine fundus.



Second Maneuver (Sides of the Maternal Abdomen)

Place one hand on each side of the woman's abdomen, aiming to capture the body of the fetus between them. Use one hand to steady the uterus and the other to palpate the fetus.



Third Maneuver (Lower Pole)

Face the woman's feet. Palpate the area just above the symphysis pubis. Note whether the hands diverge with downward pressure or stay together to learn if the presenting part of the fetus, head or buttocks, is descending into the pelvic inlet.



Fourth Maneuver (Confirmation of the Presenting Part)

With your dominant hand, grasp the part of the fetus in the lower pole, and with your nondominant hand, the part of the fetus in the upper pole. Try to distinguish between the head and the buttocks.



RECORDING YOUR FINDINGS

Recording the Physical Examination— The Pregnant Woman

"Abdomen: No surgical scars. Active bowel sounds. Soft, nontender; no palpable hepatosplenomegaly or masses. Fundus palpable 2 fingerbreadths below the umbilicus; shape is ovoid and smooth. Fetal heart rate 144. No inguinal adenopathy. External genitalia: midline episiotomy scar present. No lesions, discharge, or signs of infection. Bimanual examination: cervix midline, soft, 4 cm in length; external os admits fingertip, internal os closed. No pain elicited on movement of cervix; no adnexal masses. Fundus enlarged to 20 weeks' size, midline, smooth; vaginal tone reduced." (Describes examination of healthy pregnant woman at 20 weeks' gestation)

20

The Older Adult

Older adults now number more than 27 million in the United States, growing to 86 million by 2050. Life span at birth is currently 79 years for women and 74 years for men. The "demographic imperative" is to maximize not only life span but also "health span" of our older population, so that seniors maintain full function for as long as possible, enjoying rich and active lives in their homes and communities.

- Assessing the older adult entails a focus on healthy or "successful" aging; the need to understand and mobilize family, social, and community supports; the importance of skills directed to functional assessment, "the sixth vital sign"; and promoting the older adult's long-term health and safety.
- The aging population displays marked heterogeneity. Investigators distinguish "usual" aging, with its complex of diseases and impairments, from optimal aging. Optimal aging occurs in those people who escape debilitating disease entirely and maintain healthy lives late into their 80s and 90s. Studies of centenarians show that genes account for approximately 20% of the probability of living to 100, with healthy lifestyles accounting for approximately 20% to 30%.

THE HEALTH HISTORY



APPROACH TO THE PATIENT

As you talk with older adults, convey respect, patience, and cultural awareness. Be sure to address patients by their last name.

Adjusting the Office Environment. Make sure the office is neither too cool nor too warm. Face the patient directly, sitting at eye level. A well-lit room allows the older adult to see your facial expressions and gestures.

More than 50% of older adults have hearing deficits. Free the room of distractions or noise. Consider using a "pocket talker," a microphone that amplifies your voice and connects to an earpiece inserted by the patient. Chairs with higher seating and a wide stool with a handrail leading up to the examining table help patients with quadriceps weakness.

Shaping the Content and Pace of the Visit. Older people often reminisce. Listen to this process of life review to gain important insights and help patients as they work through painful feelings or recapture joys and accomplishments.

Balance the need to assess complex problems with the patient's endurance and possible fatigue. Consider dividing the initial assessment into two visits.

Eliciting Symptoms in the Older Adult. Older patients may overestimate healthiness even when increasing disease and disability are obvious. To reduce the risk of late recognition and delayed intervention, adopt more directed questions or *health screening tools*. Consult with family members and caretakers

Acute illnesses present differently in older adults than in younger age groups. Be sensitive to changes in presentation of myocardial infarction and thyroid disease. Older patients with infections are less likely to have fever.

Recognize the symptom clusters typical of different *geriatric syndromes*, notable for the interaction and probable synergism among multiple risk factors, for example, falls, dizziness, depression, urinary incontinence, and functional impairment. Searching for the usual "unifying diagnosis" may pertain to fewer than 50% of older adults

Cognitive impairment may affect the patient's history. Even elders with mild cognitive impairment, however, can provide

sufficient history to reveal concurrent disorders. Use simple sentences with prompts to trigger necessary information. If impairments are more severe, confirm symptoms with family members or caretakers.

Addressing Cultural Dimensions of Aging. By 2050, the older adult population will increase by 230%, and the minority older adult population by 510%. Cultural differences affect the epidemiology of illness and mental health, acculturation, the specific concerns of the elderly, the potential for misdiagnosis, and disparities in health outcomes. Review the components of self-awareness needed for cultural responsiveness, discussed in Chapter 3 (pp. 52–53). Ask about spiritual advisors and native healers. Cultural values particularly affect decisions about the end of life. Elders, family, and even an extended community group may make these decisions with or for the older patient.



₹ COMMON CONCERNS

- Activities of daily living
- Instrumental activities of daily living
- Medications
- Nutrition
- Acute and persistent pain
- Smoking and alcohol
- Advance directives and palliative care

Place symptoms in the context of your overall *functional assessment*, always focusing on helping the older adult to maintain optimal well-being and level of function.

Activities of Daily Living. Daily activities provide an important baseline for the future. You might say "Tell me about your typical day" or "Tell me about your day yesterday." Then move to a greater level of detail: "You got up at 8 AM? How is it getting out of bed?"

ACTIVITIES OF DAILY LIVING AND INSTRUMENTAL ACTIVITIES OF DAILY LIVING	
Physical Activities of	Instrumental Activities of
Daily Living (ADLs)	Daily Living (IADLs)
Bathing	Using the telephone
Dressing	Shopping
Toileting	Preparing food
Transferring	Housekeeping
Continence	Laundry
Feeding	Transportation
Managing money	Taking medicine

Medications. Adults older than 65 take approximately 30% of all prescriptions. Roughly 30% take more than eight prescribed drugs each day! Take a thorough medication history, including name, dose, frequency, and indication for each drug. Explore all components of polypharmacy, or suboptimal prescribing, including concurrent use of multiple drugs, underuse, inappropriate use, and nonadherence. Ask about use of over-the-counter medications, vitamin and nutrition supplements, and mood-altering drugs. Medications are the most common modifiable risk factor associated with falls.

Nutrition. Taking a diet history and using the Rapid Screen for Dietary Intake and the Nutrition Screening Checklist (pp. 71–72) are especially important in older adults.

Acute and Persistent Pain. Pain and associated complaints account for 80% of clinician visits, usually for musculoskeletal complaints like back and joint pain. Older patients are less likely to report pain, leading to undue suffering, depression, social isolation, physical disability, and loss of function.

Inquire about pain each time you meet with the older patient. Ask specifically, "Are you having any pain right now? How about over the past week?" Unidimensional scales such as the Visual Analog Scale, graphic pictures, and the Verbal 0–10 Scale have all been validated and are easiest to use.

Characteristics of Acute and Persistent Pain		
Acute Pain	Persistent Pain	
Distinct onset	Lasts more than 3 months	
Obvious pathology	Often associated with psychological or functional impairment	
Short duration	Can fluctuate in character and intensity over time	
Common causes: postsurgical, trauma, headache	Common causes: arthritis, cancer, claudication, leg cramps, neuropathy, radiculopathy	

(Source: Reuben DB, Herr KA, Pacala JT, et al. Geriatrics at Your Fingertips: 2004, 6th ed. Malden, MA: Blackwell Publishing, for the American Geriatrics Society, 2004:149.)

Smoking and Alcohol. At each visit, advise elderly smokers to quit.

An estimated 5% to 10% of older adults have alcohol-related problems, and this percentage is expected to rise as the population ages in coming decades. Despite the prevalence of alcohol problems among the elderly, rates of detection and treatment are low. Use the CAGE questions to uncover problem drinking (see p. 50).

Advance Directives and Palliative Care. Providers should initiate these discussions before serious illness develops. Advance care planning involves providing information, invoking the patient's preferences, identifying proxy decision makers, and conveying empathy and support. Use clear, simple language. Ask about preferences relating to written "Do Not Resuscitate" orders specifying life support measures "if the heart or lungs were to stop or give out." Seek a written health care proxy or durable power of attorney for health care, "someone who can make decisions reflecting your wishes in case of confusion or emergency." Include these discussions in office settings rather than the uncertain and stressful environment of emergency or acute care.

The goal of *palliative care* is "to relieve suffering and improve the quality of life for patients with advanced illnesses and their families through specific knowledge and skills, including communication with patients and family members; management of pain and other symptoms; psychosocial, spiritual, and bereavement support; and coordination of an array of medical and social services."

HEALTH PROMOTION AND COUNSELING

Important Topics for Health Promotion and Counseling in the Older Adult

- When to screen
- Cancer screening
- Dementia and mild cognitive impairment
- Elder mistreatment

When to Screen. As the life span for older adults extends into the 80s, new issues for screening emerge. In general, base screening decisions on each older person's particular circumstances, rather than on age alone. Consider life expectancy, time interval until benefit from screening accrues, and patient preference. The American Geriatrics Society recommends that if life expectancy is short, give priority to treating conditions that will benefit the patient in the time that remains.

- Screen for age-related changes in *vision* and *hearing*. These are included in the 10-Minute Geriatric Screener (pp. 411–412).
- Recommend regular aerobic exercise. Resistance training and Tai Chi may help to improve balance.
- Immunizations: include the pneumococcal vaccine once after age 65, annual influenza vaccinations, and the herpes zoster vaccine.
- Promote household safety. Correct poor lighting, chairs at awkward heights, slippery or irregular surfaces, and environmental hazards.

Cancer Screening. Cancer screening can be controversial because of limited evidence about adults older than age 70 to 80:

 The American Geriatrics Society recommends annual or biennial mammography for breast cancer screening up to

- age 75, then every 2 to 3 years if life expectancy remains more than 4 years.
- 40% to 50% of deaths from cervical cancer are in women older than 65 years. Provide Pap smears every 1 to 3 years until age 65 to 70 when there is no history of cervical pathology.
- Colonoscopy is recommended for colon cancer screening every 10 years, beginning at age 50. This examination is difficult for many older patients, who may decline despite encouragement.
- Check for skin and oral cancers in high-risk patients.

Dementia and Mild Cognitive Impairment. Dementia is "an acquired syndrome of decline in memory and at least one other cognitive domain such as language, visuospatial, or executive function sufficient to interfere with social or occupational functioning." It affects 16% of Americans over age 65. Prominent features include:

- Short- and long-term memory deficits and impaired judgment
- Impoverished thought processes
- Hesitant speech resulting from difficulty finding words
- Loss of orientation to place

Most dementias represent Alzheimer's disease (50% to 85%) or vascular multi-infarct dementia (10% to 20%). Dementia often has a slow, insidious onset. It may escape detection by both families and clinicians, especially in the early stages of mild cognitive impairment, which may be detected only on neurocognitive testing. Look for problems with memory, then later for changes in cognitive function or ADLs. Watch for family complaints of new or unusual behaviors. Investigate contributing factors such as medications, depression, metabolic abnormalities, or other medical and psychiatric conditions.

Elder Mistreatment. Screen older patients for possible elder mistreatment, which includes abuse, neglect, exploitation, and abandonment. Prevalence is ~1% to 5% of the older population; however, many more cases may remain undetected.

TECHNIQUES OF EXAMINATION

Assessment of the older adult does not follow the traditional format of the history and physical examination. Enhanced techniques of interviewing, special emphasis on daily function and key topics related to elder health, and a focus on functional assessment are especially important.

Functional status is the ability to perform tasks and fulfill social roles associated with daily living across a wide range of complexity.



Assessing Functional Ability. Several performance-based assessment instruments are available. The screening tool below is brief, has high inter-rater agreement, and can be used easily by office staff. It covers the three important domains of geriatric assessment: physical, cognitive, and psychosocial function. It addresses key sensory modalities and urinary incontinence, an often unreported problem that greatly affects social interactions and self-esteem in the elderly. One mnemonic that helps students assess incontinence is DIAPERS: Delirium, Infection, Atrophic urethritis/vaginitis, Pharmaceuticals, Excess urine output (e.g., due to congestive heart failure, hyperglycemia), Restricted mobility, Stool impaction.

10-Minute Geriatric Screener	
Problem and Screening Measure	Positive Screen
Vision: 2 Parts: Ask: "Do you have difficulty driving, or watching television, or reading, or doing any of your daily activities because of your eyesight?	Yes to question and inability to read greater than 20/40 on Snellen chart
	(continued)


10-Minute Geriatric Screener (continued)	
Problem and Screening Measure	Positive Screen
If yes, then: Test each eye with Snellen chart while patient wears corrective lenses (if applicable).	
Hearing: Use audioscope set at 40 dB. Test hearing using 1,000 and 2,000 Hz.	Inability to hear 1,000 or 2,000 Hz in both ears or either of these frequencies in one ear
Leg mobility: Time the patient after instructing: "Rise from the chair. Walk 20 feet briskly, turn, walk back to the chair, and sit down."	Unable to complete task in 15 seconds
Urinary incontinence: 2 Parts: Ask: "In the last year, have you ever lost your urine and gotten wet?"	Yes to both questions
If yes, then ask: "Have you lost urine on at least 6 separate dates?"	
Nutrition/weight loss: 2 parts: Ask: "Have you lost 10 lbs over the past 6 months without trying to do so?"	Yes to the question or weight <100 lbs
Weigh the patient.	
Memory: Three-item recall	Unable to remember all three items after 1 minute
Depression: Ask: "Do you often feel sad or depressed?"	Yes to the question
Physical disability: Six questions: "Are you able to :	No to any of the questions
"Do strenuous activities like fast walking or bicycling?"	
	(continued)

Problem and Screening Measure	Positive Screen
"Do heavy work around the house	
like washing windows, walls, or	
floors?"	
"Go shopping for groceries or	
clothes?"	
"Get to places out of walking	
distance?"	
"Bathe, either a sponge bath, tub	
bath, or shower?"	
"Dress, like putting on a shirt,	
buttoning and zipping, or putting	
on shoes?"	

(Source: More AA, Siu AL. Screening for common problems in ambulatory elderly: clinical confirmation of a screening instrument. Am J Med 100:438–440, 1996.)

Further Assessment of Falls. Each year ~35% to 40% of healthy community-dwelling older adults experience falls. Incidence rates in nursing homes and hospitals are almost three times higher, with related injuries in approximately 25%.

The American Geriatrics Society recommends risk factor assessment for falls during routine primary care visits, with more intensive assessment in high-risk groups—those with first or recurrent falls, nursing home residents, and those prone to fall-related injuries. Assess how the fall occurred, seeking details from any witnesses, and identify risk factors, medical comorbidities, functional status, and environmental risks. Couple your assessment with interventions for prevention, including gait and balance training and exercise to strengthen muscles, reduction of home hazards, discontinuation of psychotropic medication, and multifactorial assessment with targeted interventions. Recommendations from the American Geriatrics Society are provided on the next page.



(Source: American Geriatrics Society, British Geriatrics Society, American Academy of Orthopaedic Surgeons. Guideline for the prevention of falls in older persons. Am Geriatr Soc 49(5):664-672, 2001.)

EXAMINATION TECHNIQUES

PHYSICAL EXAMINATION OF THE OLDER ADULT

Vital Sians. Measure blood pressure, checking for increased systolic blood pressure (SBP) and widened

Isolated systolic hypertension (SBP ≥140) after age 50 triples the risk of coronary heart disease in men. PP ≥60 is a risk

POSSIBLE FINDINGS

pulse pressure (PP), defined as SBP minus diastolic blood pressure (DBP).

factor for cardiovascular and renal disease and stroke.

Assess the patient for orthostatic hypotension, defined as a drop in SBP of ≥20 mm Hg or DBP of ≥10 mm Hg within 3 minutes of standing. Measure in two positions: supine after the patient rests for up to 10 minutes, then within 2 to 3 minutes after standing up.

Orthostatic hypotension in 10%–20% of older adults and in up to 30% of frail nursing home residents, especially when they first arise in the morning. Watch for lightheadedness, weakness, unsteadiness, visual blurring, and, in 20% to 30% of patients, syncope.

Review the JNC VII categories of "prehypertension" to help with early detection and treatment. Assess for medications, autonomic disorders, diabetes, prolonged bedrest, blood loss, and cardiovascular disorders.

Measure heart rate, respiratory rate, and temperature. The apical heart rate may yield more information about arrhythmias in older patients. Use thermometers accurate for lower temperatures.

Respiratory rate ≥25 breaths per minute indicates lower respiratory infection.

Weight and height are especially important and needed for calculation of the body mass index (p. 60). Weight should be measured at every visit.

Hypothermia is more common in elderly patients.

Low weight is a key indicator of poor nutrition.

Undernutrition in depression, alcoholism, cognitive impairment, malignancy, chronic organ failure (cardiac, renal, pulmonary), medication use, social isolation, and poverty

Skin. Note physiologic changes of aging, such as thinning, loss of elastic tissue and turgor, and wrinkling.

Check the extensor surface of the hands and forearms.

Look for changes from sun exposure. There may be actinic lentigines, or "liver spots," and actinic keratoses, superficial flattened papules covered by a dry scale (p. 104).

Inspect for the benign comedones, or blackheads, on the cheeks or around the eyes; cherry angiomas (p. 103); and seborrheic keratoses, (p. 104).

Inspect for painful vesicular lesions in a dermatomal distribution.

In older bedbound patients, especially when emaciated or neurologically impaired, inspect for damage or ulceration.

HEENT. Inspect the eyelids, the bony orbit, and the eye.

Dry, flaky, rough, and often itchy

White depigmented patches (pseudoscars); welldemarcated, vividly purple macules or patches that may fade after several weeks (actinic purpura)

Distinguish such lesions from a basal cell carcinoma and squamous cell carcinoma (p. 105). Dark, raised, asymmetric lesion with irregular borders in melanoma

Herpes zoster from reactivation of latent varicella-zoster virus in the dorsal root ganglia

Pressure sores if obliteration of arteriolar and capillary blood flow to the skin or shear forces with movement across sheets or lifting upright incorrectly

Senile ptosis arising from weakening of the levator

POSSIBLE FINIDINGS

palpebrae, relaxation of the skin, and increased weight of the upper eyelid

Ectropion or entropion of lower lids (p. 130)

Yellowing of the sclera and arcus senilis, a benign whitish ring around the limbus

Test visual acuity, using a pocket Snellen chart or wall-mounted chart.

Examine the lenses and fundi

Inspect each lens for opacities.

Assess the cup-to-disc ratio, usually $\leq 1:2$.

Inspect the fundi for colloid bodies causing alterations in pigmentation called *drusen*. These may be hard and sharply defined, or soft and confluent with altered pigmentation.

Test hearing by the whispered voice (see p. 121) or audioscope. Inspect ear canals for cerumen.

More than 40 million Americans have refractive errors—presbyopia.

Cataracts, glaucoma, and macular degeneration all increase with aging.

Cataracts are the world's leading cause of blindness.

Increased cup-to-disc ratio suggests open-angle *glaucoma* and possible loss of peripheral and central vision, and blindness. Prevalence is three to four times higher in African-Americans.

Macular degeneration causes poor central vision and blindness. Types include dry atrophic (more common but less severe) and wet exudative, or neovascular.

Removing cerumen often quickly improves hearing.

Examine the oral cavity for odor, appearance of the gingival mucosa, any caries, mobility of the teeth, and quantity of saliva.

Inspect for lesions on mucosal surfaces. Ask patient to remove dentures so you can check gums for denture sores.

Thorax and Lungs. Note subtle signs of changes in pulmonary function.

Cardiovascular System. Review blood pressure and heart rate.

Inspect the jugular venous pulsation (JVP), palpating the carotid upstrokes, and listen for any overlying carotid bruits.

Assess the point of maximal impulse (PMI), and then heart sounds.

Malodor in poor oral hygiene, periodontitis, or caries

Gingivitis if periodontal disease

Dental plaque and cavitation if caries. Increased tooth mobility; risk of tooth aspiration

Decreased salivation from medications, radiation, Sjögren's syndrome, or dehydration

Oral tumors, usually on lateral borders of tongue and floor of mouth

Increased anteroposterior diameter, purse-lipped breathing, and dyspnea with talking or minimal exertion in chronic obstructive pulmonary disease

Isolated systolic hypertension and a widened pulse pressure are cardiac risk factors. Search for left ventricular hypertrophy (LVH).

A tortuous atherosclerotic aorta can raise pressure in the left jugular veins by impairing drainage into right atrium.

Carotid bruits in possible carotid stenosis.

Sustained PMI is found in LVH: diffuse PMI is found within congestive heart failure (see p. 170).

POSSIBLE FINDINGS

Listen for cardiac murmurs in all areas (see p. 172). Describe timing, shape, location of maximal intensity, radiation, intensity, pitch, and quality of each murmur.

In older adults, S₃ in dilatation of the left ventricle from congestive heart failure or cardiomyopathy; S₄ in hypertension

A systolic crescendo decrescendo murmur in the second right interspace in aortic sclerosis or aortic stenosis. Both carry increased risk of cardiovascular disease and death.

A harsh holosystolic murmur at the apex suggests *mitral* regurgitation, also common in the elderly.

For systolic murmurs over the clavicle, check for delay between the brachial and radial pulses. Delay during simultaneous palpation (but not compression) of brachial and radial pulses in appring stemps is

Breasts and Axillae. Palpate the breasts carefully for lumps or masses.

Possible breast cancer

Abdomen.

- Check for any bruits over the aorta, renal arteries, and femoral arteries.
- *Bruits* in atherosclerotic vascular disease
- Inspect the upper abdomen; palpate to the left of the midline for aortic pulsations.

Widened aorta and pulsatile mass may be found in abdominal aortic aneurysm.

Female Genitalia and Pelvic Examination.

Take special care to explain the steps of the examination and allow time for careful positioning. For the woman with arthritis or spinal deformities who cannot flex her hips or knees, an assistant can gently raise and support the legs, or help the woman into the left lateral position.

Inspect the vulva for changes related to menopause; identify any labial masses. Bluish swellings may be varicosities. Benign masses include condylomata, fibromas, leiomyomas, and sebaceous cysts.

Bulging of the anterior vaginal wall below the urethra in urethrocele

Inspect the urethra for *caruncles*, or prolapse of fleshy erythematous mucosal tissue at the urethral meatus.

Clitoral enlargement in androgen-producing tumors or use of androgen creams

Speculum Examination.Inspect vaginal walls, which may be atrophic, and cervix.

Estrogen-stimulated cervical mucus with ferning in use of hormone replacement therapy, endometrial hyperplasia, and estrogen-producing tumors

Obtain endocervical cells for the Pap smear. Use a blind swab if the atrophic vagina is too small.

Removing speculum, ask patient to bear down.

Perform the bimanual examination.

Uterine prolapse, cystocele, urethrocele, or rectocele.

See Table 14-6, Positions of the Uterus, and Uterine Myomas, p. 257.

POSSIBLE FINDINGS

Mobility of cervix restricted if inflammation, malignancy, or surgical adhesion

Palpable ovaries in ovarian cancer.

Perform the rectovaginal examination.

Enlarged, fixed, or irregular uterus if adhesions or malignancy. Rectal masses in colon cancer.

Male Genitalia and Prostate. Examine the penis; retract foreskin if present. Examine the scrotum, testes, and epididymis.

Smegma, penile cancer, and scrotal hydroceles

Do a rectal examination.

Rectal masses in colon cancer. Prostate hyperplasia if enlargement; prostate cancer if nodules or masses.

Peripheral Vascular System.

Auscultate the abdomen for aortic, renal, femoral artery bruits.

Bruits over these vessels in atherosclerotic disease.

Palpate pulses.

Diminished or absent pulses in *arterial occlusion*. Confirm with an office ankle–brachial index.

Musculoskeletal System.

Screen general range of motion and gait. If joint deformity, deficits in mobility, or pain with Review the techniques for examining individual joints in Chapter 16.

Degenerative joint changes in *osteoarthritis;* joint

movement, conduct a more thorough examination.

inflammation in rheumatoid or aouty arthritis. See Tables 16-1 to 16-4.

Nervous System. Begin evaluation with the 10-Minute Geriatric Screener, pp. 411-412.

Pursue further examination if any deficits. Focus especially on memory and affect.

Assess gait and balance, particularly standing balance: timed 8-foot walk: stride characteristics like width, pace, and length of stride; and careful turning.

Although neurologic abnormalities are common in older adults, their prevalence without identifiable disease increases with age, ranging from 30% to 50%.

Assess any tremor, rigidity, bradykinesia, micrographia, shuffling gait, and difficulty turning in bed, opening jars, and rising from a chair.

Learn to distinguish delirium from depression and dementia. See Table 20-1.

Abnormalities of gait and balance, especially widening of base, slowing and lengthening of stride, and difficulty turning, are correlated with risk of falls.

Physiologic changes of aging: unequal pupil size, decreased arm swing and spontaneous movements, increased leg rigidity and abnormal gait, presence of the snout and grasp reflexes, and decreased toe vibratory sense.

Tremor is slow frequency and at rest, with a "pill-rolling" quality, and aggravated by stress and inhibited during sleep or movement.

Essential tremor if bilateral, symmetric with positive family history, and diminished by alcohol

RECORDING YOUR FINDINGS

As you read through this physical examination, you will notice some atypical findings. Try to test yourself. See if you can interpret these findings in the context of all you have learned about the examination of the older adult.

Recording the Physical Examination—The Older Adult

2/1/08

Mr. J is an older adult who appears healthy but underweight, with good muscle bulk. He is alert and interactive, with good recall of his life history. He is accompanied by his son.

Vital Signs: Ht (without shoes) 160 cm (5'). Wt (dressed) 65 kg (143 lb). BMI 28. BP 145/88 right arm, supine; 154/94 left arm, supine. Heart rate (HR) 98 and regular. Respiratory rate (RR) 18. Temperature (oral) 98.6°F.

10-Minute Geriatric Screener: (see pp. 411-412)

Vision: Patient reports difficulty reading. Visual acuity 20/60 on Snellen chart.

Needs further evaluation for glasses and possibly hearing aid.

Hearing: Cannot hear whispered voice in either ear. Cannot hear 1,000 or 2,000 Hz with audioscope in either ear.

Leg Mobility: Can walk 20 feet briskly, turn, walk back to chair, and sit down in 14 seconds.

Urinary Incontinence: Has lost urine and gotten wet on 20 separate days.

Needs further evaluation for incontinence, including "DIAPER" assessment (see p. 410), prostate examination, and postvoid residual, which is normally ≤50 mL (requires bladder catheterization).

Nutrition: Has lost 15 lbs over the past 6 months without trying.

(continued)

Needs nutritional screen (see pp. 71–72).

Memory: Can remember three items after 1 minute.

Depression: Does not often feel sad or depressed.

Physical Disability: Can walk fast but cannot ride a bicycle. Can do moderate but not heavy work around the house. Can go shopping for groceries or clothes. Can get to places out of walking distance. Can bathe each day without difficulty. Can dress, including buttoning and zipping, and can put on shoes.

Consider exercise regimen with strength training.

Physical Examination: Record the vital signs and weight. Carefully describe your findings for each relevant segment of the peripheral examination, using terminology found in the "Recording Your Findings" section of the prior chapters.

AIDS TO INTERPRETATION

TABLE 20-1 Delirium and Dementia

	Delirium	Dementia
Clinical Features		
Onset	Acute	Insidious
Course	Fluctuating, with lucid intervals; worse at night	Slowly progressive
Duration	Hours to weeks	Months to years
Sleep/Wake Cycle	Always disrupted	Sleep fragmented
General Medical Illness or Drug Toxicity	Either or both present	Often absent, especially in Alzheimer's disease
Mental Status		
Level of Consciousness	Disturbed. Person less clearly aware of the environ- ment and less able to focus, sustain, or shift attention	Usually normal until late in the course of the illness
Behavior	Activity often abnormally decreased (somnolence) or increased (agitation, hypervigilance)	Normal to slow; may become inappropriate
Speech	May be hesitant, slow or rapid, incoherent	Difficulty in finding words, aphasia
Mood	Fluctuating, labile, from fearful or irritable to normal or depressed	Often flat, depressed

(table continues next page)

TABLE 20-1 Delirium and Dementia (continued)

	Delirium	Dementia
Thought Processes	Disorganized, may be incoherent	Impoverished. Speech gives little information.
Thought Content	Delusions common, often transient	Delusions may occur.
Perceptions	Illusions, hallucinations, most often visual	Hallucinations may occur.
Judgment	Impaired, often to a varying degree	Increasingly impaired over the course of the illness
Orientation	Usually disoriented, especially for time. A known place may seem unfamiliar.	Fairly well maintained, but becomes impaired in the later stages of illness
Attention	Fluctuates. Person easily distracted, unable to concentrate on selected tasks	Usually unaffected until late in the illness
Memory	Immediate and recent memory impaired	Recent memory and new learning especially impaired
Examples of Cause	Delirium tremens (due to withdrawal from alcohol) Uremia	Reversible: Vitamin B ₁₂ deficiency, thyroid disorders
		Irreversible:
	Acute hepatic failure	Alzheimer's
	Acute cerebral vasculitis	disease, vascular dementia (from
	Atropine poisoning	multiple infarcts), dementia due to head trauma

TABLE 20-2

Screening for Dementia: The Mini-Cog

Administration

The test is administered as follows:

- 1. Instruct the patient to listen carefully to and remember 3 unrelated words and then to repeat the words.
- 2. Instruct the patient to draw the face of a clock, either on a blank sheet of paper or on a sheet with the clock circle already drawn on the page. After the patient puts the numbers on the clock face, ask him or her to draw the hands of the clock to read a specific time.
- 3. Ask the patient to repeat the 3 previously stated words.

Scoring


Give 1 point for each recalled word after the clock drawing test (CDC) distractor.

Patients recalling none of the three words are classified as demented (Score = 0).

Patients recalling all three words are classified as nondemented (Score = 3).

Patients with intermediate word recall of 1-2 words are classified based on the CDT (Abnormal = demented; Normal = nondemented).

Note: The CDT is considered normal if all numbers are present in the correct sequence and position, and the hands readably display the requested time.



MINI-COG SCORING ALGORITHM

(From Borson S, Scanlan J, Brush M, et al. The Mini-Cog: a cognitive 'vital signs' measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry 15*(11);1021–1027, 2000. Copyright John Wiley & Sons Limited. Reproduced with permission.)

INDEX

NOTE: Page numbers followed by b indicate boxed material; those followed by t indicate tables.

A	Acromioclavicular arthritis, 307t
ABCD screen, 93, 106t	Actinic keratoses, 104t, 415
Abdomen, 193–213. See also specific	Actinic lentigines, 415
structures	Actinic purpura, 415
back pain referred from, 302t	Activities of daily living, 405, 406b
in children, 364, 372	Adams bend test, 375
common or concerning symptoms	Adduction stress test, 294
of, 193 <i>b</i>	Adenomas, breast, 182b
disorders of, 209 <i>t</i> –213 <i>t</i>	Adnexa (ovaries), 15, 249, 400
examination of, 14, 201-208	Adolescents. See also Children
fullness in, 195	examining, 352, 374-375
health history of, 193-199	hypertension in, 357b, 380t
health promotion or counseling	substance use in, 51
for, 199–200	Advance directives, 407
in older adults, 418	Adventitious breath sounds, 146,
painful or tender, 193-196, 198,	147 <i>b</i> , 150, 161 <i>t</i>
213 <i>t</i>	Affect, 79 <i>b</i> , 81
during pregnancy, 396-398	Age
recording exam findings of, 208b,	gestational, 353, 354b
377 b	of patient, 28
Abdominal aortic aneurysm,	Agenda, for interview, 39–40
217–218, 418	Agoraphobia, 90 <i>t</i>
Abdominal reflexes, 330, 340t	AIDS/HIV infection, 105 <i>t</i> , 231–232
Abducens nerve, 317b, 318	Air and bone conduction test, 122,
Abduction stress test, 296	135 <i>t</i> , 320
Abrasions, genital, 373	Airway sounds. See Breath sounds
Abscess	Alcohol use and abuse
brain, 112	health promotion and counseling
lung, 157t	for, 80, 199
peritonsillar, 371	history of, 2 <i>b</i> , 4, 50–51
Abstract thinking, 84	liver disease in, 193, 197b
Abuse	in older adults, 407
of alcohol, 2 <i>b</i> , 4, 50–51, 199, 407	screening for, 80
of drugs, 4, 50–51, 80	Alertness, patient, 77, 81, 333b
of older adults, 409	Alkalosis, respiratory, 396
physical, 51, 392–393	Allen test, 222
sexual, 373, 388 <i>b</i>	Allergies, 2b, 4, 114, 350
Achilles tendon, 297	Alopecia areata, 107t
Acoustic nerve, 317 <i>b</i> , 319–320	Alternative health care, 5
Acquired immunodeficiency syndrome	Alzheimer's disease, 409
(AIDS), 105t. See also HIV	Amblyopia, 368
Acrocyanosis, 358	Amenorrhea, 242

Anal reflex, 331	Arcus senilis, 416
Analgesic rebound headaches, 126t	Argyll Robertson pupils, 117
"Anatomic snuffbox," 284	Arms
Androgen-producing tumors, 419	examination of, 218-219
Anemia of pregnancy, 395	flaccid, 334
Aneurysm, aortic, 152t, 163, 418	pain in, 215
Angina pectoris, 152t, 163	Arousal (reticular activating) system,
Angioedema, 136t	312
Angioma, 103t, 415	Arrhythmias, 64, 176 <i>t</i> –180 <i>t</i> , 364.
Angry patient, 47	See also specific arrhythmias
Angular cheilitis, 136t	Arterial insufficiency, 223, 224t, 225t
Anhedonia, 80	Arterial occlusion, 219
Ankle-brachial index, 217, 226t–227t	Arterial pulses, 218b
Ankles, 297–298	Arteries, retinal, 119, 120 <i>b</i>
muscles of, 323	Arthralgia, 275
reflexes of, 329, 400	Arthritis
ulcers on, 216, 225 <i>t</i>	acromioclavicular, 307t
Ankylosing spondylitis, 302t	in ankles or feet, 297, 298
Anorexia, 60, 70 <i>t</i> , 196, 242	degenerative, 275, 283, 305 <i>t</i> –306 <i>t</i> ,
Anteflexion, 257 <i>t</i>	308t, 421
Anterior cruciate ligament (ACL),	
	gonococcal, 274, 283
294–295, 309 <i>t</i>	gouty, 421
Anterior drawer sign, 294, 309 <i>t</i>	in hips, 290, 291
Anus, 263, 264, 265 <i>b</i> , 398	in knees, 293, 308 <i>t</i>
Anxiety, 57, 90 <i>t</i> –91 <i>t</i> , 153 <i>t</i>	in older adults, 420–421
Aorta	patterns of pain in, 305 <i>t</i> –306 <i>t</i>
aneurysm of, 141, 152 <i>t</i> , 163,	rheumatoid, 274–275, 276, 283,
217–218	284, 305 <i>t</i> –306 <i>t</i> , 308 <i>t</i> , 421
coarctation of, 363	septic, 275
examination of, 14, 205	in shoulders, 307t
Aortic valve	in spine, 289
regurgitation of (insufficiency),	in wrists or hands, 283, 284
169, 171, 173, 219	Articular structures, 271, 275
sclerosis of, 418	Asbestosis, 155 <i>t</i>
stenosis of, 169, 170, 174, 175,	Ascites, 206–207
381 <i>t</i> , 418	Assessments, 17–36. See also specific
Apgar score, 352–353	assessments
Aphasia, 82	case study of, 19 <i>b</i> –28 <i>b</i>
Aphthous ulcer, 138t	comprehensive vs. focused, 1b, 37
Apical pulse (impulse)	organizing data for, 28-30
in children, 363–364	and problem list, 36
measuring, 64, 168, 170	process of, 17-19
in older adults, 417	recording, 33-36, 69
in pregnancy, 396	test selection in, 30b-32b
Apnea, 361	tips for, 29b
Appearance	Asterixis, 332
of clinician, 38	Asthma, 154t, 158t, 161t
of patient, 58, 78, 81	Astigmatism, 126t
Appendicitis, 194, 196, 207–208	Ataxia, 313
' '	

Ataxic (Biot's) breathing, 74t	Biceps reflexes, 328
Atelectasis, 161t	Bicipital tendonitis, 281, 307t
Atherosclerosis, 420	Biliary obstruction, 198b
Athetosis, 341t	Bimanual examination, 15, 249
Atrial fibrillation, 64, 168	in older adults, 419-420
Atrial septal defect, 364	in pregnancy, 399-400
Attention, 77, 78, 83, 425t	Biot's (ataxic) breathing, 74t
Auditory acuity. See Hearing; Hear-	Birth control, 245
ing loss	Birth history, 350
Auricles, 120–121, 360	Birth weight, 353, 354 <i>b</i>
Auscultation	Bladder disorders, 198-199,
of abdomen, 201-202	211 <i>t</i> –212 <i>t</i> , 274
of chest, 146-148, 150	Bleeding
of fetal heart, 397-398	gastrointestinal, 260
of heart, 171-173, 364, 396	nose, 114, 395
Auscultatory gap, 62b, 168	postmenopausal, 243
Autonomy, 54	rectal, 260
Axillae, 12, 186, 418	subarachnoid, 112, 127t, 313
Axillary temperature, 65	vaginal, 373
	vitreous, 112
	Blindness, 129t
В	Bloating, abdominal, 195
Babinski response, 328, 330	Blood
Back	in stool, 260
examination of, 12, 287-289	in urine, 198
pain in, 274, 276, 301 t-302 t	Blood pressure. See also Hypertension;
Bacterial pneumonia, 156t	Hypotension, orthostatic
Bacterial vaginosis, 255t	adult classifications of, 63b, 164
"Baker's" (popliteal) cyst, 293, 310t	in children and adolescents, 357,
Balance, 313, 421	368, 380t
Balanitis, 232	cuffs for measuring, 61b, 368
Balloon sign, 294	and diet, 73t
Ballottement	health promotion for, 164
abdominal, 207	measuring, 61-64, 168
patellar, 295	in older adults, 413-414
Barlow test, 365	in pregnancy, 394, 395b, 400
Barrel chest, 159t	Body mass index (BMI)
Bartholin's glands, 247, 398	calculating, 59-61
Basal cell carcinoma, 105t, 415	of children, 368
Basal ganglia, 311, 340t	excessively low, 70t
Behavior. See also Mental status	of older adults, 414
of clinician, 38	of pregnant women, 395
in delirium and dementia, 424t	Body odors, 58
of patient, 46-47, 81	Body position
Bell's palsy, 339t	observation of, 321
Beneficence, 54	for physical exam, 9b, 11, 394b, 396
Benign prostatic hyperplasia, 260,	Bone density, 277
264, 267 <i>t</i> , 268 <i>t</i> , 420	Bottle-feeding, 350
Biases, 52–53	Bouchard's nodes, 284

Bowel habits, 196, 260	Bronchovesicular breath sounds, 147b
Bowel sounds, 201 <i>b</i> –202 <i>b</i>	Brudzinski's sign, 331
Bowlegs, 355	Bruits
Brachial pulse, 219	abdominal, 201b-202b, 418, 420
Brachioradialis reflex, 329	carotid, 169, 417
Bradycardia, 358	Bulge sign, 293
Brain, 311–312	Bulimia, 60, 70t
abscess of, 112	Bullae, 99t
tumors of, 112, 127 <i>t</i> –128 <i>t</i> , 357	Burrows (scabies), 100t
Brainstem, 311	Bursae, 271
Branchial cleft cyst, 361	Bursitis, 275
Breast, 181-192	in hips, 289–290
cancer of, 182–183, 189 <i>t</i> –192 <i>t</i> ,	in knees, 292, 293, 308t
408-409	in shoulders, 307t
in children and adolescents, 364,	
375, 383 <i>t</i>	
common or concerning symptoms	C
of, 181 <i>b</i>	Café-au-lait spots, 358
discharge from, 186	CAGE questions, 50b, 199
examination of, 12-13, 183-188	Calcium, sources of, 72t
health history of, 6, 181	Calculating abilities, 84
health promotion and counseling	Cancer
for, 181–183	breast, 182–183, 189 <i>t</i> –192 <i>t</i>
masses in, 181, 182b	cervical, 244, 254t, 409
in older adults, 418	colorectal, 200, 209t, 261-262,
in pregnancy, 396	420
recording exam findings of, 189b,	lung, 155t, 158t
377 <i>b</i>	neck, 361
self-exam of, 183, 187b-188b	oral, 136t, 138t, 417
Breast-feeding, 350	ovarian, 420
Breath odor, 58	penis, 232, 236t
Breath sounds, 146, 147 <i>b</i>	prostate, 260-261, 269t, 420
in chest disorders, 161t	rectal, 269t
in newborns, 362 <i>b</i> , 363 <i>b</i>	screening for, in older adults,
Breathing. See also Respiratory system	408-409
abnormalities of, 74t	skin, 104 <i>t</i> –105 <i>t</i> , 106 <i>t</i> , 415
in comatose patient, 332, 333	testicular, 238t
in newborns, 358, 361, 362 <i>b</i>	vulvar, 252t
in pregnancy, 396	Candidiasis
rate and rhythm of, 64–65, 143	in infants, 361
shortness of breath, 142,	of tongue, $137t$
154 <i>t</i> –155 <i>t</i> , 163	vaginitis, 255 <i>t</i>
Breech presentation, 400	Canker sores, 138t
Bronchi, 141	Capacity, altered, 46–47
Bronchial breath sounds, 147b	Carcinoma. See also Cancer
Bronchiectasis, 157t	basal cell, 105 <i>t</i> , 415
Bronchiolitis, 358, 362b	of cervix, 254t
Bronchitis, 154t, 157t, 161t	of lip, 136 <i>t</i>
Bronchophony, 148b	squamous cell, 104 <i>t</i> , 105 <i>t</i> , 415

of tongue or floor of mouth, $138t$	Cheilitis, angular, 136t
of vulva, 252t	Chemicals, irritating, 159t
Cardiac syncope, 314	Cherry angioma, 103t, 415
Cardiomyopathy, 170, 174, 175	Chest pain, 141–142, 152 <i>t</i> –153 <i>t</i> ,
Cardiovascular system, 163-180	163–164. See also Thorax
chest pain caused by disorders of,	Cheyne-Stokes breathing, 74t
152 <i>t</i> –153 <i>t</i>	Chief complaints, 2b, 3, 349
common or concerning symptoms	Children, 349–388
of, 57, 163 <i>b</i>	development of, 349
cough caused by disorders of,	examining, 352-375 (See also spe-
158 <i>t</i> –159 <i>t</i>	cific system or structure)
disorders of, 176t–180t	head circumference in, 357
examination of, 13, 64, 168-175	health history of, 2b, 4, 349-351
health history in, 6, 163-164	health promotion and counseling
health promotion and counseling	for, 351
for, 164–167	heart murmurs in, 371-372,
in older adults, 417-418	381 <i>t</i> –382 <i>t</i>
in pregnancy, 396	hypertension in, 357b, 368, 380t
recording exam findings, 175b,	interviewing, 366b-367b
376 <i>b</i> –377 <i>b</i>	newborn maturity level classifica-
Carotid pulse, 169	tion, 378t
Carotid stenosis, 417	preventive care recommendations
Carpal tunnel syndrome, 283, 286,	for, 379 <i>t</i>
322	recording exam findings for,
Cartilaginous joints, 272b	375–377
Caruncles, 247, 419	sex maturity ratings for, 383t-387t
Cataracts, 113, 115, 416	sexual abuse in, 388t
Cauda equina, 274, 312	Chills, 198
Caviar lesions, 138t	Chloasma, 395
Cellulitis, 216	Choanal atresia, 361
Central nervous system, 311–312,	Cholecystitis, 208
339 t, 340 t	Cholesterol, 166
Cephalohematoma, 359	Chondromalacia, 292
Cerebellum, 14, 312, 340t	Chorea, 341t
Cerebrovascular disease, 315	Chronic obstructive pulmonary dis-
Cerumen, 416	ease, 154t, 161t, 163, 417
Cervix	Cirrhosis, 197b
abnormalities of, 254t	"Clasp-knife" resistance, 343t
cancer of, 244, 254t, 409	Claudication, 215, 224t
examination of, 15, 247-249	Clavicle fracture, 365
inflammation of (cervicitis), 254t	Clinical breast examination, 182
in pregnancy, 399	Clinical reasoning, 17-19. See also
shapes of, 253t	Assessments
Chalazion, 131t	Clinician
Chancre	behavior and appearance of, 38
lip, 136 <i>t</i>	relationship with patient, sexuality
penile, 232, 237t	in, 53
vulvar, 252t	self-awareness of, 52-53
Chart, medical. See Records, patient	Clitoris, 247

Clonus, 330	Cornea, 117
Clubbing, nail, 108t	Corneal light reflex test, 368
Cluster headaches, 125t	Corneal reflexes, 319
Coarctation of the aorta, 363	Coronary heart disease, 165-167
Cognitive function	Costovertebral angle tenderness, 205
assessment of, 83-84	Cough, 142, 156 <i>t</i> –159 <i>t</i>
higher, 79b, 84	Counseling. See Health promotion
in older adults, 404-405, 409	and counseling
Cogwheel rigidity, 343t	Cover-uncover eye test, 368
Cold sore, 136t	Crackles, 147b
Colitis, 210t	Cranial nerves, 312
Colles' fracture, 283	assessment of, 14, 318-320
Colonoscopy, 262, 409	function of, 316b-317b
Colorectal cancer	Cranial neuralgias, 128t
and diarrhea, 209t	Crohn's disease, 210t
in older adults, 420	Croup, 362b
screening for, 200, 260, 261-262,	Crust (skin), 101t
409, 420	Crying patient, 47
Coma, 332-334, 346t, 347t	Cryptorchidism, 233, 238t
Comedones, 415	Cultural differences, 52-53, 405
Communication, with patient, 44,	Cup-to-disc ratio, 416
53	Cutis marmorata, 358
Complaints, chief, 2b, 3, 349	Cyanosis, 363
Comprehensive health history. See	Cyanotic congenital heart disease,
Health history	358
Comprehensive physical examination.	Cyclothymic episode, 89t
See Physical examination	Cystocele, 247, 256t, 398
Concentration, difficulty with, 80	Cystourethrocele, 247, 256t
Conception age, 390	Cysts
Conductive hearing loss, 113, 135t,	"Baker's," 293, 310t
319	branchial cleft, 361
Condylar joints, 273b	breast, 182 <i>b</i>
Confidentiality, patient, 39, 46–47, 54	congenital, 361
Confusing patient, 46	epidermoid, 251t
Congestion, nasal, 114	epididymal, 239t
Congestive heart failure, 170, 364	periauricular, 361
Conjunctiva, 117	pilar, 116
Consciousness, level of	pilonidal, 263
assessing, 57, 78, 81	popliteal, 293
defined, 78b, 333b	skin, 98 <i>t</i>
in delirium or dementia, 424 <i>t</i>	thyroglossal, 361
loss of, 314	
Consolidation (lung), 161t	D
Constipation, 196, 197, 260	_
Constructional ability, 84	Data
Continuers, 40, 44	identifying, 2 <i>b</i> , 349
Contraception, 245	organizing, 28–30
Convergence, eye, 117	subjective and objective, 2b
Coordination, 323–325, 340 <i>t</i>	Deaf patient, 48

Davids and Julius 51, 52	D: 1 : 1 201 :
Death and dying, 51–52	Disc, herniated, 301t
Decision-making	Discharge
evidence-based, 18b	breast, 186
patient's capacity for, 46–47, 80	penile, 231
Deep tendon reflexes, 328, 340t	vaginal, 243, 255 <i>t</i> , 373
Defecation, pain with, 260	Disease, defined, 42
Degenerative joint disease, 305 <i>t</i> –306 <i>t</i> ,	Disease/illness model, 42
308 <i>t</i>	Disruptive patient, 47
Dehydration, 358	Dissecting aortic aneurysm, 152t,
Delirium, 46, 424 <i>t</i> –425t	163
Delusional disorder, 92t	Distress, signs of, 57
Dementia, 46, 409, 424t-426t	Diverticulitis, 196
Dental caries, 115, 371, 417	Dizziness, 313
Depression	Do Not Resuscitate (DNR) orders,
in older adults, 411b, 423b	52, 407
screening for, 57, 79-80, 411b	Doll's eye movement, 334
symptoms of, 88 <i>t</i> –89 <i>t</i>	Domestic violence, 51, 392–393
DeQuervain's tenosynovitis, 322	Dorsalis pedis pulse, 220
Dermatomes, 325, 344 <i>t</i> –345 <i>t</i>	Double vision, 113, 313
Development of children, 350, 356	Draping patient, 10
Diabetes, 199	Drawer sign, anterior, 294, 309t
Diabetic retinopathy, 133t	Dress, patient's, 58, 81
Diagnoses, establishing, 17 <i>b</i> –19 <i>b</i>	"Drop-arm" test, 281, 307t
DIAPERS assessment, 410	Drugs
	abuse of, 4, 50-51, 80
Diaphragm, 146	diarrhea due to, 209t
Diarrhea, 196, 209 <i>t</i> –210 <i>t</i> , 260	incontinence due to, 212t
Diastolic blood pressure, 63 <i>b</i> , 414	medication history, 2b, 4
Diet	use in older adults, 406
changes in, for hypertension, 73t	Drusen, 120b, 416
in health history, 5, 350	Dullness to percussion, 145b
health promotion and counseling	Durable power of attorney for health
for, 56–57	care, 47, 407
in older adults, 406	Dying patient, 51-52
Diethylstilbestrol (DES), fetal expo-	Dysarthria, 82, 313
sure to, $254t$	Dysesthesias, 314
Diffuse esophageal spasm, 153t	Dyskinesias, oral-facial, 342t
Diffuse idiopathic skeletal	Dyslipidemias, 166
hyperostosis, 302t	Dysmenorrhea, 242
Diffuse interstitial lung diseases,	Dyspareunia, 243
155 <i>t</i>	Dyspepsia, 195
Digit span, 83	Dysphagia, 196
Digital rectal examination, 261. See	Dysphonia, 82
also Rectum	Dyspnea, 142, 154 <i>t</i> –155 <i>t</i> , 163
Diphtheria, 139t	Dysrhythmias, 64, 176 <i>t</i> –180 <i>t</i> , 364.
Diplegias, spastic, 366	See also specific dysrhythmias
Diplopia, 113, 313	Dysthymic disorder, 89t
Disability, physical, 411 <i>b</i> –412 <i>b</i> ,	Dystonia, 342t
423 <i>b</i>	Dysuria, 198

E	Epitrochlear nodes, 12, 219
Ear canal, 121, 369	Epstein's pearls, 361
Eardrum, 121, 134t, 369-370	Erectile dysfunction, 230
Ears. See also Hearing; Hearing loss	Erosion, skin, 102t
in children, 360, 369–370	Esophagus, 142, 153t
disorders of, $135t$	Estrogen-producing tumors, 419
examination of, 12, 120–122	Ethics, 53–54
health history of, 6, 113–114	Ethnicity, 52
in older adults, 416	Evidence-based decision making, 181
pain in (earache), 113	Excoriation, skin, 102t
recording exam findings, 124b, 376b	Exercise
Eating disorders, 60, 70t	health promotion and counseling
Ecchymosis, 104t	for, 57, 276
Echoing, 44	history of, 5
Ectopic pregnancy, 196, 400	in older adults, 408
Ectropion, 130 <i>t</i> , 416	during pregnancy, 392
Edema, 164	Expected date of delivery (EDD),
of breast skin, 192 <i>t</i>	390–391
dependent, 164	Expiratory time, forced, 150
pitting, 220	Expressions, facial, 58, 81
in pregnancy, 395, 400	Extinction sensation, 327
scrotal, 236t	Extra-articular structures, 271, 275
Effusion	Extraocular muscles, 117, 318
around knee, 293-294, 295	Extremities, 14, 400. See also specific
behind eardrum, 134t	structures
pleural, 146, 161 <i>t</i>	Eyebrows, 116
Egophony, 148b	Eyelids
Ejaculation, 230	disorders of, 130t
Elbows, 282, 322	inspection of, 116
Elderly patients. See Older adults	in older adults, 415–416
Embolism, pulmonary, 155t, 158t	Eyes. See also Vision
Emotional cues, of patient, 40	in children, 360, 368–369
Empathy, for patient, 44	disorders of, 115, 126t, 129t-133t
"Empty can test," 280	examination of, 12, 116-120
Endocervical polyp, 254t	and headaches, 112, 126t
Endocrine disorders, 7, 356	health history of, 6, 112-113
Endometrial hyperplasia, 419	health promotion and counseling
Endometriosis, 242	for, 115
Engagement (fetal), 401	in older adults, 415-416
Entropion, 130t, 416	during pregnancy, 395
Environment for examination, 10, 38,	recording exam findings, 124b, 376b
39, 404	
Epidermoid cysts, 251t	
Epididymis, 233, 239t, 420	F
Epiglottitis, 362b	Face
Episcleritis, 130t	edema of, 395
Episiotomy, 398	examination of, 116
Epistaxis, 114	expressions of, 58, 81

of infants, 359	Fever blister, 136t
paralysis of, 116, 339 t	Fibroadenoma of breast, 182b
during pregnancy, 395	Fibrocystic breast disease, 182b
Facial nerve, 317 <i>b</i> , 319	Fibromyalgia, 275
Failure to thrive, 356	Fibrous joints, 272b
Fainting, 314	Fingernails, 95, 108 <i>t</i> –109 <i>t</i>
Fallot, tetralogy of, 381 <i>t</i> –382 <i>t</i>	Fingers, 216, 284–286, 322, 365
Falls, in older adults, 276, 412, 413 <i>f</i>	Fissures
False positive or negative results,	skin, 102 <i>t</i>
30 <i>b</i> , 31 <i>b</i>	tongue, 137t
Family history, 3 <i>b</i> , 5, 51, 390	Flaccidity, 334, 343 <i>t</i>
Family planning, 245	Flail chest, 160t
Farsightedness (hyperopia), 112, 126t	Flank pain, 198
Fasciculations, 341t	Flatness to percussion, 145 <i>b</i>
Fatigue, 55	Folate, sources of, 72t
Fecal occult blood test, 261–262	Fontanelles, 358-359
Feeding history, 350	Forced expiratory time, 150
Feet	Foreign bodies
coldness, numbness or pallor in,	in airway, 362 <i>b</i>
216	in vagina, 373
deformities of, 366	Fovea, 120 <i>b</i>
examination of, 297-298	Fractures
swelling of, 216	of clavicle, 365
ulcers on, 225t	Colles', 283
Females	scaphoid, 284
axillae in, 12, 186	Fremitus, tactile, 145
breast examination in, 12-13,	Friction rubs, 171, 201
183–185, 187 <i>b</i> –188 <i>b</i>	Fullness, abdominal, 195
epitrochlear nodes in, 12	Functional ability, 410
genital exams in, 7, 15, 241-257	Fundal height, 397
in children, 365, 373, 375	Fundi, eye, 118-119, 416
in older adults, 419-420	Funnel chest (pectus excavatum),
in pregnancy, 398-400	160 <i>t</i>
recording results of, 250b, 377b	
Pap smears in, 244, 247-248,	
394, 399, 409, 419	G
rectal exams in, 15, 250, 264	Gag reflex, 320
sexual maturity of, 246, 383t,	Gait
386 <i>t</i> –387 <i>t</i>	inspection of, 58, 289, 291, 313,
Femoral hernia, 234, 240t, 250	324
Femoral pulse, 220	in older adults, 420-421
Fetus	Gallbladder, 142
engagement of, 401	Gases, irritating, 159t
exposure to DES, 254t	Gastroesophageal reflux disease, 158t,
heart of, 397-398	195
lie of, 400	Gastrointestinal system
movement of, 397	bleeding in, 260
presentation of, 400-401	chest pain caused by disorders of,
Fever, 56, 65–66, 198	153 <i>t</i>

Gastrointestinal system (continued) common or concerning symptoms of, 193b examination of, 201–203 health history of, 6, 193–197 Gaze, six cardinal directions of, 117 Gegenhalten, 343t General survey, 55–61 examination techniques for, 11, 57–66 health history in, 5, 55–56 health promotion and counseling in, 56–57 in infants, 356–357 interpreting findings of, 70t–74t recording findings of, 69 Genital examinations in females, 7, 15, 241–257 in children, 365, 373, 375, 386t–387t, 388b common concerns of, 241b	Giant cell (temporal) arteritis, 128t Gingivitis, 417 Glaucoma, 113, 115, 126t, 416 Glaucomatous cupping, 132t Glossopharyngeal nerve, 317b, 320 Goals for interview, 38 Goiter, 115, 124, 140t Gonococcal infections, 231, 283 Gout, 275, 421 Granulomatous colitis, 210t Graves' disease, 140t Gray matter, 311 Great arteries, transposition of, 363, 382t Grip, 322 Grooming, patient, 58, 81 Growth of children, 350, 356 Guarding, 202 Gums, 122 Gynecologic history, 4
common concerns of, 241 <i>b</i> health promotion and counseling for, 244–245 interpreting results of, 251 <i>t</i> –257 <i>t</i> in older adults, 419–420 in pregnancy, 398–400 recording results of, 250 <i>b</i> , 377 <i>b</i> techniques for, 246–250 in males, 6–7, 15, 229–240 in children, 365, 372–373, 375, 384 <i>t</i> –385 <i>t</i> , 388 <i>b</i> common or concerning symptoms, 229 <i>b</i> health promotion and counseling for, 231–232 interpreting results of, 236 <i>t</i> –240 <i>t</i> in older adults, 420 recording results of, 235 <i>b</i> , 377 <i>b</i> techniques for, 232–235 Genital herpes, 232, 237 <i>t</i> , 251 <i>t</i> Genitalia, ambiguous, 365 Geographic tongue, 137 <i>t</i> Geriatric syndromes, 404. <i>See also</i> Older adults Gestation, expected weeks of, 390–391	Hair in infants, 358 inspection of, 95, 116 loss of, 93b, 107t-108t, 216 during pregnancy, 395 pubic, 384t-385t, 386t-387t Hairy leukoplakia, 137t Hairy tongue, 137t Hand-washing, 16 Hands, 283-287 arterial supply to, 222 flaccid, 334 in pregnancy, 400 Hashimoto's thyroiditis, 140t Head circumference of, 357 common or concerning symptoms of, 111b examination of, 12, 116 health history of, 6, 111-112 of infants, 357, 358-359 in pregnancy, 395-396 recording exam findings, 124b,
Gestational age, 353, 354 <i>b</i> , 378 <i>t</i> Gestational hypertension, 395 <i>b</i> , 400	376 <i>b</i> Headaches, 111–112, 125 <i>t</i> –129 <i>t</i> , 313 Health care proxy, 48, 52, 407

Health history. See also Interview	for smoking, 142, 392
of children, 349-351	for STDs or HIV, 231-232, 245
components of, 1, 2b-3b	for stroke prevention, 165,
comprehensive, 1, 3-7, 37	315–316
focused, 1, 10, 37	for suicidality, 80
format of, 37	for testicular self-examination, 232
interviewing process, 38-54	for transient ischemic attacks,
of older adults, 403-405	315-316
in pregnancy, 389-391	for vision, 115b
recording findings of, 33-36	for weight, 56b
Health Insurance Portability and	Hearing
Accountability Act (HIPAA),	health history of, 113-114
46–47	in infants, 360
Health maintenance, 4, 18b	in older adults, 408, 411 b, 416,
Health promotion and counseling	422b
for abdominal aortic aneurysm,	testing, 12, 121-122, 319-320
217–218	Hearing loss and deafness
for alcohol and substance abuse,	history of, 113
80, 199	interviewing patient with, 48
for back pain, 276	in older adults, 115
for breast health, 181-183	patterns of, 135t
for cardiovascular disease,	Heart. See also Cardiovascular system
164–167	in children, 363-364, 371-372
for children, 351	coronary heart disease, 165-167,
for colon cancer screening, 200,	358
261–262	examination of, 13, 169-175
for depression, 79-80	health history of, 6
for diet, 56b	in older adults, 417-418
for domestic violence, 392-393	in pregnancy, 396
for dyslipidemias, 166	Heart failure
for exercise, 56b, 276, 392	congestive, 170
for fall prevention, 276	left ventricular, 154t, 158t, 161t,
for family planning, 245	163
for hearing loss, 115b	Heart murmurs
for hepatitis, 199-200	causes of, 180t
for hypertension, 164	in children, 371–372, 381 <i>t</i> –382 <i>t</i>
for immunizations, 393	gradations of, 173b
for menopause, 245	identifying, 171-173
for nutrition, 56b, 276, 391	in older adults, 418
for older adults, 408-409	in pregnancy, 396
for oral health, 115b	Heart rate
for osteoporosis, 277	fetal, 397-398
for Pap smear screening, 244-245	in infants, 358
for peripheral arterial disease, 217	measuring, 64, 168
for pregnancy, 391–393	in older adults, 414
for prostate cancer, 260–261	Heart rhythm, 64, 364, 398
for renal artery disease, 217	Heart sounds
for skin care, 93b	in infants, 364

Heart sounds (continued)	genital, 232, 237t, 251t
interpreting, 171, 176 <i>t</i> –180 <i>t</i>	in infants, 358
in older adults, 417-418	oral (cold sore), 136t
Heartburn, 195	Herpes zoster, 325, 415
Heberden's nodes, 284	Hinge joints, 273b
HEENT (Head, eyes, ears, nose	Hips, 289–291
throat)	dysplasia of, 365
examining, 12, 116-124	muscles of, 323
health history of, 6, 111–115	in newborns, 365–366
health promotion and counseling	History. See Health history
for, 115	HIV (human immunodeficiency
interpreting findings for, 125t–140t	virus), 231-232, 243, 245
in older adults, 415–417	Hoarseness, 114
recording findings for, 124b, 376b	"Housemaid's knee," 292, 308t
Hegar's sign, 399	HPV (human papillomavirus), 244
Height, 58	Humility, cultural, 52
of children and adolescents, 356	Hydrocele, 233, 236t, 365
of older adults, 414	Hydrocephalus, 116, 357, 358
of pregnant women, 395	Hygiene, personal, 58, 81
Hematologic system, 7	Hymen, 247, 373
Hematoma, subdural, 357	Hyperemesis, 395
Hemianopsia, 113	Hyperopia (farsightedness), 112,
Hemolytic disease, 358	126 <i>t</i> , 368
Hemoptysis, 142, 156 <i>t</i> –159 <i>t</i>	Hyperpnea, 74t
Hemorrhage	Hyperpyrexia, 65
subarachnoid, 112, 127 <i>t</i> , 313	Hyperresonance, 145 <i>b</i>
vitreous, 112	Hypertension
Hemorrhoids, 268t, 398	in children and adolescents, 357 <i>b</i> ,
Hepatitis	368, 380 <i>t</i>
alcoholic, 193, 197 <i>b</i>	chronic, 395b
health promotion and counseling	classification of, 63b, 164
for, 199–200	dietary changes for, 74 <i>t</i>
and liver disease, 197 <i>b</i>	gestational, 395b, 400
symptoms of, 195	in older adults, 413–414
types of, 197 <i>b</i> , 199–200	pulmonary, 364
Hereditary hemorrhagic telangiecta-	screening for, 164
sia, 136 <i>t</i>	Hyperthyroidism, 115, 116
Hernias	Hypertonia, 343 <i>t</i>
in children, 355, 365, 372	Hypertrophic cardiomyopathy, 174,
in females, 250	175
femoral, 234, 240t, 250	Hyperventilation, 74t
inguinal, 234, 236t, 240t, 250,	Hypoglossal nerve, 317b, 320
365, 372	Hypomanic episode, 89t
in males, 15, 233, 234, 236t, 240t,	Hypoplastic left heart syndrome, 363
365	Hypospadias, 232, 236t
scrotal, 236t	Hypotension, orthostatic, 63 <i>b</i> , 168,
umbilical, 355	414
Herniated disc, 301t	Hypothalamus, 311
Herpes simplex infection	Hypothermia, 65, 66

Hypotheses, generating and testing, 18b, 41–42	for children, 366 <i>b</i> –367 <i>b</i> comprehensive <i>vs.</i> focused, 1 <i>b</i> , 37
Hypothyroidism, 115	dealing with sensitive topics, 49–54
Hypotonia, 343t, 366	sequence of, 39–43
Hysterectomy, 244	for specific situations, 46–49
	techniques for, 43–45
and the second s	Intracranial pressure, 358
	Introitus, 247
Icterus (jaundice), 197, 358	Involuntary movements, 321, 340t,
Identifying data, 2b, 349	341 <i>t</i> -342 <i>t</i>
Iliopectineal bursa, 289	Iris, 117
Iliotibial band bursitis, 308t	Iron, sources of, 72t
Illnesses. See also specific illnesses	Irritable bowel syndrome, 209t
defined, 42	Ischiogluteal bursa, 290
history of, $2b$, 4	Isolated systolic hypertension, 64
patient perspective on, 40b, 42b	Itching
present, 2b, 3-4, 350	in eyes, nose, throat, 114
Immunizations, 4, 350, 393, 408	vaginal, 243
Impotence, 230	
Incontinence, urinary	
in older adults, 410, 411b, 422b	J
types of, 199, 211 <i>t</i> –212 <i>t</i>	Jaundice, 197, 358
Infants. See also Children	Joints. See also specific joints
examining, 352-366	assessing, 271-273
head circumference of, 357	common or concerning symptoms
hypertension in, 357b, 368, 380t	of, 273 <i>b</i>
maturity classification of, 378t	examination of, 278-310
Infections. See also specific infections	health promotion and counseling
and diarrhea, 209t, 210t	for, 276–277
sexually transmitted, 231-232,	pain in and around, 273-276,
243, 245, 373	305 <i>t</i> –306 <i>t</i>
Inflammation, 202, 275. See also	Judgment, patient, 77, 78b, 83, 425t
Arthritis	Jugular veins, 168
Inflammatory bowel disease, 210t	Jugular venous pressure, 168, 417
Information assessment, 84	
Inguinal hernia	
in children, 365, 372	K
direct, 234, 240t	Kaposi's sarcoma, 105t
indirect, 234, 236t, 240t, 250	Keloids, 101t
Inguinal ligament, 289	Keratoses, 104 <i>t</i> –105 <i>t</i> , 415
Insight, patient, 77, 78b, 83	Kernig's sign, 331
Intelligence, limited, 49	Kidneys, 14, 198, 205
Intercourse, pain during, 243	Klinefelter's syndrome, 238t
Intermittent claudication, 215, 224t	Knees
Interpreters, 47, 48b	examination of, 291-296
Interstitial lung diseases, diffuse, 155t	extension and flexion of, 323
Interview, 37-54. See also Health	painful, tender, 308t-310t
history	reflexes, 329, 400
approach to, 38, 403-405	Koplik's spots, 139t

Korotkoff sounds, 62b, 174 Kubler-Ross stages of dying, 51 Kussmaul breathing, 74t	Libido, 230 Lichenification, 101 <i>t</i> Lifestyle habits, 5, 167
Kyphoscoliosis, thoracic, 160t	Ligaments, 271
	knee, 294–295, 296, 309 <i>t</i> –310 <i>t</i>
L	Light reflex, 120 <i>b</i> Lighting for examination, 10
Labia, 246	Lipid therapy, 166
Lachman test, 295	Lips, 122, 136 <i>t</i>
Lacrimal apparatus, 116	Listening, active, 43
Lactating adenomas, 182b	Literacy, low, 48
Lactose intolerance, 210t	Liver
Language, 41, 79b, 82	in children, 372
Language barriers, 47, 48b	disease of, 197–198
Laryngitis, 156t	examination of, 14, 203–204
Laryngotracheobronchitis (croup),	toxic damage to, 198 <i>b</i>
362 <i>b</i>	"Liver spots," 415
Lateral collateral ligament (LCL),	LMP (last menstrual period),
292, 294	390–391
Lateralization test, 122, 319	Low-density lipoproteins (LDL),
Laxatives, abuse of, 210t	166
LDL (low-density lipoproteins), 166	Lumbar spinal stenosis, 301t
Lead-pipe rigidity, 343t	Lung sounds. See Breath sounds
Learning ability, 83	Lungs
Left ventricular heart failure, 154 <i>t</i> ,	abscess of, 157t
158 <i>t</i> , 161 <i>t</i> , 163	assessing function of, 150
Left ventricular hypertrophy, 417	cancer of, 155 <i>t</i> , 158 <i>t</i>
Legs	common or concerning symptoms
coldness, numbness or pallor in,	of, 141 <i>b</i>
216	disorders of, 154 <i>t</i> –159 <i>t</i> , 161 <i>t</i>
examination of, 14, 219–221	and dyspnea, 154 <i>t</i> –155 <i>t</i>
flaccid, 334 hair loss on, 216	examination of, 12, 13, 143-150
	health history of, 141-142
measuring length of, 299 mobility in older adults, 411 <i>b</i> ,	health promotion and counseling
422 <i>b</i>	for, 142
pain in, 215	of newborns, 361-363
in pregnancy, 400	in older adults, 417
ulcers on, 216	in pregnant women, 396
and vascular disease, 215–216,	recording exam findings of, 151b,
224 <i>t</i> –225 <i>t</i>	376b
Lens, eye, 117, 416	Lymph nodes
Leopold's maneuvers, 398, 400-402	axillary, 12, 186
Lesions	cervical, 12
oral, 136 <i>t</i> –139 <i>t</i>	in children, 361, 376b
skin, 57, 94, 97 <i>t</i> –104 <i>t</i>	epitrochlear, 12, 219
Lethargy, 333b	examination of, 123, 124b
Leukonychia, 109t	inguinal, 221
Leukoplakia, hairy, 137t	Lymphadenopathy, 361

M	Meningeal signs, 331
Macula, 119, 120 <i>b</i>	Meningitis, 112, 127t, 358
Macular degeneration, 113, 115,	Menisci of knee, 292, 296, 309t
120 <i>b</i> , 416	Menopause, 243, 245
Macules, 97 <i>t</i>	Menstrual cycle, 242, 390
Malabsorption syndromes, 210t	Mental status, 75–92
Males	of children, 355–356, 367, 377b
axillae in, 12, 186	common or concerning symptoms
breast exam in, 13, 186	of, 77 <i>b</i>
epitrochlear nodes in, 12	in delirium and dementia,
genital exams in, 6–7, 15,	424 <i>t</i> –425 <i>t</i>
229–240	disorders of, 46-47, 75b-76b,
in children, 365, 372–373, 375,	87 <i>t</i> –92 <i>t</i>
384 <i>t</i> –385 <i>t</i> , 388 <i>b</i>	examination of, 14, 77b, 78b-79b,
common or concerning symp-	81-85
toms of, 229 <i>b</i>	health history of, 4, 7, 50, 77-79
health promotion and counsel-	health promotion and counseling
ing for, 231–232	for, 79-80
interpreting results of,	recording findings of, 86b, 377b
236 <i>t</i> –240 <i>t</i>	Microcephaly, 357, 358
in older adults, 420	Migraine headaches, 111, 112, 125 <i>t</i>
recording results of, 235b, 377b	Mini-Cog test, 426t
techniques for, 232-235	Mini-Mental State Examination, 85
rectal exam in, 15, 262-264	Mitgehen, 343t
sexual maturity in, 384t-385t	Mitral valve
Malocclusion, 371	prolapse of, 174, 175
Malodor, 417	regurgitation of, 171, 418
Mammography, 182-183, 408-409	stenosis of, 154t, 158t, 163, 171
Manic episodes, 88 <i>t</i> –89 <i>t</i>	Mixed mood disorder, 89t
Manner, patient, 81	Moles, 93, 106 <i>t</i>
Masseter muscles, 278, 319	Mongolian spots, 358
Mastitis, 182b	Monoarticular joint pain, 274
Mastoiditis, 121	Mood. See also Depression
McMurray test, 296	assessing, 77, 78, 82
Medial collateral ligament (MCL),	defined, 79b
292, 296, 310 <i>t</i>	in delirium and dementia, 424t
Median nerve compression (carpal	disorders of, 88 <i>t</i> –89 <i>t</i>
tunnel syndrome), 283, 286,	Motor system (motor activity)
322	disorders of, 340t
Medical chart. See Records, patient	examination of, 14, 58, 321–325
Medications. See Drugs	and mental status, 81
Melanoma, 93, 106t, 415	Mouth
Melena, 197	abnormalities of, 136 <i>t</i> –139 <i>t</i>
Memory, 78 <i>b</i>	cancers of, 138 <i>t</i> , 409, 417
assessing, 77, 78, 83	candidiasis in, 137 <i>t</i> , 361
in delirium and dementia, 425 <i>t</i>	in children, 361, 370–371
in older adults, 409, 411 <i>b</i> , 423 <i>b</i>	examination of, 12, 122–123
Ménière's disease, 114	health history of, 6, 114-115

Mouth (continued)	Nasal congestion, 114
health promotion and counseling	Nausea, 112, 195
for, 115	Near reaction, 117
in older adults, 417	Near-sightedness, 113
in pregnant women, 395	Neck
recording exam findings of, 124b,	and chest pain, 142
376 <i>b</i>	common or concerning symptoms
Movements	of, 111 <i>b</i>
coordinating, 323-324	examination of, 12, 123-124
involuntary, 321, 340t, 341t-342t	health history of, 6, 114-115
MRSA precautions, 15-16	of infants, 361
Multiple sclerosis, 322	pain in, 274, 303 <i>t</i> –304 <i>t</i>
Murmurs. See Heart murmurs	range of motion in, 289
Murphy's sign, 208	recording exam findings of, 124b,
Muscles	376 <i>b</i>
bulk of, 321, 340t	Negative predictive value, 31 <i>b</i> –32 <i>b</i>
disorders of, 343t	Negatives, true and false, 30b, 31b
strength or weakness of, 55,	Neoplasms. See Cancer; Tumors
321–323, 340 <i>t</i>	Nerves
tone of, 321, 340t, 343t	cranial, 14, 312, 316b-317b,
Musculoskeletal system, 271–310.	318–320
See also specific structures	peripheral, 312
abnormalities of, 301 <i>t</i> –310 <i>t</i>	root compression of, 325
assessing joints, 271-273	spinal, 312
in children, 365–366, 373	Nervous system, 311–347
common or concerning symptoms	in children, 355, 366, 374
of, 273 <i>b</i>	common or concerning symptoms
examination of, 278-310	of, 313–315
health history of, 7, 273-276	components of, 311-312
health promotion counseling for,	disorders of, 336t-347t
276–277	examination of, 14, 316-334
in older adults, 420-421	health history of, 7
range of motion in, 13, 275, 299	health promotion and counseling
recording exam findings, 300b, 377b	for, 315–316
Myalgia, 275	in older adults, 421
Myasthenia gravis, 314, 322	recording exam findings, 335b, 377b
Mycoplasma infections, 156t	Neuralgias, cranial, 128t
Myelopathy, cervical, 304t	Neuropathic ulcer, 225t
Myocardial infarction, 152t, 163, 194	Neuropathy, peripheral, 316
Myocardium, 141, 364	Nevi, 93, 106t
Myoma, uterine, 257t	Newborns. See also Children; Infants
Myopathy, 314, 322	Apgar score of, 352–353
Myopia, 113, 368	assessing, 352–355
Myxedema, 116	birth weight of, 353, 354b, 378t
	gestational age of, 353, 354b, 378t
	head circumference of, 357
N	height and weight of, 356
Naegele's rule, 390	hypertension in, 357b, 380t
Nails, 95, 108 <i>t</i> –109 <i>t</i>	maturity classification of, 354b, 378t

	01 . 200
vital signs in, 357–358	Obturator sign, 208
Night sweats, 56	Occlusion, arterial, 420
Nipples	Occult blood test, 261–262
discharge from, 186	Ocular fundus, 118–119
inspection of, 184, 185	Ocular movements, 333
Paget's disease of, 192t	Oculocephalic reflex, 334
in pregnancy, 396	Oculomotor nerve, 316 <i>b</i> , 318
retraction and deviation of, 191 <i>t</i>	Odors, body and breath, 58, 417
Nocturia, 199	Odynophagia, 196
Nodules	Older adults, 403–426
breast, 185	adjusting environment for, 404
in rheumatoid arthritis, 284	common concerns of, 405–408
skin, 98 <i>t</i>	content and pace of visit with, 404
thyroid, 140t	cultural differences in, 405
Nonarticular joint pain, 275	delirium and dementia in, 409,
Nonmaleficence, 53	424 <i>t</i> –426 <i>t</i>
Nonverbal communication, 44	eliciting symptoms in, 404–405
Nose	examining, 410–421
bleeding from, 114, 395	falls in, 276, 412, 413f
congestion in, 114	health history in, 403–405
examination of, 12, 122	health promotion and counseling
health history of, 6, 114	for, 408–409
in newborns, 361	mistreatment of, 409
in pregnant women, 395	osteoporosis in, 277
recording exam results of, 124b,	recording exam findings of,
376 <i>b</i>	422–423
Note-taking, 38	Olfactory nerve, 316b, 318
Nuchal rigidity, 361	Onycholysis, 109t
Number identification, 327	Open-angle glaucoma, 113, 416
Numbness, in legs or feet, 216	Opening snap, 176t
Nutrition	Ophthalmoscope, 118b
health promotion and counseling	Optic atrophy, 132t
for, 56–57, 72 <i>t</i> –73 <i>t</i> , 276	Optic disc, 118, 119 <i>b</i> –120 <i>b</i> , 132 <i>t</i> , 318
and hypertension, 73t	Optic nerve, 316 <i>b</i> , 318
in older adults, 406, 411 <i>b</i> ,	Oral cavity. See Mouth
422 <i>b</i> –423 <i>b</i>	Oral-facial dyskinesias, 342t
during pregnancy, 391	Oral temperature, 65–66
screening checklist for, 71t	Orchitis, 238t
sources of nutrients, 72t	Orgasm, 230
Nystagmus, 360	Orientation to identity, place, and
	time, 78 <i>b</i>
	assessing, 77, 83
0	in delirium and dementia, 425 t
Obesity, 56–57	in older adults, 409
Objective data, 2 <i>b</i>	Orthopnea, 163
Obsessive-compulsive disorder,	Orthostatic hypotension, 63b, 168,
90 <i>t</i> –91 <i>t</i>	414
Obstetric history, 4, 390	Ortolani test, 365
Obtundation, 333b	Osteoarthritis, 305 <i>t</i> –306 <i>t</i> , 420

Osteoporosis, 277	Patients
Otitis externa, 113, 120, 121, 370	with altered capacity, 46-47
Otitis media, 113, 121, 134 <i>t</i> , 370	angry or disruptive, 47
Otoscope, pneumatic, 370	approaching, 10, 403–405
Ovaries	comfort of, 10, 39, 394 <i>b</i>
cancer of, 420	confidentiality of, 39, 46–47, 54
examination of, 15, 249, 400	confusing, 46
Overweight patients, 56–57	crying, 47
Ovulation, 245	culture and ethnicity of, 52–53
Ovination, 243	dying, 51–52
	emotional cues of, 40
P	empowering, 45
Paget's disease of the nipple, 192 <i>t</i>	
Pain, 56, 57. See also specific types of	establishing rapport with, 39, 366b
pain	facilitating story of, 40, 41
* .	with impaired hearing, 48
acute and persistent, 67–69, 406, 407 <i>b</i>	with impaired vision, 49
	interviewing, 37–54
response to, 325	with language barriers, 47, 48 <i>b</i>
Palliative care, 407–408	with limited intelligence, 49
Palpitations, 163	with low literacy, 48
Panic attack, 90t, 153t	partnerships with, 45, 53
Panic disorder, 90 <i>t</i>	with personal problems, 49
Pap smears	perspective on illness, 40 <i>b</i> , 42 <i>b</i>
in older adults, 244, 409, 419	position of, 9 <i>b</i> , 11, 394 <i>b</i> , 396
in pregnancy, 394, 399	sexual relationship with clinician, 53
schedule of testing for, 244	silent, 46
technique for, 247–248	talkative, 47
Papilledema, 120 <i>b</i> , 132 <i>t</i>	Peau d'orange sign, 192 <i>t</i>
Papillomavirus, human (HPV), 244	Pectoriloquy, whispered, 148b
Papules, 97t	Pectus carinatum (pigeon chest), 160t
Paradoxical pulse, 174	Pectus excavatum (funnel chest), 160t
Paralanguage, 44	Pediatric patients. See Children
Paralysis, 116, 313, 334	Pelvic examination, 246–250
Paratonia, 343 <i>t</i>	in older adults, 419–420
Paravertebral muscles, 288	in pregnancy, 399–400
Paresthesias, 314	Pelvic floor relaxations, 256t
Parietal pain, 194	Pelvic inflammatory disease, 196, 242,
Parkinsonism, 340t	249
Paronychia, 108 <i>t</i>	Pelvis, back pain referred from, 302t
Paroxysmal nocturnal dyspnea, 163	Penis
Paroxysmal supraventricular tachy-	abnormalities of, 231, 236t
cardia, 358, 364	cancer of, 232, 236 <i>t</i>
Partnering, with patient, 45, 53	in children and adolescents,
Patches, skin, 97t	384 <i>t</i> –385 <i>t</i>
Patella, 292, 295	discharge from, 231
Patellofemoral disorder, 292, 309t	examination of, 15, 232
Pathologic processes, 17b	in older adults, 420
Pathophysiologic processes, 18b	Perceptions, patient's, 77, 78b, 83,
Patient records See Records patient	425t

Percussion, chest, 145–146, 149–150,	interpreting findings of, 70t–74t
161 <i>t</i>	lighting and environment for, 10,
Periauricular cysts and sinuses, 361	38, 39, 404
Pericardial friction rubs, 171	in older adults, 413-421, 423b
Pericarditis, 141, 152t	patient comfort in, 10, 394b
Perineal irritation, 373	positions for, 9b, 11, 394b, 396
Peripheral arterial disease, 215-216,	preparing for, 8
217	recording findings of, 33-36, 69
Peripheral nervous system, 311, 312.	sequence for, 9b, 11, 352
See also Nervous system	Pigeon chest (pectus carinatum), 160t
disorders of, 340t	Pigmentation, skin, 358
lesion of, 339t	Pilar cysts, 116
Peripheral neuropathy, 316	Pilonidal cyst, 263
Peripheral vascular system, 215–227	Pinguecula, 130t
abnormalities of, 224t-225t	Plan, developing, 19b, 42
common or concerning symptoms	Plantar reflexes, 330, 340t
of, 215 <i>b</i>	Plaques, skin, 97t–98t
examination of, 13, 218-223,	Pleura
226 <i>t</i> –227 <i>t</i>	inflammation of (pleurisy), 194,
health history of, 7, 215-216	213 <i>t</i>
health promotion or counseling	pain in, 141, 153 <i>t</i>
for, 216–218	Pleural effusion, 146, 161t
in older adults, 420	Pneumonia, 155t, 156t, 358, 362b
recording exam results of, 223b	Pneumothorax, 155t, 161t
Peritoneal inflammation or tender-	Point localization, 327
ness, 202, 213t	Point of maximal impulse (PMI)
Peritonsillar abscess, 371	in children, 363-364
Personal history, 3b, 5	measuring, 64, 168, 170
Personal problems, patient with, 49	in older adults, 417
Pes anserine bursitis, 293, 308t	in pregnancy, 396
Petechia, 103t	Polyarticular joint pain, 274-275
Peutz-Jeghers syndrome, 136t	Polydactyly, 365
Peyronie's disease, 236t	Polymyalgia rheumatica, 275, 276
Phalen's sign, 287	Polyneuropathy, 314, 322
Pharynx	Polyps
abnormalities of, 139t	endocervical, 254t
in children, 361, 370–371	rectal, 268t
examination of, 12, 123	Polyuria, 199
health history of, 6	Popliteal ("Baker's") cyst, 293, 310t
sore (pharyngitis), 114, 139t	Popliteal pulse, 220
Phimosis, 232	Position, patient's, 9 <i>b</i> , 11, 394 <i>b</i> , 396
Phobias, 90t	Position sense, 326
Physical abuse, 51, 392–393	Positive predictive value, 31 <i>b</i>
Physical examination, 8–16. See also	Positives, true and false, 30 <i>b</i> , 31 <i>b</i>
specific system or structure	Posterior cruciate ligament (PCL), 296
approach to, 10, 403–405	Posterior tibial pulse, 221
in children, 352–375	Postictal state, 315
comprehensive vs. focused, 1, 10	Postnasal drip, 157t
health history in, 1 <i>b</i> , 2 <i>b</i> –3 <i>b</i> , 3–7	Posttraumatic stress disorder, 91t

Postural hypotension, 63b, 168	Psychotic disorders, 92t
Posture, 58, 81, 333	Pterygoid muscles, 278
Precautions, standard and universal,	Ptosis, 130t, 415-416
15–16	Puberty, 246, 372
Precision, 30b	Pubic hair, 246, 384 <i>t</i> –385 <i>t</i> ,
Precocious puberty, 372	386 <i>t</i> –387 <i>t</i>
Predictive value, 31 <i>b</i> –32 <i>b</i>	Pulmonary embolism, 155t, 158t
Preeclampsia, 395b	Pulmonary fibrosis, idiopathic,
Pregnancy, 389-402	155 <i>t</i>
common concerns of, 389-391	Pulmonary function, 150
ectopic, 196, 400	Pulmonary hypertension, 364
examination in, 394-402	Pulmonary tuberculosis, 157t
health promotion and counseling	Pulmonary valve
in, 391–393	atresia of, 363, 382t
planning or preventing, 245	stenosis of, 170, 363, 381 <i>t</i>
recording exam findings in, 402b	Pulse pressure, 414
symptoms of, 242, 389	Pulses
Prehypertension, 63 <i>b</i> , 164, 414	apical, 64, 168, 170
Premature ejaculation, 230	brachial, 219
Prenatal care. See Pregnancy	carotid, 169
Prepatellar bursitis, 292, 308t	dorsalis pedis, 220
Presbyopia, 112-113, 416	femoral, 220
Present illness, 2b, 3-4, 350	in infants, 358, 363–364
Presentation, fetal, 401	in older adults, 420
Pressure sores, 415	palpation and grading of,
Preventive care, pediatric, 379t	218–219, 220–221
Primum non nocere (first, do no	paradoxical, 174
harm), 53	popliteal, 220
Privacy, 10, 38	posterior tibial, 221
Problem identification, 17b–19b, 36	pulsus alternans, 174
Pronator drift, 324	*
Prostate gland	radial, 64, 168, 219
benign hyperplasia of, 260, 264,	Pulsus alternans, 174
267t, 268t, 420	Pupils, 117, 318, 333, 347t
cancer of, 260–261, 269t, 420	Purging, 70t
examination of, 263-264	Purpura, 103t
inflammation of (prostatitis), 198,	Pustules, 99 <i>t</i> –100 <i>t</i>
264, 269t	Pyelonephritis, 198
Prostate-specific antigen (PSA) test,	Pyloric stenosis, 364
261	Pyrexia (fever), 56, 65–66, 198
Prostatitis, 198, 269t	
Proteinuria, 395b	Q
Pseudoclaudication, 301t	
Pseudoscars, 415	Quadriceps, 291, 292
Psoas sign, 207–208	Questions, interview
Psoriasis, 116	adaptive questioning, 43–44
Psychiatric disorders. See Mental	directed, 41, 43
status	open-ended, 40, 41
Psychopathologic processes, 18b	for sensitive topics, 49–53

R	grading, 328b
Radial nerve, 322	in infants, 366
Radial pulse, 64, 168, 219	and motor disorders, 340t
Radiculopathy, 301t, 303t, 322, 331	in pregnancy, 400
Rales, 147 <i>b</i>	Regional enteritis, 210t
Range of motion (ROM), 13	Reliability (test), 30b
at ankle, 297-298	Renal artery disease, 217
at hip, 290–291	Renal system, 193 <i>b</i>
limitations to, 275	Resonance, 145 <i>b</i>
measuring, 299	Respiratory alkalosis, 396
at neck and spine, 289	Respiratory infections, 361, 362
in older adults, 420–421	Respiratory rate and rhythm
at shoulder, 279–280	abnormalities of, 74t
at wrists and hands, 285–287	assessment of, 64–65, 143
Rapport with patient, 39, 367 <i>b</i>	in children, 358, 361, 362 <i>b</i>
Raynaud's phenomenon, 216	in older adults, 414
Reading ability, 48	Respiratory system
Reasoning, clinical, 17–19. See also	assessing function of, 150
Assessments	common or concerning symptoms
Reassurance, of patient, 45	of, 57, 141 <i>b</i>
Rebound tenderness, 202–203	disorders of, 153 <i>t</i> , 154 <i>t</i> –159 <i>t</i> ,
Records, patient. See also specific	161 <i>t</i> , 163
examinations	and dyspnea, 154 <i>t</i> –155 <i>t</i> , 163
checklist for, 33b–36b	examination of, 12, 143-150
for children, 375 <i>b</i> –377 <i>b</i>	health history of, 6, 142
of history and physical examina-	health promotion and counseling
tions, 33–36, 69	for, 142
for older adults, 422 <i>b</i> –423 <i>b</i>	in pregnancy, 396
	recording findings of, 151b
organizing, 33–36	Reticular activating system, 312
reviewing before exam, 38	Retina, 112, 119 <i>b</i> –120 <i>b</i>
Rectal temperature, 65, 66	Retinal arteries and veins, 112, 119,
Rectocele, 247, 256 <i>t</i> , 398	120b
Rectovaginal exam, 250, 400, 420	Retinopathy, diabetic, 133t
Rectum, 259–269	Review of systems, 3b, 5–7. See also
bleeding in, 260	specific systems
cancer of, 269t	Rheumatic fever, 275
examination of	Rheumatoid arthritis
abnormalities in, 268 <i>t</i> –269 <i>t</i>	in knees, 308t
in men, 15, 262–264	in older adults, 421
in older adults, 420	pain in, 305 <i>t</i> –306 <i>t</i>
in women, 15, 250, 264	symptoms of, 274–276
polyps of, 268t	in wrists or hands, 283, 284
Red reflex, 118	Rhinitis, 114
Referral, source of, 2b	Rhinorrhea, 114
Referred pain, 194	Rhonchi, 147b
Reflex esophagitis, 153t	Rigidity, 343 <i>t</i>
Reflexes. See also specific reflexes	Ringworm (tinea capitis), 108t
assessing, 14, 328-331	Rinne test, 122, 135 <i>t</i> , 320

Pombous test 224	Self-examinations
Romberg test, 324 Potator suff 280, 281, 207;	
Rotator cuff, 280–281, 307 <i>t</i>	breast, 182, 183, 187 <i>t</i> –188 <i>t</i>
Rovsing's sign, 207	testicular, 232, 234 <i>b</i> –235 <i>b</i> Self-reflection, 38
	Senile ptosis, 415–416
S	Sensations
	discriminative, 326–327
Sacroiliac joints, 288	loss of, 314
Safety, 5, 408	testing, 318, 325–326
Salpingitis, 213t	Sensitivity (test), 30 <i>b</i> –31 <i>b</i> , 32 <i>b</i>
Sarcoidosis, 155 <i>t</i>	Sensorineural hearing loss, 113, 135 <i>t</i> ,
Sarcoma, Kaposi's, 105t	319
Satiety, early, 195	Sensory system, 14, 325–327
Scabies, 100 <i>t</i>	Serial 7s, 83
Scales, skin, 100t	Sexual abuse, 373, 388 <i>b</i>
Scalp, 116	Sexual history and function, 49
Scaphoid fracture, 284	in females, 243
Scapula, winging of, 332	in males, 230–231
Scars, 101 <i>t</i>	Sexual maturity
Schizoaffective disorder, 92t	in boys, 384 <i>t</i> –385 <i>t</i>
Schizophrenia, 92t	in girls, 246, 383 <i>t</i> , 386 <i>t</i> –387 <i>t</i>
Schizophreniform disorder, 92t	Sexuality, in clinician-patient rela-
Sciatic nerve, 288	tionship, 53
Sciatica, 301 <i>t</i>	Sexually transmitted diseases
Sclera, 117	in children, 373
Scoliosis, 299, 375	counseling for, 231-232, 245,
Screening tests	263
for breast cancer, 182–183	in females, 243, 245
for cervical cancer, 244	in males, 231-232, 237t
for children, 351	Shaking chills, 56
for colorectal cancer, 200,	Shortness of breath, 142, 154 <i>t</i> –155 <i>t</i> ,
261–262, 409, 420	163
for depression, 57, 79–80, 411 <i>b</i>	Shoulders, 279–281, 307t
history of, 4	Sighing breathing, 74t
in older adults, 408–409,	Sigmoidoscopy, 261-262
410 <i>b</i> -412 <i>b</i> , 422 <i>b</i>	Signs, 17b. See also Symptoms
for prostate cancer, 261–262	Silent patient, 46
Scrotum	Sinuses, 6, 12, 114, 122
abnormalities of, 236t	Sinusitis, 126 <i>t</i> –127 <i>t</i>
in children, 365, 372	Skene's glands, 398
edema in, 236 <i>t</i>	Skin
examination of, 15, 233	cancer of, 93, 104 <i>t</i> –105 <i>t</i> , 106 <i>t</i> ,
hernia in, 236t	409,415
in older adults, 420	color of, 57, 94, 96t, 362b
Seborrheic keratoses, 104 <i>t</i> –105 <i>t</i> ,	common or concerning symptoms
415	of, 93 <i>b</i>
Seizures, 315	dimpling of, 191t
Self-awareness, 52–53	examination of, 12, 94, 116
Self-esteem, low, 80	health history of, 5

health promotion and counseling	Spontaneous pneumothorax, 155t
for, 93 <i>b</i>	Sprains, 275, 297–298, 309t
in infants, 358	Squamous cell carcinoma, 104t,
lesions of, 57, 94, 97t–104t	105 <i>t</i> , 415
nevi (moles), 93, 106t	Stance, 289, 324
in older adults, 415	Standard precautions, 15-16
recording exam results of, 95b	Staphylococcus aureus, methicillin-
tags, 365	resistant, 15–16
tumors of, 104 <i>t</i> –105 <i>t</i>	Stereognosis, 326-327
Skull, 116	Sternomastoid muscles, 143, 320
Sleep disorders, 80	Stiffness, joint, 275, 302t
Smell, sense of, 318	Stomach, 142
Smoking	Stools, 197, 260
counseling for, 142	Strabismus, 360, 368
history of, $2b$, 4	Strep throat, 114
in older adults, 407	Stress disorders, 91t
during pregnancy, 392	Stridor, 144, 362b
Smooth tongue, 137t	Stroke
Sneezing, 114	facial paralysis in, 339t
Social development, 350	health promotion and counseling
Social history, 3 <i>b</i> , 5	for, 165, 315–316
Social phobia, 90t	symptoms of, 313, 322, 338t
Somatoform disorders, 46, 87t	types of, 336 <i>t</i> –337 <i>t</i>
Spasmodic torticollis, 342t	vascular territories of, 338t
Spastic diplegias, 366	Stupor, 332–334
Spasticity, 343t	Sty, 131 <i>t</i>
Specificity (test), 31 b, 32 b	Subacromial bursa, 307t
Speculum examination, 247–248,	Subarachnoid hemorrhage, 112,
399, 419	127 <i>t</i> , 313
Speech	Subdeltoid bursitis, 307t
aphasia, 82	Subdural hematoma, 357
in delirium and dementia, 424 <i>t</i>	Subjective data, 2b
development of, in children, 350	Substance abuse, 2b, 4, 50–51, 80,
and mental status, 78, 82	199
in older adults, 409	Subtalar joint, 298
Spelling backwards, 83	Suicide, 80
Spermatic cord, 233, 239t, 372	Summarization, 45
Spermatocele, 239 <i>t</i>	Sun exposure, 93
Spheroidal (ball and socket) joints, 273 <i>b</i>	Supinator reflex, 329
	Surgical history, 4
Spider angioma, 103 <i>t</i> Spider veins, 103 <i>t</i>	Surrogate informant, 47
Spinal accessory nerve, 317 <i>b</i> , 320	Sutures, cranial, 357, 358–359
Spinal cord, 312	Swallowing, 196, 320
Spinal nerve, 312	Sweating, 56, 115
Spinal stenosis, 215, 301 <i>t</i>	Swelling
Spine, 12, 287–289	of feet or legs, 216
Spinous processes, 288	in infant head, 359
Spleen, 14, 204	in joints, 275

Tendonitis, 275, 307t

Symptoms. See also specific structures	Tendons, 271
or systems	Tenosynovitis, 275
characterization of, 4	Tension headaches, 111, 125 <i>t</i>
eliciting, 41, 404–405	Terry's nails, 109t
identifying, 17b	Test selection, 30–32
multiple, 46	Testes
sequence and time course of,	abnormalities of, 238t
28–29, 41, 56	in children and adolescents, 372,
seven attributes of, 41 b	384 <i>t</i> –385 <i>t</i>
unexplained, 75 <i>b</i> –76 <i>b</i> , 87 <i>t</i>	examination of, 233
Syncope, 314	in older adults, 420
Syndactyly, 365	self-examination of, 232,
Synovial joints, 272b, 273b	234 <i>b</i> –235 <i>b</i>
Syphilis	tumor of, $238t$
lip chancre in, 136t	Tetralogy of Fallot, 381 <i>t</i> –382 <i>t</i>
mucous patch of, on tongue, 138t	Thalamus, 311
penile chancre in, 232, 237t	Thermometers, 65–66
secondary, 252t	Thoracic kyphoscoliosis, 160t
vulvar chancre in, 252t	Thorax (chest), 141–161
Systemic lupus erythematosus, 276	anterior, 13, 149–150
Systems review, 3b, 5–7. See also spe-	common or concerning symptoms
cific systems	of, 141 <i>b</i>
Systolic blood pressure, 63 <i>b</i> , 168,	disorders of, 152 <i>t</i> –161 <i>t</i>
413–414	examination of, 12, 13, 143–150
Systolic clicks, 171, 176t	health history of, 141–142
	in newborns, 361–363
т	in older adults, 417
	pain or discomfort in, 141–142,
Tachycardia, 358, 364	152 <i>t</i> –153 <i>t</i> , 163–164
Tachypnea, 74t	posterior, 12, 144–148
Tactile fremitus, 145	in pregnancy, 396
Talkative patient, 47	recording findings of, 151 <i>b</i>
Talocalcaneal joint, 298	shape of, 143, 159 <i>t</i> –160 <i>t</i>
Tavistock Principles, 54 <i>b</i>	Thought content, 78b
Teeth	abstract, 84
in children, 361, 371	assessment of, 79, 82
examination of, 122	in delirium and dementia, 425 <i>t</i>
health promotion for, 115	Thought processes, 78b
Telangiectasia, hereditary hemor-	assessment of, 82
rhagic, 136t	in delirium and dementia, 409, 425 <i>t</i>
Temperature	Thoughts, recurring or unusual, 77,
intolerance to, 115	79, 82
measuring, 65–66	Thrills, 170, 364
in older adults, 414	Throat
sensitivity to, 325, 326	abnormalities of, 139 <i>t</i>
Temporamandibular joint (TMI) 278	in children, 361, 370–371
Temporomandibular joint (TMJ), 278	examination of, 12, 123, 124 <i>b</i>
Tenderness, 202–203, 213 <i>t</i>	health history of, 6, 114–115

sore, 114, 139t

Thromboangiitis obliterans, 219	Tumors. See also Cancer
Thrush, 361	androgen-producing, 419
Thumbs, 285-286, 322	brain, 112, 127 <i>t</i> -128 <i>t</i>
Thyroglossal duct cysts, 361	estrogen-producing, 419
Thyroid gland, 115, 123–124, 140t,	myoma of uterus, 257t
396	skin, 104 <i>t</i> –105 <i>t</i>
TIA (transient ischemic attack), 313,	of testis, 238t
315–316	Turgor, skin, 358
Tibeofemoral joint, 292	Two-point discrimination, 327
Tibial pulse, posterior, 221	Tympanic membrane, 66, 121, 134 <i>t</i> ,
Tibial torsion, 366	369–370
Tibiotalar joint, 297	Tympanosclerosis, 134t
Tics, 342t	Tympany, 145 <i>b</i>
Tinea capitis (ringworm), 108t	
Tinel's sign, 287	
Tinnitus, 114	U
Toenails, 95	Ulcerative colitis, 210t
Tongue, 123, 137 <i>t</i> -138 <i>t</i> , 320, 361	Ulcers
Tonsils, 123, 371	aphthous, 138t
Torsion, spermatic cord, 239t, 372	of feet and ankles, 225t
Torticollis, spasmodic, 342t	on lower legs, 216
Tortuous atherosclerotic aorta, 417	neuropathic, 225t
Total anomalous pulmonary venous	skin, 102 <i>t</i>
return, 363	venous stasis, 216
Touch sensation, 325, 326	Ulnar nerve, 322
Trachea, 123, 141, 161t	Umbilical artery, 355
Tracheal breath sounds, 147b	Umbilical cord, 355
Tracheitis, bacterial, 362b	Umbilical hernias, 355
Tracheobronchitis, 153t, 156t	Universal precautions, 16
Transient ischemic attack, 313,	Urethra, 247
315–316	caruncle of, 247, 419
Transmitted voice sounds, 148, 150, $161t$	infection of (urethritis), 198, 231, 232, 247
Transposition of great arteries, 363,	Urgency, urinary, 198
382 <i>t</i>	Urinary incontinence
Transverse tarsal joints, 298	in older adults, 410, 411b, 422b
Trapezius muscles, 320	types of, 199, 211 <i>t</i> –212 <i>t</i>
Tremors, 341t, 421	Urinary system, 198–199
Triceps reflex, 328	common or concerning symptoms
Trichomonas vaginitis, 255t	of, 193 <i>b</i>
Trichotillomania, 107t	health history of, 6, 198-199
Tricuspid regurgitation, 168	Urination, 198-199, 260
Trigeminal nerve, 317b, 318-319	Urine
Trigeminal neuralgia, 128t	blood in, 198
Trochanteric bursa, 289, 290f	color of, 197
Trochlear nerve, 317b, 318	protein in, 395b
True negatives or positives, 30b, 31b	Uterus
Trunk, 323	bicornuate, 400
Tuberculosis, pulmonary, 157t	contractility of, 397

Uterus (continued)	Venous pulsations, 120b
examination of, 15, 249	Venous stasis ulcers, 216
myoma of, 257t	Ventricular failure, left, 154t, 158t,
positions of, 257t	161 <i>t</i> , 163
in pregnancy, 390, 397, 399–400	Ventricular hypertrophy, left, 417
prolapsed, 256t	Ventricular septal defect, 382t
	Vertigo, 114, 313
	Vesicles, 99t
V	Vesicular breath sounds, 147b
Vagina	Vibration sense, 326
adenosis of, 254t	Violence, domestic, 51, 392-393
bleeding from, 373	Viral infections, 156t, 209t
discharge from, 243, 255t, 373	Visceral pain or tenderness,
examination of, 15, 249	193–194, 213 <i>t</i>
infection of (vaginosis), 255t	Vision
inflammation (vaginitis), 255t, 373	acuity of, 116, 318, 368-369, 416
walls of, during pregnancy, 399	changes in, 112-113
Vagus nerve, 317b, 320	in children, 360, 368-369
Valgus stress test, 296	disorders of, 115
Validation, of patient's feeling, 45	double vision, 113, 313
Validity (test), 30b	examination of, 12, 116, 360,
Valsalva maneuver, 174	368–369, 416
Values, defined, 52	field defects, 129t
Varicocele, 239t	and headaches, 112, 126t
Varicose veins, 138 <i>t</i> , 221, 400	health history of, 112–113
Varus stress test, 294	health promotion and counseling
Vascular disease, 363	for, 115
Vascular ring, 362b	interviewing patient with impair-
Vascular system, peripheral, 215–227	ment of, 49
abnormalities of, 224t-225t	in older adults, 115, 408,
common or concerning symptoms	410 <i>b</i> -411 <i>b</i> , 416, 422 <i>b</i>
of, 215 <i>b</i>	Vital signs
examination of, 13, 218–223,	blood pressure, 61–64
226 <i>t</i> –227 <i>t</i>	heart rate and rhythm, 64
health history of, 7, 215–216	measuring, 11, 61–66
health promotion and counseling	in newborns, 357
for, 216–218	in older adults, 413–414, 422 <i>b</i>
in older adults, 420	in pregnancy, 394–395
recording exam results of, 223 <i>b</i>	recording findings of, 69, 376b
Vasodepressor syncope, 314	respiratory rate and rhythm,
Vasomotor rhinitis, 114	64–65
Vasovagal syncope, 314	"sixth" (functional status), 410–413
Veins	temperature, 65–66
jugular, 168	Vitamin D, sources of, 72t
retinal, 112, 119, 120 <i>b</i>	Vitreous floaters, 113
spider, 103 <i>t</i>	Vitreous hemorrhage, 112
varicose, 138 <i>t</i> , 221, 400	Vocabulary assessment, 84
Venereal warts, 232, 237t, 251t, 260	Voice sounds, 148, 150, 161 <i>t</i> , 320
Venous insufficiency, 224t, 225t	Vomiting, 112, 195, 395

Vulva, 251*t*–252*t* Vulvovaginitis, 198



Waist circumference, 61
Warts, venereal, 232, 237*t*, 251*t*, 260
Weakness, muscle, 55, 313–314, 321–322
"Weaver's bottom," 290
Weber test, 122, 135*t*, 319
Weight
at birth, 353, 354*b*

and body mass index, 59-61

changes in, 55
of children and adolescents, 356, 368, 378t
general survey of, 59
health counseling for, 56–57
in older adults, 411b, 414
during pregnancy, 391, 392b, 395
Wheals, 98t
Wheezing, 142, 144, 147b
Whiplash, 303t
Whispered pectoriloquy, 148b
White lines in nails, 109t
White matter, 311
Wrists, 283–287, 322